



# HISTOPATHOLOGY OF THE SKIN

WALTER F LEVER, M D

*Clinical Associate in Dermatology, Harvard  
Medical School Assistant Dermatologist  
Massachusetts General Hospital Associate in  
Dermatology, Peter Bent Brigham Hospital  
Consulting Dermatologist Massachusetts Eye  
and Ear Infirmary, The Children's Hospital  
and Robert Breck Brigham Hospital*

SECOND EDITION

281 Illustrations

With 8 in Color

LONDON

PITMAN MEDICAL PUBLISHING CO LTD

*First Published in the United States of America in 1924 by*  
**J B LIPPINCOTT COMPANY**

*Published in Great Britain by*  
**PITMAN MEDICAL PUBLISHING CO LIMITED**  
**45 NEW OXFORD STREET LONDON W C 1**

**ASSOCIATED COMPANIES**

**SIR ISAAC PITMAN & SONS LTD**  
**PITMAN HOUSE PARKER STREET KINGSWAY LONDON W C 2**  
**THE PITMAN PRESS BATH**  
**PITMAN HOUSE LITTLE COLLINS STREET MELBOURNE**  
**27 BECKETTS BUILDING PRESIDENT STREET JOHANNESBURG**

**PITMAN PUBLISHING CORPORATION**  
**2 WEST 45TH STREET NEW YORK**

**SIR ISAAC PITMAN & SONS (CANADA) LTD**  
**(INCORPORATING THE COMMERCIAL TEXT BOOK COMPANY)**  
**PITMAN HOUSE 381 383 CHURCH STREET TORONTO**

In Memory of My Father  
DR ALEXANDER LEVER  
1877-1916

My First Teacher in Dermatology





# Preface to the Second Edition

Descriptive histopathology is regarded by some as a static science in which nowadays but few concepts change. But this is not so. Many advances have been made in the field of dermatopathology in only the five years that have passed since the appearance of the First Edition of this book. Consequently many changes were necessary for this Second Edition.

Important advances have been made in recent years in the histologic diagnosis of the vesicular and the bullous diseases especially of pemphigus. Therefore the descriptions of these diseases have been entirely rewritten and brought together into a new separate chapter. Because of changes in the concept of the histogenesis of the nevus cell the chapter on nevi and melanomas also has been rewritten. The use of the periodic acid Schiff reaction for the demonstration of fungi in tissue is an important advance which is taken due notice of in the chapter on fungus diseases. Extensive changes also have been made for instance in the descriptions of kraurosis vulvae the purpuras and verruca. Furthermore discussion of the following diseases has been incorporated because recent work has increased the interest of dermatologists in them: beryllium granuloma, papular myxedema, porphyria, ochronosis, hibernoma and hemangiopericytoma. Many changes have been made in the Bibliographies that follow each chapter in order to keep them up to date. Fifty six new photomicrographs have been added and 14 of the old photomicrographs have been replaced by better ones.

It is with great regret that I record the death of Dr. Tracy B. Mallory, former chief of the Pathology Laboratory at the Massachusetts General Hospital. To him I owe a great debt of gratitude. Dr. Benjamin Castleman, the present chief of the laboratory, has advised and helped me in many ways and I thank him. I also wish to express my thanks to Mr. Richard W. St. Clair for producing with great skill the new photomicrographs and to Mr. Walter Kahoe, Director of the Medical Department of J. B. Lippincott Company for his many courtesies and his unfailing co-operation.

WALTER F. LEVER



# Preface to the First Edition

This book is based on the courses of dermatopathology which I have been giving in recent years to graduate students of dermatology enrolled at Harvard Medical School and Massachusetts General Hospital. The book is written primarily for dermatologists. I hope however that it may be useful also to pathologists since dermatopathology is given little consideration in most textbooks of pathology.

I have attempted to keep this book short. Emphasis has been placed on the essential histologic features. Minor details and rare aberrations from the typical histologic picture have been omitted. I have allotted more space to the cutaneous diseases in which histologic examination is of diagnostic value than to those in which the histologic picture is not characteristic. In spite of my striving for brevity I have discussed the histogenesis of several dermatoses because knowledge of the histogenesis often is of great value for the understanding of the pathologic process.

Primarily for the benefit of pathologists who usually are not too familiar with dermatologic diseases I have preceded the histologic discussion of each disease with a short description of the clinical features.

A fairly extensive bibliography has been supplied for readers who are interested in obtaining additional information. In the selection of articles for the bibliography preference has been given whenever possible to those written in English.

I wish to express my deep gratitude to Dr. Tracy B. Mallory and Dr. Benjamin Castleman of the Pathology Laboratory at the Massachusetts General Hospital for the training in pathology they have given me. It has been invaluable to me. Their teaching is reflected in this book. Furthermore I wish to thank Mr. Richard W. St. Clair who with great skill and patience produced all the photomicrographs in this book.

WALTER F. LEVER



# Contents

1	INTRODUCTION	1
	Technic for Biopsy	1
	Limitations of Histologic Diagnosis	1
2	EMBRYOLOGY OF THE SKIN	3
	The Epidermis	3
	The Dermis	5
3	HISTOLOGY OF THE SKIN	7
	The Epidermis	7
	The Epidermal Appendages	13
	The Dermis	19
4	LABORATORY METHODS	28
	Fixation Embedding and Staining	28
	Histochemical Staining	30
	Polariscopic Examination	31
5	MORPHOLOGY OF THE MESODERMAL CELLS	32
	Myeloid Group	32
	Lymphoid Group	33
	Reticular or Histiocytic Group	34
	Plasma Cells	36
	Mast Cells	37
6	CONGENITAL DISEASES (GENODERMATOSES)	40
	Ichthyosis	40
	Congenital Ichthyosiform Erythroderma	41
	Keratosis Palmaris et Plantaris	42
	Keratosis Punctata Palmaris et Plantaris	42
	Pachyonychia Congenita	42
	Porokeratosis Mibelli	43
	Xeroderma Pigmentosum	44
	Congenital Ectodermal Defect	44
	Rothmund's Syndrome Werner's Syndrome and Progeria of Children	45
	Hydroa Vacciniiforme and Hydroa Aestivale	46
	Epidermolysis Bullosa	47
	Darier's Disease (Keratosis Follicularis)	48

6	CONGENITAL DISEASES (GENODERMATOSES)—( <i>Continued</i> )	
	Familial Benign Chronic Pemphigus (Hailey and Hailey)	51
	Epidermodysplasia Verruciformis (Lewandowsky and Lutz)	52
	Acrokeratosis Verruciformis (Hopl)	53
	Pseudoxanthoma Elasticum	54
	Cutis Hyperelastica (Ehlers Danlos Syndrome)	56
	Urticaria Pigmentosa	57
	Incontinentia Pigmenti	59
7	NONINFECTIOUS VESICULAR AND BULLOUS DISEASES	64
	Dermatitis Eczema	66
	Dermatopathic Lymphadenitis	75
	Miliaria	76
	Pompholyx (Dyshidrotic Eruption)	76
	Pemphigus	77
	Pemphigus Vulgaris	77
	Pemphigus Vegetans	80
	Pemphigus Foliaceus	82
	Pemphigus Erythematosus	84
	Bullous Pemphigoid	85
	Benign Mucous Membrane Pemphigoid (Pemphigus Con-	
	junctivae)	87
	Dermatitis Herpetiformis	88
	Erythema Multiforme	89
	Burn	90
8	NONINFECTIOUS INFLAMMATORY DISEASES	94
	Urticaria	94
	Prurigo Nodularis	95
	Erythema Nodosum	95
	Nodular Vasculitis	97
	Psoriasis	98
	Pustulosis Palmaris et Plantaris (Pustular Psoriasis Pus-	
	tular Bacterid)	101
	Acrodermatitis Continua (Hallopeau)	102
	Impetigo Herpetiformis	104
	Parapsoriasis	104
	Pityriasis Rosea	105
	Lichen Planus	106
	Lichen Nitidus	110
	Lichen Striatus	111
	Pityriasis Rubra Pilaris	111

8	NONINFECTIOUS INFLAMMATORY DISEASES—(Continued)	
	Granuloma Faciale (Eosinophilic Granuloma of the Face)	111
	Erythema Elevatum Diutinum	114
	Granuloma Annulare	114
	Subcutaneous Nodules of Rheumatic Fever and Rheumatoid Arthritis	117
	Acrodermatitis Chronica Atrophicans	118
	Radiodermatitis	119
	Solar Dermatitis	122
	Hydrocystoma	123
	Acne Vulgaris	123
	Acne Varioliformis (Acne Necrotica Miliaris)	124
	Acne Rosacea	124
	Fox Fordyce Disease	125
	Alopecia Areata	125
	Alopecia Cicatricata (Pseudopelade Brocq)	126
	Purpura	126
	Noninflammatory Purpura	127
	Inflammatory Purpura (Vasculitis)	127
	Purpura Pigmentosa Progressiva (Majocchi Schamberg)	129
	Relapsing Febrile Nodular Nonsuppurative Panniculitis (Weber Christian Disease)	131
	Sclerema Neonatorum	133
	Hereditary Edema of the Legs (Milroy's Disease)	135
	Chondrodermatitis Nodularis Chronica Helicis	135
	Scabies	136
	Insect Bites	137
	Lichen Urticatus (Papular Urticaria)	138
	Foreign Body Granulomas	139
	Lipid Granuloma (Paraffinoma)	140
	Tattoo Granuloma	140
	Silicon Granuloma	140
	Beryllium Granuloma	141
	Swimming Pool Granuloma	142
9	ERUPTIONS DUE TO DRUGS	150
	Fixed Drug Eruption	150
	Quinacrine Hydrochloride (Atabrine) Dermatitis	150
	Exfoliative Dermatitis Due to Drugs	151
	Bromoderma	151
	Iododerma	151
	Argyria	152
	Chrysisis	153
		154



6	CONGENITAL DISEASES (GENODERMATOSES)—(Continued)	
	Familial Benign Chronic Pemphigus (Hailey and Hailey)	51
	Epidermodysplasia Verruciformis (Lewandowsky and Lutz)	52
	Acrokeratosis Verruciformis (Hopl)	53
	Pseudoxanthoma Elasticum	54
	Cutis Hyperelastica (Ehlers Danlos Syndrome)	56
	Urticaria Pigmentosa	57
	Incontinentia Pigmenti	59
7	NONINFECTIOUS VESICULAR AND BULLOUS DISEASES	61
	Dermatitis Eczema	66
	Dermatopathic Lymphadenitis	75
	Miliaria	76
	Pompholyx (Dyshidrotic Eruption)	76
	Pemphigus	77
	Pemphigus Vulgaris	77
	Pemphigus Vegetans	80
	Pemphigus Foliaceus	82
	Pemphigus Erythematosus	84
	Bullous Pemphigoid	85
	Benign Mucous Membrane Pemphigoid (Pemphigus Con-	
	junctivae)	87
	Dermatitis Herpetiformis	88
	Erythema Multiforme	89
	Burn	90
8	NONINFECTIOUS INFLAMMATORY DISEASES	94
	Urticaria	94
	Prurigo Nodularis	95
	Erythema Nodosum	95
	Nodular Vasculitis	97
	Psoriasis	98
	Pustulosis Palmaris et Plantaris (Pustular Psoriasis Pus-	
	tular Bacterid)	101
	Acrodermatitis Continua (Hallopeau)	102
	Impetigo Herpetiformis	104
	Parapsoriasis	104
	Pityriasis Rosea	105
	Lichen Planus	106
	Lichen Nitidus	110
	Lichen Striatus	111
	Pityriasis Rubra Pilaris	111

FUNGUS DISEASES—(Continued)	
Moniliasis	220
North American Blastomycosis (Gilchrist's Disease)	222
Torulosis (Cryptococcosis, European Blastomycosis of Busse and Buschke)	224
Chromoblastomycosis	225
Coccidioidomycosis (San Joaquin Valley Fever)	226
Actinomycosis	228
Sporotrichosis	229
Histoplasmosis	231
4 DISEASES CAUSED BY PROTOZOA	235
Leishmaniasis	235✓
5 DISEASES CAUSED BY VIRUSES	238
Variola Varicella, Herpes Zoster, Herpes Simplex	238
Eczema Vaccinatum and Kaposi's Varicelliform Eruption	244
Molluscum Contagiosum	244
Verruca	246
Verruca Vulgaris	247
Verruca Plana	249
Verruca Plantaris	251
Condyloma Acuminatum	251
Lymphogranuloma Venereum (Lymphogranuloma In guinale Lymphopathia Venerea)	252
Milkers Nodules	253
16 METABOLIC DISEASES	257
Lipoidoses	257
Lipoidoses with Increased Blood Lipids	257
Lipoidoses with Normal Blood Lipids	262
Localized Lipoidoses	269
Amyloidosis	272
Primary Systemic Amyloidosis	272
Primary Localized Amyloidosis of the Skin	274✓
Secondary Systemic Amyloidosis	275
Calcinosis Cutis	275
Metastatic Calcification	275
Metabolic Calcification	276
Gout	277
Porphyrria	278
Myxedema	279
Generalized Myxedema	279

9	ERUPTIONS DUE TO DRUGS—( <i>Continued</i> )	
	Arsenical Keratosis and Carcinoma	155
10	DEGENERATIVE DISEASES	157
	Senile Degeneration	157
	Lichen Sclerosus et Atrophicus	158
	Kraurosis Vulvae	161
	Balanitis Xerotica Obliterans	162
	Striae Distensae	162
	Macular Atrophy (Anetoderma)	163
	Atrophoderma Reticulatum (Folliculitis Ulerythematosi Reticulata)	163
	Colloid Milium (Colloid Degeneration of the Skin)	163
11	BACTERIAL DISEASES	167
	Impetigo	167
	Erysipelas	168
	Folliculitis (Impetigo Bockhart, Furuncle, Folliculitis Barbae)	168
	Folliculitis Keloidalis (Dermatitis Papillaris Capillitii)	169
	Hidradenitis Suppurativa	169
	Pyoderma Gangrenosum (Chronic Undermining Burrow ing Ulcer)	170
	Subacute Bacterial Endocarditis	170
	Meningococcemia (Waterhouse Friderichsen Syndrome)	171
	Keratitis Blepharorrhagica (Gonorrheal Keratitis)	172
	Tuberculosis	174
	Primary Tuberculosis	176
	Reinfection Tuberculosis	177
	Sarcoidosis	188
	Leprosy	193
	Diphtheria of the Skin	197
	Anthrax	198
	Tularemia	199
	Chancroid	199
	Granuloma Inguinale	203
	Rhinoscleroma	202
12	SPIROCHETAL DISEASES	208
	Syphilis	208
13	FUNGUS DISEASES	217
	Dermatophytosis (Tinea)	217

## 1 EPIDERMAL TUMORS—(Continued)

## 2 Tumors of the Epidermal Appendages—(Continued)

Hair Nevus	346
Sebaceous Adenoma	347
Syringocystadenoma Papilliferum (Nevus Syringo- cystadenomatosus Papilliferus)	349
Hidradenoma Papilliferum	352
Sebaceous Epithelioma	353
Syringoma	354
Cylindroma	357
Myo-epithelioma (Myo-epithelial Sweat Gland Tu- mor)	359
Trichoepithelioma (Epithelioma Adenoides Cysti- cum Multiple Benign Cystic Epithelioma)	363
Calcifying Epithelioma (Mallherbe)	366
Basal Cell Epithelioma	369
Basal Cell Papilloma (Verruca Senilis Keratosis Se- borrheica)	382
Carcinoma of Sebaceous Glands	385
Carcinoma of Eccrine Sweat Glands	386
Carcinoma of Apocrine Glands	389
Metastatic Carcinoma of the Skin	389

## 19 MESODERMAL TUMORS

1 Tumors of Fibrous Tissue	403
Dermatofibroma Histiocytoma (Sclerosing Heman- gioma Nodular Subepidermal Fibrosis)	403
Keloid	408
Fibrosarcoma	408
2 Tumors of Mucoid Tissue	411
Myxoma	411
Myxosarcoma	412
Synovial Cyst of the Skin (Myxomatous Degenera- tion Cyst)	413
3 Tumors of Fatty Tissue	413
Lipoma	413
Hibernoma	414
Liposarcoma	414
Malignant Hibernoma	415
4 Tumors of Nerve Tissue	416
Neuroma	416
Neurofibromatosis (von Recklinghausen's Disease)	417
Neurolemoma	421

16	METABOLIC DISEASES—(Continued)	
	Myxedema—(Continued)	
	Circumscribed Myxedema	279
	Papular Myxedema (Papular Mucinosiſ)	281
	Scleredema Adultorum (Buschke)	281
	Addison's Disease	281
	Acanthoſiſ Nigricans	282
	Riehl's Melanoſiſ, Melanodermatitiſ Toxică, Poikiloderma Reticulare (Civatte)	283
	Vitiligo	285
	Hemochromatotiſ (Bronze Diabetes)	285
	Ochronoſiſ	286
	Vitamin A Deficiency (Phrynoderma)	286
	Pellagra	287
	Vitamin C Deficiency (Scurvy)	288
17	SYSTEMIC DISEASES OF UNKNOWN CAUSE	293
	Lupus Erythematoſuſ	293
	Lupus Erythematoſuſ Profunduſ (Kaposi Irring)	303
	Dermatomyoſitiſ	303
	Poikiloderma Atrophicuſ Vasculare (Jacobi)	305
	Scleroderma	307
	Periarteritiſ Nodosa	310
	Allergic Granulomatotiſ	312
18	EPIDERMAL TUMORS	317
	Histoſiſ and Claſſificatiſ of Epidermal Tumors	317
	1 Tumors of the Surface Epidermiſ	321
	Nevuſ Verrucoſuſ	321
	Epidermal Cyst (Wen), Sebaceous Cyst, Milium, Dermoid Cyst	324
	Keratoſiſ Seniliſ	326
	Leukoplakia	328
	Squamous Cell Carcinoma (Epidermoid Carcinoma)	329
	Bowen's Disease	336
	Paget's Disease	339
	2 Tumors of the Epidermal Appendages	343
	Nevuſ Sebaceuſ (Jadassohn)	343
	Adenoma Sebaceum (Pringle) (Tuberouſ Scleroſiſ)	344
	Senile Sebaceous Nevuſ (Senile Sebaceous Hyperplasia)	346
	Fordyce's Disease	346
	Apocrine Gland Nevuſ	346

# 1

## Introduction

### TECHNIC FOR BIOPSY

It is important to select a proper site for biopsy. In most instances histologic examination of a fully developed lesion will give more information than examination of an early or an involuting lesion. An exception to this rule represents vesicular, bullous and pustular lesions. For their histologic examination a very early lesion is required otherwise, secondary changes (such as regeneration, degeneration or secondary infection) may obscure essential features and make recognition of their mode of formation impossible. Generally, it is inadvisable to include normal tissue in the biopsy specimen unless a large specimen is taken or the physician personally supervises the processing of the specimen because improper sectioning by the technician may result in only normal skin being seen in the section. The specimen should include subcutaneous fat, because, in many dermatoses characteristic histologic features are found in the lower dermis or in the subcutaneous fat. If several types of lesions are present and the diagnosis hinges on the histologic findings much time may be saved by taking specimens for biopsy from more than one lesion.

In the author's experience a specimen obtained with a 6 mm biopsy punch nearly always has proved adequate for histologic study. Two sutures are sufficient to close the wound.

Before placing the specimen in the routine fixative, which in many hospitals is Zenker's or Helly's solution, one should consider which stains are desirable. Some of the special stains can be performed only if the specimen has been fixed in the appropriate fixative. A tabulation of staining methods with the required fixatives is found in Table 1 on page 29.

### LIMITATIONS OF HISTOLOGIC DIAGNOSIS

Although histologic study is one of the most valuable means of diagnosis in dermatology it has its limitations. Often no definitive diagnosis can be made. The reason for this is that few dermatoses, aside from the tumors, are associated regularly with a diagnostic his

## 19 MESODERMAL TUMORS—(Continued)

5 Tumors of Vascular Tissue	422
Hemangioma	422✓
Granuloma Pyogenicum	424✓
Angiokeratoma Mibelli Angiokeratoma Corporis Diffusum	426
Senile Hemangioma	426
Nevus Araneus	427
Osler's Disease (Familial Hemorrhagic Telangiectasia)	427
Lymphangioma	427
Glomus Tumor	429
Hemangiopericytoma	430
Kaposi's Sarcoma (Multiple Idiopathic Hemorrhagic Sarcoma)	431
Hemangioendothelioma (Hemangiosarcoma)	435
Post Mastectomy Lymphangiosarcoma	437
6 Tumors of Muscular Tissue	437
Leiomyoma	437✓
Granular Cell Myoblastoma	440
7 Tumors of Osseous Tissue	442
Osteoma Cutis	442

## 20 PIGMENTED NEVI AND MALIGNANT MELANOMA

Pigmented Nevus	450✓
Malignant Melanoma	458✓
Lentigo	464
Mongolian Spot	465
Blue Nevus	466✓

## 21 LYMPHOMA AND MYELOSIS

Lymphoma	470
1 Stem Cell Lymphoma	474
2 Reticulum Cell Lymphoma	474
3 Lymphoblastic Lymphoma	477
4 Lymphocytic Lymphoma	478✓
5 Follicular Lymphoma (Brill Symmers)	480
6 Hodgkin's Disease	481✓
7 Mycosis Fungoides	484
Lymphocytoma Cutis (Lymphadenosis Benigna Cutis Spiegler Fendt Sarcoid)	491
Myelosis	493

# Introduction

## TECHNIC FOR BIOPSY

It is important to select a proper site for biopsy. In most instances histologic examination of a fully developed lesion will give more information than examination of an early or an involuting lesion. An exception to this rule represents vesicular, bullous, and pustular lesions. For their histologic examination a very early lesion is required; otherwise secondary changes (such as regeneration, degeneration, or secondary infection) may obscure essential features and make recognition of their mode of formation impossible. Generally it is inadvisable to include normal tissue in the biopsy specimen unless a large specimen is taken or the physician personally supervises the processing of the specimen because improper sectioning by the technician may result in only normal skin being seen in the section. The specimen should include subcutaneous fat because in many dermatoses characteristic histologic features are found in the lower dermis or in the subcutaneous fat. If several types of lesions are present and the diagnosis hinges on the histologic findings, much time may be saved by taking specimens for biopsy from more than one lesion.

In the author's experience a specimen obtained with a 6 mm biopsy punch nearly always has proved adequate for histologic study. Two sutures are sufficient to close the wound.

Before placing the specimen in the routine fixative which in many hospitals is Zenker's or Helly's solution, one should consider which stains are desirable. Some of the special stains can be performed only if the specimen has been fixed in the appropriate fixative. A tabulation of staining methods with the required fixatives is found in Table 1 on page 29.

## LIMITATIONS OF HISTOLOGIC DIAGNOSIS

Although histologic study is one of the most valuable means of diagnosis in dermatology, it has its limitations. Often no definitive diagnosis can be made. The reason for this is that few dermatoses, aside from the tumors, are associated regularly with a diagnostic his-



19 MESODERMAL TUMORS—(*Continued*)

5 Tumors of Vascular Tissue	42
Hemangioma	42
Granuloma Pyogenicum	42
Angiokeratoma Mibelli, Angiokeratoma Corporis Diffusum	42
Senile Hemangioma	42
Nevus Araneus	42
Osler's Disease (Familial Hemorrhagic Telangiectasia)	42
Lymphangioma	42
Glomus Tumor	42
Hemangiopericytoma	43
Kaposi's Sarcoma (Multiple Idiopathic Hemorrhagic Sarcoma)	43
Hemangio endothelioma (Hemangiosarcoma)	43
Post Mastectomy Lymphangio sarcoma	43
6 Tumors of Muscular Tissue	43
Leiomyoma	43
Granular Cell Myoblastoma	44
7 Tumors of Osseous Tissue	44
Osteoma Cutis	44

## 20 PIGMENTED NEVI AND MALIGNANT MELANOMA

Pigmented Nevus	45
Malignant Melanoma	45
Lentigo	46
Mongolian Spot	46
Blue Nevus	46

## 21 LYMPHOMA AND MYELOSIS

Lymphoma	47
1 Stem Cell Lymphoma	47
2 Reticulum Cell Lymphoma	47
3 Lymphoblastic Lymphoma	47
4 Lymphocytic Lymphoma	47
5 Follicular Lymphoma (Brill Symmers)	18
6 Hodgkin's Disease	48
7 Mycosis Fungoides	48
Lymphocytoma Cutis (Lymphadenosis Benigna Cutis, Spiegler Fendt Sarcoid)	49

Myelosis	49
----------	----

# 2

## Embryology of the Skin

### THE EPIDERMIS

In the earliest period of fetal life the epidermis consists of a single layer of cells. Between the fifth and the seventh weeks of fetal life this becomes a double layer consisting of an inner layer the stratum




FIG 1 The skin of an embryo 4 months old. The epidermis consists of three layers: the stratum germinativum (SG), the stratum intermedium (SI), and the periderm (P). Two primary epithelial germs (E.G.) are shown. The fetal dermis shows many more fibroblasts than the adult dermis. (X400)

germinativum or palisade layer and an outside layer the periderm or epitrichial layer. The stratum germinativum is composed of large cuboidal cells while the periderm consists of flat cells. In the third month single cells appear between the two layers and later form a complete line the stratum intermedium (Fig 1). The cells of the

tologic picture. Instead, the histologic features may be merely suggestive of a diagnosis or may be entirely nonspecific. Even in the case of tumors, difficulties in diagnosis may arise. For instance, distinction of squamous cell carcinoma from pseudo epitheliomatous hyperplasia is not always possible. In cases of infectious granulomas such as syphilis, tuberculosis and the deep mycoses, a specific diagnosis often cannot be made unless the causative organism can be demonstrated. Great difficulties may also be encountered in the histologic study of the large group of noninfectious inflammatory dermatoses. In the diseases of this group, such as psoriasis, lichen planus and lupus erythematosus, in which the histologic picture is diagnostic as a rule, sometimes it may be merely suggestive. In other diseases of this group, such as the various types of dermatitis or eczema, the histologic picture is at best, only suggestive. In still others, such as pityriasis rosea and parapsoriasis, it is always nonspecific.

Nevertheless, frequently, when the histologic picture is not diagnostic, a correlation of the histologic with the clinical findings will make a diagnosis possible.

In many instances, the chief value of histopathologic study lies in corroborating the clinical diagnosis or in ruling out possible diseases which are being considered on the basis of clinical appearance. It is obvious that the histopathologist can give the clinician a maximum amount of information only if every specimen submitted for histologic diagnosis is accompanied by detailed clinical information, including a differential diagnosis.

except in a few areas where they persist (See The Apocrine Glands \* page 16)

**Melanocytes (Dendritic or Clear Cells)** Whereas in the past the melanin producing dendritic or clear cells were considered to be modified basal cells there is now almost complete agreement that they are neural cells which originate in the neural crest and migrate from there with the nerves to the epidermis during early fetal life (DuShane Rawles) Although this origin of the melanocytes has not been proved experimentally in man it has been proved in mice by Rawles who transplanted tissue from the neural crest of mouse embryos to the coelom of albino chicken embryos and observed the development of pigmented melanocytes

In human embryos argentaffine melanocytes first are identified within the epidermis early in the third month They possess long branching dendritic processes which in subsequent months become more numerous and longer up to 100 microns in length joining neighboring melanocytes (Zimmermann and Cornbleet) (For a further discussion of the melanocytes see pages 9 12)

## THE DERMIS

The dermis is of mesodermal origin During the first months of fetal life it consists of closely packed spindle shaped cells (mesenchymal cells) During the third month fibrils appear at first as a delicate anastomosing argyrophilic network (reticulum fibers) As the fibers increase in number and thickness they arrange themselves in bundles which no longer can be impregnated with silver and which instead begin to stain with the methods for collagenous fibers (Maximow and Bloom) Simultaneously the mesenchymal cells develop into fibroblasts The elastic fibers appear much later than the collagenous fibers usually in the sixth month (Lynch) The subcutaneous fat first becomes apparent in the third month

It is not as yet settled whether collagen and elastin develop intracellularly by a direct transformation of living substance of mesenchymal cells or extracellularly by precipitation of the ground substance under the influence of an enzyme derived from mesenchymal cells Observations on the development of fibers in tissue cultures favor the theory of their extracellular origin (Maximow and Bloom)

## BIBLIOGRAPHY


- DuShane C H The development of pigment cells in vertebrates In The  
 Fitcher " Acad Sci 1948  
 de " in the histogenesis of the epidermis

stratum intermedium are large and because of their clear cytoplasm have a ballooned appearance

At about the fourth month the periderm separates to aid in the formation of the vernix caseosa. At the same time the stratum intermedium becomes multilayered and develops into the squamous cell layer or stratum malpighii. Intercellular bridges become recognizable only after the epidermis has become stratified with several layers.

The embryonal stratum germinativum differentiates into the following types of cells (1) the basal cells (2) the eccrine sweat gland germ cells and (3) the primary epithelial germ cells (Chart 1)

CHART 1—EMBRYOLOGY OF THE EPIDERMIS



**Basal Cells** While the cells of the embryonal stratum germinativum possess no intercellular bridges the mature basal cells do. By progressive differentiation the basal cells develop into squamous cells granular cells and horny cells thus forming the multilayered surface epidermis.

**Epidermal Appendages** The epidermal appendages develop from the primary epithelial germ cells and the eccrine sweat gland germ cells. These cells possess no intercellular bridges and form cells without intercellular bridges. Primary epithelial germs first appear in the third month of fetal life as epithelial buds projecting into the dermis (Fig. 1). Eccrine sweat gland germs are first observed in the fifth or the sixth month.

two inner sheaths (the Huxley and the Henle layers) develop from the hair matrix. The outer sheath of the hair and the ducts of the sebaceous glands which are composed of prickle cells originate from cells with the potentiality of forming prickle cells namely from basal cells (Lever). The apocrine glands which begin to form with the hair and the sebaceous gland involute before they reach full development.

except in a few areas where they persist (See The Apocrine Glands \* page 16)

**Melanocytes (Dendritic or Clear Cells)** Whereas, in the past the melanin producing dendritic or clear cells were considered to be modified basal cells there is now almost complete agreement that they are neural cells which originate in the neural crest and migrate from there with the nerves to the epidermis during early fetal life (DuShane Rawles) Although this origin of the melanocytes has not been proved experimentally in man it has been proved in mice by Rawles who transplanted tissue from the neural crest of mouse embryos to the coelom of albino chicken embryos and observed the development of pigmented melanocytes

In human embryos argentaffine melanocytes first are identified within the epidermis early in the third month They possess long branching dendritic processes which in subsequent months become more numerous and longer, up to 100 microns in length joining neighboring melanocytes (Zimmermann and Cornbleet) (For a further discussion of the melanocytes see pages 9 12)

## THE DERMIS

The dermis is of mesodermal origin During the first months of fetal life it consists of closely packed spindle shaped cells (mesenchymal cells) During the third month fibrils appear at first as a delicate anastomosing argyrophilic network (reticulum fibers) As the fibers increase in number and thickness they arrange themselves in bundles which no longer can be impregnated with silver and which instead begin to stain with the methods for collagenous fibers (Maximow and Bloom) Simultaneously the mesenchymal cells develop into fibroblasts The elastic fibers appear much later than the collagenous fibers usually in the sixth month (Lynch) The subcutaneous fat first becomes apparent in the third month

It is not yet settled whether collagen and elastin develop intracellularly by a direct transformation of living substance of mesenchymal cells or extracellularly by precipitation of the ground substance under the influence of an enzyme derived from mesenchymal cells Observations on the development of fibers in tissue cultures favor the theory of their extracellular origin (Maximow and Bloom)

## BIBLIOGRAPHY

- DuShane G H The development of pigment cells in vertebrates In The Biology of Melanomas p 1 New York N Y Acad Sci 1948  
 Eichenlaub F J and Osbourn R L Studies in the histogenesis of the epidermis Arch Dermat & Syph 61 700 1951

## 6 Embryology of the Skin

- Lever W F Pathogenesis of benign tumors of the cutaneous appendages and of basal cell epithelioma Arch Dermat & Syph 57 679 1948
- Lynch F W Elastic tissue in fetal skin Arch Dermat & Syph 29 57 1934 (A good review of the embryology of the skin )
- Maximow A A and Bloom W A Textbook of Histology ed 4 pp 105 107 Philadelphia Saunders 1913
- Rawles M E Origin of pigment cells from the neural crest in the mouse embryo Physiol Zool 20 248 1947
- Zimmermann A A and Cornbleet T The development of epidermal pigmentation in the negro fetus J Invest Dermat 11 383 1948

## 3

## Histology of the Skin

## THE EPIDERMIS

In histologic sections of normal skin, the border between the epidermis and the dermis is irregular because of the fact that numerous cone shaped dermal papillae reach upward and indent the inner surface of the epidermis. The ridges of epidermis separating the papillae appear in histologic sections as pegs, and, therefore, often are referred to as rete pegs although the term rete ridges is preferable.

**Layers of the Epidermis.** The epidermis may be divided into four layers: (1) the *keratin*, (2) the *granular*, (3) the *spinous*, and (4) the *basal* layers.

located 1/2 mi. N. W. 1/4.

**U.S. E-Learning**

**THE BASAL LAYER** Two types of cells occur in the basal layer basal cells and melanocytes

Basal cells are columnar in shape and lie with their long axis vertical to the dividing line between the epidermis and the dermis. They have a deeply basophilic cytoplasm and a darkly staining oval or elongated nucleus. In sections stained with routine stains, the basal cells seem to contain melanin granules often concentrated at the

161

to

370

conclusion that the melanin granules were entirely within melano-  
cytes and that the

2) These bridges

~ ~ ~ 1/3 one third



## 6 Embryology of the Skin

- Lever, W. F. Pathogenesis of benign tumors of the cutaneous appendages and of  
Lync . . . . . 29-57, 1934 (A
- Maximow, A. A., and Bloom W. A textbook of histology, ed. 4 pp. 105-107  
Philadelphia: Saunders, 1943
- Rawles, M. E. Origin of pigment cells from the neural crest in the mouse em-  
bryo, *Physiol. Zool.* 20:218, 1947
- Zimmermann, A. A. and Cornbleet, F. The development of epidermal pig-  
mentation in the negro fetus. *J. Invest. Dermat.* 11:383, 1948

# 3

## Histology of the Skin

### THE EPIDERMIS

In histologic sections of normal skin the border between the epidermis and the dermis is irregular because of the fact that numerous cone shaped dermal papillae reach upward and indent the inner surface of the epidermis. The ridges of epidermis separating the papillae appear in histologic sections as pegs and therefore often are referred to as rete pegs although the term rete ridges is preferable.

**Layers of the Epidermis** The epidermis may be divided into four layers:

1. **Stratum lucidum**

2. **Stratum granulosum**

3. **Stratum spinosum** located between the granular and the horny layers. The cells in the various layers represent different stages in the gradual evolution of the basal cell into cornified cells and do not represent different types of cells.

**THE BASAL LAYER** Two types of cells occur in the basal layer: basal cells and melanocytes.

Basal cells are columnar in shape and lie with their long axis vertical to the dividing line between the epidermis and the dermis. They have a deeply basophilic cytoplasm and a darkly staining oval or elongated nucleus. In sections stained with routine stains the basal cells seem to contain melanin granules. Their nuclei are supranuclear and Montgomery, using silver

s a greater number of mitotic figures

epidermis to the dermis is accomplished by the firm interlocking of cytoplasmic processes of the basal cells with reticulum fibers located in the uppermost dermis. Whereas,




FIG. 2 Normal epidermis, dorsum of the hand. Four layers can be recognized (1) basal layer (2) stratum malpighii (3) granular layer and (4) horny layer. No stratum lucidum is present. Note the presence of intercellular bridges between the basal cells. Several clear cells (melanocytes) are present in the basal layer. They possess a small dark nucleus and clear cytoplasm. ( $\times 400$ )

in perpendicular sections these reticulum fibers appear like the bristles of a brush (Fig. 3), horizontal sections reveal them to form a continuous fibrillar meshwork around the cytoplasmic processes of the basal cells (Odland). In contrast with formerly expressed views, elastic fibers do not contribute to the coherence of the dermis and the epidermis since they do not extend high enough to reach the epidermis (Diek).

Whereas routine stains do not show a basement membrane, stains with the periodic acid Schiff stain after Hotchkiss and McManus (see page 30) show a thin homogeneous dense band (Fig. 1) at the dermal epidermal junction, indicating the presence of a relatively large amount of polysaccharide material in this zone (Stoughton and

Wells) This band represents a relative barrier to the diffusion of large particles as proved by the fact that dye injected intradermally beneath a subepidermal bulla will not appear in the bulla fluid and



FIG 3 (Top) for reticulum set work into which in a firm attach

FIG 4 (Bottom) Junction between the epidermis and dermis

(1820)

thorax

The  
ban

Melanocytes (Clear or Dendritic Cells) Melanocytes are of neural origin (see Embryology page 5) They stain with Bloch's dopa

stain (because they have the ability to form melanin) and usually also with silver stains (because they contain melanin) (see page 12) They also stain with gold In sections stained with hematoxylin and eosin melanocytes appear as clear cells having a small dark staining nucleus and a clear, slightly basophilic cytoplasm (Figs 2 and 254) On the other hand, in sections impregnated with silver they appear as dendritic cells with numerous long, branching processes provided a sufficient amount of melanin granules are present within the processes to show their outline (Ebert, Becker Jr, Fitzpatrick and Montgomery)

**THE STRATUM MALPIGHII** The cells of the stratum malpighii which are called squamous cells or prickle cells, are polygonal and form a mosaic They become flattened toward the surface (Fig 2) The cells are separated by spaces which are traversed by intercellular bridges or prickles They separate the squamous cells so that lymphatic fluid can circulate around them and supply them with nourishment The intercellular bridges are formed by the cytoplasm of adjoining cells and tonofibrils which extend through them from cell to cell As studies by phase contrast and polarization microscopy have shown, tonofibrils pass uninterruptedly from one cell to another all the way from the basal cell layer to the stratum corneum Roentgen spectography suggests that the tonofibrils consist of keratin (Nelmans Keuning van Rijssel and Ruiter)

Each of the intercellular bridges in the stratum malpighii contains a small nodular thickening called the nodule of Bizzozero These nodules are stained selectively by Heidenhain's iron hematoxylin which does not stain the epidermal cells and bridges (Tavre) This special staining reaction suggests that the nodules of Bizzozero are special structures and not just a thickening of the intercellular bridges Nieuwmeijer on the other hand believes that they merely represent an optical effect produced by the crossing of tonofibrils in the intercellular bridges

If a gold stain is employed one may observe interspersed between the cells of the stratum malpighii gold impregnated cells with den

so far about their nature Masson regards them as without melanocytes which are being carried away passively toward the surface Ferreira Marques points out that schwannian cells generally are auto

represent intra epidermal receptors of pain Becker Jr Fitzpatrick and Montgomery, on the other hand regard the Langerhans cells as ordinary melanocytes and maintain that they appear to lie in the

superficial epidermis only because of distortion and wrinkling of the tissue produced by the acid used in the gold impregnation. In separated epidermis, which is less distorted by acid than whole skin, they found gold impregnated cells only at the junction of the epidermis and the dermis and never in the upper layers of the epidermis.

THE GRANULAR LAYER The cells of the granular layer are diamond shaped and filled with granules (Fig. 2). The granules are




FIG. 5 Epithelium of the oral mucosa. No keratin is formed. The epithelial cells in their migration from the basal layer to the surface first become vacuolated then shrink and finally desquamate ( $\times 200$ ).

coarse, irregular in size and shape, and strongly light refractile. The chemical nature of the granules is not known. Unna called them keratohyaline granules, but the substance differs in its chemical behavior from both keratin and hyalin. The thickness of this layer

... varies from 0.02 mm on the flexor surface of the forearm to 0.5 mm and more on the soles. The cells are closely packed in this layer.

The mucous membrane of the mouth normally possesses no gran

ular cells and no horny layer. There the epithelial cells in their migration from the basal layer to the surface first become vacuolated, then shrink and finally desquamate (Fig. 5).

**THE STRATUM LUCIDUM** This layer occurs only on the palms and the soles. It is located above the granular layer and is composed of two or three layers of flat, anuclear cells of homogeneous, transparent appearance. The substance filling the cells and giving them the only appearance is called eleidin, which is believed to be formed by liquefaction of the keratohyaline granules. Ordinarily, this layer cannot be demonstrated in sections stained by routine staining methods. However, it can be shown by two rather complicated staining methods described by MacLeod and Muende.

**Pigment of the Epidermis** For the study of cutaneous pigmentation, two stains are necessary: the silver stain and the dopa stain. Silver stains melanin black, thus indicating its presence. Dopa stains

Becker, Praver and Thatcher, consists in bathing sections of skin in a 0.1 per cent solution of levorotatory 3,4-dihydroxyphenylalanine (called "dopa" for short). Following Bloch's suggestion, the cells that stain dark with the dopa stain are called dopa positive. Such cells stain dark because the enzyme tyrosinase which they contain changes the colorless dopa of the stain into a dark, insoluble product, the dopa melanin (Bloch, 1927). The dopa stain closely imitates physiologic melanin formation, which occurs because the amino acid tyrosine in cells containing the enzyme tyrosinase is transformed to dopa and further to melanin by the action of this enzyme (Lerner and Fitzpatrick).

The dopa stain shows that in the basal layer only the neural

layer of the epidermis but also are found interspersed between the hair matrix cells of the hair bulb. When active melanin formation occurs (e.g., after exposure to the sun) the number of dopa positive melanocytes in the basal layer of the epidermis increases greatly.

The silver stain does not, like the dopa stain, demonstrate the site of formation of melanin but instead demonstrates the presence of melanin. Thus, in the also mesodermal phagomatophores stain pos

*those new secretory cells migrate from the dermis only found in the layers of epidermis*

dermis melanin is seen only in the region of the basal layer. However in a deeply pigmented skin as in the Negro, melanin is found also in the upper layers of the epidermis and in phagocytic cells of the dermis. Studies by Becker Jr, Fitzpatrick and Montgomery on separated epidermis have revealed that even in deeply pigmented skin melanin is never found in basal cells or squamous cells. It is contained exclusively in melanocytes and their dendritic processes until the stratum granulosum is reached. At this level the dendrites terminate and free melanin granules are seen in the granular layer and the stratum corneum.

The Nerve of the Skin

... intra-epidermal terminal nerve fibers mediating pain have been demonstrated extending upward from the dermis into the lower layers of the stratum malpighii and in the fingers even into the stratum granulosum (Woollard, Weddell and Harpman). With the methylene blue method these nerve fibers subserving pain have been demonstrated in the lower layers of the epidermis.

... they are most abundant on the fingers where they may show connections by means of medullated fibers with Meissner's tactile corpuscles located in the papillae (Woollard).

Formerly two types of cells within the epidermis were regarded as nerve end cells, the Merkel-Ranvier tactile cells and the Langerhans cells. The Merkel-Ranvier cell is identical with the clear cell and as such is a melanocyte.

## THE EPIDERMAL APPENDAGES

The Sweat or Eccrine Glands Sweat glands are present everywhere in the human skin. They are found in greatest abundance on the palms and the soles and in the axillae. They are tubular glands whose secretory cells during the process of secretion do not change in size or shape and do not release any cellular material into the lumen of the gland. Schiefferdecker called the sweat glands eccrine glands because of his belief that their secretory cells are of the same type as those of the lacrimal glands.

Sweat glands are composed of a secretory portion and a duct portion.

The eccrine gland is found in the skin.



tissue which connects with the subcutaneous fat. Although eccrine glands are formed embryologically with two layers of epithelial cells throughout, only one distinct layer, composed of secretory cells, lines



FIG 6 Normal skin, back of neck. On the left side of the illustration, a sweat duct (SD) enters the epidermis. In the center, a large sebaceous gland (SG) leads into a follicle containing a long hair. On the right side, a large hair (H) lies within a follicle surrounded by sebaceous glands. An arrector pili muscle (AP) is situated in the obtuse angle of the hair. Beneath the large sebaceous gland, a coiled up eccrine sweat gland (SWG) is present ( $\times 50$ ).

the secretory portion in postembryonal life (Fig 7), because the second, outer layer of epithelial cells has become differentiated into myo-epithelial cells. The secretory cells are large, cylindrical cells with clear, slightly basophilic cytoplasm. The myoepithelial cells are small, spindle-shaped cells inserted sporadically between the secretory cells at their bases. Their long axis extends at a right angle to that of the secretory cells, i.e., parallel with the circumference of the

gland. They possess long processes composed of cytoplasmic fibrils with the same staining properties as smooth muscle and are contractile. By their contraction they drive the secretion out of the secretory cells into the lumen. Peripheral to the myoepithelial cells lies a thin hyaline basement membrane.



The ducts are composed of two layers of small cuboidal dark staining cells. The lumina of the ducts are lined with a homogeneous cuticle. (X400)

The ductal part of the sweat glands leads into the epidermis (Fig 6). Up to the epidermis the duct is composed of two layers of small cuboidal deeply basophilic epithelial cells. No peripheral basement membrane is present but the lumen of the duct is lined with a deeply eosinophilic, homogeneous membrane or cuticle. Some authors state that the duct is lined with myoepithelial cells in the dermis and with epidermal cells in the epidermis.

Others however like Pinkus and Memmesheimer, 1936) maintain that sweat ducts have their own lining cells within the epidermis. The observation by Holyoke and Lobitz that the cells lining the intra epidermal portion of the

sweat duct maintain their normal structure when surrounded by carcinoma tends to support Pinkus' contention.

The lumen of sweat glands is small, measuring approximately 20 microns in diameter in the secretory portion, and less still in the ductal portion. In the horny layer of the epidermis, it measures only from 5 to 10 microns in diameter. (The diameter of an erythrocyte is 7.6 microns.)

The Apocrine Glands. The apocrine glands differ from the eccrine glands in origin, distribution, mode of secretion, size and staining reactions. They represent vestigial scent glands.



FIG 8 Apocrine glands and eccrine sweat glands in the axilla. Note the great difference in the size of the lumina of the apocrine glands (left), and of the eccrine glands (right). The apocrine gland cells show "decapitation secretion" ( $\times 100$ )

directly to the epidermis

Apocrine glands are encountered in only a few areas in the axillae, around the nipples, in the perigenital and the perianal regions, and, in the external ear canal (ceruminous glands), in the breast (mammary glands). Occur also on the abdomen and the

chest.

name apocrine for these glands because part of the cytoplasm of their secretory cells is pushed off (apo = off)

because the outer layer has become differentiated just as in the sweat glands into myo epithelial cells. The secretory cells vary greatly in height depending on the stage of their secretory cycle. The secretory cells stain distinctly eosinophilic in contrast with the secretory cells of sweat glands which stain slightly basophilic. In addition they frequently contain granules which react positively to iron stains (2) (Honna) The lumen of the apocrine glands is filled with cellular (2)

on the modified skin of the vermilion border of the lips the glans penis the inner surface of the prepuce the labia minora and the they (2)  
on of  
duct (3)

a hair. On modified skin where hair is absent, sebaceous glands lead into follicles devoid of hair. The meibomian glands of the eyelids are modified sebaceous glands.

ination when the disks are crossed (see page 31) microscopic exam

Hair. The hair consists of the hair shaft, composed of keratinized cells, and the hair root, composed of nonkeratinized cells. The hair root terminates in a knoblike expansion, the hair bulb, containing the hair matrix cells. A small connective tissue structure, the papilla,




FIG 9 Lower part of a hair. The hair papilla (P) composed of connective tissue, protrudes into the hair bulb. The various linings of the hair can be recognized. They are, from the inside to the outside (1) the hair cuticle, (2) the sheath cuticle, (3) the Huxley layer, (4) the Henle layer (which stains darkly) (5) the outer hair sheath and (6) the cutaneous layer ( $\times 200$ )

protrudes into the hair bulb (Fig 9). The papilla is richly supplied with blood vessels and nerves and, in individuals with dark hair, contains considerable amounts of melanin situated largely in macrophages.

Each hair is surrounded by two inner sheaths and an outer sheath all composed of epithelial cells. The two inner sheaths develop, together with the hair, from hair matrix cells, while the outer sheath represents a downward extension of the epidermis.

The hair root consists of a medulla and a cortex. Its cells possess large, vesicular nuclei and contain variable amounts of melanin. Dopa positive melanocytes are interspersed between the hair cells.

where the hair rests on the papilla. Peripheral to the cortex is the hair cuticle which is composed of a single row of nucleated cells (Fig 9.)

The hair shaft is composed of fully keratinized cells closely united with one another and containing either no nuclei or elongated greatly shrunken nuclei. No medulla and no hair cuticle are recognizable since they have lost their identity in the process of keratinization. It may be pointed out that keratinization in the hair proceeds without the interposition of granular cells.

The two inner hair sheaths accompany the hair to approximately the level at which keratinization begins. They are separated from the hair cuticle by the sheath cuticle (Fig 9.) The sheath cuticle consists of a single layer of the cells of the hair cuticle.

granules the same layer contain darkly because they are granules. On this account, some authors have even stated that the Henle layer contained no nuclei. However they can be recognized easily in hair cut on a diagonal plane.

The ward is the epibulb. Its cells have a clear vacuolated appearance due to the presence of considerable amounts of glycogen in the cytoplasm. Outside of the outer hair sheath lies a thin membrane of condensed connective tissue the glassy or vitreous layer (Fig 9.)

Nails. The nail is composed of keratin. It grows from the nail matrix which is located beneath the nail fold. The nail matrix consists of epidermis without a stratum granulosum. Thus just as in the hair keratinization in the nail proceeds without the interposition of a granular layer. The epithelium of the nail bed has no stratum granulosum either. Its rete ridges are oriented not as a network of anastomosing ridges as elsewhere in the skin but as parallel longitudinal ridges.

## THE DERMIS

Three types of fibers occur in the dermis, collagenous, elastic and reticulum fibers.

Collagenous Fibers. Collagen forms more than 98 per cent of the connective tissue of the dermis. It occurs in bundles of fibers held

Hair. The hair consists of the hair shaft, composed of keratinized cells, and the hair root, composed of nonkeratinized cells. The hair root terminates in a knoblike expansion, the hair bulb, containing the hair matrix cells. A small connective tissue structure, the papilla,




FIG 9 Lower part of a hair. The hair papilla (P) composed of connective tissue, protrudes into the hair bulb. The various linings of the hair can be recognized. They are, from the inside to the outside (1) the hair cuticle, (2) the sheath cuticle (3) the Huxley layer (4) the Henle layer (which stains darkly) (5) the outer hair sheath and (6) the vitreous layer ( $\times 200$ )

protrudes into the hair bulb (Fig 9). The papilla is richly supplied with blood vessels and nerves and, in individuals with dark hair, contains considerable amounts of melanin situated largely in macrophages.

Each hair is surrounded by two inner sheaths and an outer sheath, all composed of epithelial cells. The two inner sheaths develop, together with the hair, from hair matrix cells, while the outer sheath represents a downward extension of the epidermis.

The hair root consists of a medulla and a cortex. Its cells possess large, vesicular nuclei and contain variable amounts of melanin. Dopa positive melanocytes are interspersed between the hair cells.

are penetrated more easily by the colloidal silver (Nageotte and Guyon).

namely, reticulum cells, histiocytes, lymphocytes, vascular endothelial cells, smooth as well as striated muscle cells and fat cells (Dublin).

of sweat glands and around the blood vessels. In addition, reticulum fibers are present in the uppermost dermis (Fig 3). In perpendicular sections, these reticulum fibers appear arranged like the bristles of a brush, but horizontal sections show a continuous fibrillar mesh work (Odland). Due to the fact that cytoplasmic processes of the basal cells extend into the mesh work, the

the abundance of reticulum fibers in pathologic conditions in which there is formation of young mesodermal cells. Large amounts of reticulum fibers are present

#### any can become converted to collagen

**Nerves and Nerve End Organs** On sections stained by routine methods, one easily can recognize the large nerves and the Pacini and Meissner end organs. The fine nerves and the other nerve end organs require special staining, either impregnation with silver salts, e.g. by the Bodian stain (Bodian), or vital staining with methylene blue (Woollard, Weddell and Harpman).

The skin is endowed with nerves from both the cerebrospinal and the autonomic or vegetative, system. The sensory nerves belong to the cerebrospinal system, while the vasomotor nerves and the nerves supplying the smooth muscles and the sweat glands

are undifferentiated to a certain degree, since the former up to their terminal ramifications always possess a myelin sheath while the latter usually do not. A few sympathetic fibers may possess a myelin sheath (Jaeger)



together by an interfibrillary ground substance. The fibers and the bundles are coarsest toward the subcutaneous tissue and finest toward the outermost portion of the dermis. In the papillary and the subpapillary portions of the dermis, the collagen bundles run in an apparently haphazard manner and do not interlace. In the lower dermis, the bundles are arranged nearly parallel to the surface of the skin and do interlace. Collagen bundles are only slightly extensible, but, since they are wavy, they permit stretching of the skin. A small number of cells, the fibroblasts, are interspersed between the collagen bundles. They have oval or spindle shaped, pale nuclei and do not stain with routine stains. The membrane (see page 36)

The bundles are interwoven with a web of elastic fibers. The fibers are slightly wavy, and, therefore, only a small portion of any fiber is seen in any histologic section. The fibers are thickest and densest in the lower portion of the dermis where they are arranged just as the collagen bundles, chiefly parallel to the surface of the skin and are up to 200 microns long. In the uppermost dermis, there are only few small fibers running horizontally, obliquely and sometimes vertically. From these fibers still finer fibrils spread out toward the epidermis but do not reach it (Dick). Therefore, elastic fibers do not contribute to the attachment of the epidermis to the dermis, as had been believed formerly.

turn of the skin into its normal position after stretching, thus supplying elasticity to the skin. Their rigidity prevents overdistention. When the skin is overstretched as in pregnancy, the elastic fibers may break and degenerate.

Reticulum Fibers Reticulum fibers (or lattice fibers "Gitterfasern") form a third system of fibers. They are not visible as such with routine stains but stain with silver (Foot's stain). It now is agreed quite generally that collagen is merely compacted reticulum fibers.

easily can see, in sections containing reticulum fibers and stained with Foot's stain, that in areas where the reticulum fibers are densest they tend to aggregate into collagen bundles (Fig. 89). The essential identity of collagen and reticulum is based on the fact that both react alike to all stains with the exception of silver stains (Mallory and Parker). The difference in argyrophilic properties is due to the fact that reticulum fibrils are finer than collagen fibrils and therefore

detected easily in microscopic sections (Fig 10) Located in the subcutis especially of the palms and the soles they aid in the mediation of the sense of pressure They are large oval omonlike structures composed of a cortex a core and a myelinated nerve which enters the structure at its lower pole The cortex consists of from 20 to 60 concentric layers of fibrous tissue the core consists of semisolid material in which the nerve is embedded The myelin sheath accompanies the nerve up to the upper pole of the Pacini corpuscle There the nerve ends in numerous ramifications

#### MEISSNER TACTILE BODIES

These are located in papillae (Fig 11) and mediate a sense of touch They are present almost exclusively on the hands and the feet and are especially numerous on the palmar surfaces of the fingers with their number increasing distally On the finger tips they lie in groups of two or three in adjoining papillae thus supplying a spatial relationship to the sense of touch (Weddell) They are cone shaped and lie with their long axis perpendicular to the surface of the skin They are of such size that they occupy the greater part of the papilla in which they are located They are composed of a connective tissue capsule and flat



FIG 11 A Meissner tactile body located in a papilla of the fingertip It is composed of flattened (schwannian) cells which have their longest diameter in a transverse direction Since this is a hematoxylin and eosin stain the nerve fibrils cannot be visualized ( $\times 400$ )

tened cells probably schwannian cells which have their longest diameter in a transverse direction One to four myelinated nerves lead to each body They lose their myelin sheath on entering the body and divide into fibrils which pass upward in a spiral fashion

**RUFFINI BODIES** The Ruffini bodies are located in the deeper dermis and in the subcutis Some believe that they mediate a sensation of heat They present brushlike ramifications of a nonmyelinated nerve within a thin connective tissue capsule

**KRAUSE BODIES** The Krause bodies are irregularly shaped spherical formations located in the dermis close to the epidermis They

The basic unit of any nerve is the neurofibril, several of which form a neurite or axon. An axon may or may not have a myelin sheath, but always is surrounded, except at its terminal ramifications, by the neuro ectodermal schwannian sheath and the mesodermal endoneurium. In medium-sized nerves, several such units are bound together by the perineurium. Large nerve trunks have several such components and an epineurium as a cover. All autonomic nerves and



FIG 10. Pacini corpuscles in the subcutaneous fat of a fingertip. Their largeness becomes apparent if one compares their size with that of the eccrine sweat glands and their ducts which are located on the right side of the field ( $\times 50$ )

most cerebrospinal nerves end freely in numerous ramifications. However, some of the latter end in special nerve end organs.

Nerve end organs are not present to an equal degree everywhere. They abound in areas of refined sensations, such as the palms and the soles, the lips and the genital region, and are sparse elsewhere. The function of some is not established clearly. Several types of nerve end organs occur: the Pacini bodies, the Meissner tactile bodies, the Ruffini bodies, the Krause bodies and the genital corpuscles. These end organs all have a similar structure: a connective-tissue capsule surrounding a core in which the afferent nerve splits up into numerous branches.

**PACINI CORPUSCLES.** The Pacini corpuscles are the largest of the end organs. They measure up to 2 mm. in diameter and thus are

collagen bundles Lymphatic vessels begin as loops in the papillae and lead down to a lymphatic plexus in the subpapillary layer from which they pass down through the dermis to a deeper plexus at the junction of the dermis and the subcutis Lymphatic vessels are lined by only one layer of endothelial cells

pilorum the tunica dartos of the scrotum and the muscle fibers in the areola of the nipple are smooth muscles The muscle fibers of the arrectores pilorum are anchored in the connective tissue of the papillae and are attached to the hair follicles below the sebaceous gland They are situated in the obtuse angle of the follicle, so that on contracting they make the follicle more vertical and produce gooseflesh (Fig 6)

Striated or voluntary muscle shows cross striation of its fibers The nuclei are located at the periphery of the fibers Striated muscle is found in the skin of the neck (platysma) and the face (muscles of expression)

#### BIBLIOGRAPHY

##### Histology of the Epidermis

- Becker S W Jr Fitzpatrick T B and Montgomery H Human melanogenesis cytology and histology of pigment cells (melanodendrocytes) Arch Dermat & Syph 63 511 1952 (Important observations.)
- Becker S W Prayer L L and Thatcher H An improved (paraffin section) method for <sup>1</sup>
- Bloch B Das P  
krankheiten
- and Schaaf F Pigmentstudien Biochem Ztschr 162 181 1925
- Dick J C Observations on the elastic tissue of the skin with a note on the reticular layer at the junction of the dermis and epidermis J Anat 81 201 1917
- Ebert M H Multiple pigmented nevi Arch Dermat & Syph 37 1 1938 (Discussion of clear cell and dendritic cell)
- Favre M Le nodule de Bizzozero dans l'épiderme norm 1 et 1 ne 1
- Ferr
- Haythorn S R Studies on the histogenesis of the so-called basal cell carcinoma Am J Cancer 15 1969 1931
- Lerner A B and Fitzpatrick T B Biochemistry of melanin formation Physiol Rev 30 91 1950
- MacLeod J M H and Muende L Practical Handbook of the Pathology of the Skin ed 3 p 49 New York Hoeber 1946
- Masson P Les naevi pigmentaires tumeurs nerveuses Ann d anat path 3 417 657 1926

have a thin fibrous capsule within which nonmyelinated nerves branch. They are believed to mediate a sensation of cold.

**GENITAL CORPUSCLES** These have the same general structure as the Krause bodies, but are larger and have a thicker capsule.

**MERKEL-RANVIER TACTILE CELLS** These cells, located in the epidermis, are discussed on page 13.

**Blood Vessels.** The arrangement of the cutaneous blood vessels is inconstant and varies in different parts of the skin. Nevertheless a deep plexus at the junction of the dermis and the subcutaneous tissue and a superficial plexus in the subpapillary layer are always present. The deep plexus has arterioles which possess three layers: adventitia, media and intima. The media and the adventitia gradually become thinner as the arteriole ascends through the dermis. The capillaries in the papillary layer are composed merely of one layer of endothelial cells surrounded by a few histiocytes (perithelial cells). In some instances, an additional type of cell, called the pericyte, is present. This cell has branching processes which encircle the capillary and are contractile. They can thus change the caliber of the lumen. They represent modified smooth muscle cells (Stout and Murray).

A special structure, the glomus, is distributed widely throughout the dermis, but occurs most abundantly on the tips of the fingers and the toes and in the nailbeds. The glomus is concerned with temperature regulation and represents a special short circuit device connecting, without the interposition of capillaries, an arteriole with a venule. It consists of an arterial and a venous segment. The arterial segment, called the Sucquet Hoyer canal, branches from an arteriole and has a narrow lumen and a thick wall. The wall consists of an inner endothelial layer and of several layers of glomus cells which are large cells with clear cytoplasm resembling epithelioid cells. The glomus cells are in intimate association with a rich network of nonmyelinated nerve fibrils demonstrable by silver stain. Although myofibrils are sparse or absent in the glomus cells, they are contractile. The glomus cell generally has been regarded as a modified smooth muscle cell (Popoff Weidman). On the basis of tissue culture experiments, Murray and Stout believe that the glomus cell is derived from the pericyte. The venous segment of the glomus has a wide lumen. The blood is emptied from the venous segment into subpapillary venules and through the latter into deeper veins.

**Lymphatic Vessels.** Lymphatic fluid circulates in the epidermis around the squamous cells, which are kept apart by intercellular bridges or prickles. Similarly, lymphatic fluid circulates between the

Odland G The morphology of the attachment between the dermis and the epidermis *Anat Rec* 108 399 1950

Popoff N W The digital vascular system *Arch Path* 18 295 1931

Woollard H H Weddell G and Harpman J A Observations on neuro-histological basis of cutaneous pain *J Anat* 74 413 1910

## 26 Histology of the Skin

- Nelemans T G Keuning F J van Rijssel T G and Ruiter M Histologic changes in the tonofibrils in vesicular and bullous diseases of the skin *Brit J Dermat* 61 177 1952
- Nieuwmeijer A H *Tonofibrils in bullous dermatoses A histo and cytopathologic study* *Dermatol* 106 379 1953
- Odlund G The morphology of the attachment between the dermis and the epidermis *Anat Rec* 108 399 1950
- Percival G H and Hannay R W Observations on the structure and formation of bullae *Brit J Dermat* 61 41 77 1919
- Stoughton R and Wells G A histochemical study on polysaccharides in normal and diseased skin *J Invest Dermat* 14 37 1950
- Thüringer J M The mitotic index of the palmar and plantar epidermis in response to stimulation *J Invest Dermat* 2 313 1939
- Woollard H H Intra epidermal nerve endings *J Anat* 71 54 1936/37
- Weddell G and Harpman J A Observations on the neurohistological basis of cutaneous pain *J Anat* 74 113 1940

### Histology of the Epidermal Appendages

- Grynfeldt E Un cas d'épithélioma baso sébacé *Bull de l'Assn franç pour l'étude du cancer* 13 174 1924
- Holyoke J B and Lobitz W C Jr Histologic variations in the structure of human eccrine sweat glands *J Invest Dermat* 18 147 1952
- Homma H On apocrine sweat glands in white and Negro men and women *Bull Johns Hopkins Hosp* 38 365 1926
- Pinkus H Notes on the anatomy and pathology of the skin appendages *J Invest Dermat* 2 175 1939
- Schnefferdecker P Die Hautdrüsen des Menschen und der Säugetiere ihre biologische und rassenanatomische Bedeutung sowie die Muscularis sexualis *Biolog Zentralbl* 37 534 1917 and Stuttgart F Enke 1922
- Wry S C The sebaceous glands *Arch Dermat & Syph* 24 353 1931
- and Memmesheimer A The sudoriparous glands—I The eccrine glands *Arch Dermat & Syph* 34 797 1936
- and — The sudoriparous glands—II The apocrine glands *Arch Dermat & Syph* 38 373 1938

### Histology of the Dermis

- Bodian D A new method for staining nerve fibers and nerve endings in mounted paraffin sections *Anat Rec* 65 89 1936
- Dick J C Observations on the elastic tissue of the skin with a note on the reticular layer at the junction of the dermis and epidermis *J Anat* 81 201 1947
- Dublin W B Reticulum *Arch Path* 41 299 1946
- Foot N C and Day H A The occurrence of reticulum in tumors *Am J Path* 14 51 1925
- Jaeger H Les méthodes d'imprégnation métallique aux sels d'argent et d'or en neurohistologie cutanée *Dermatol* 88 334 1943
- Levin O L Silvers S H and Berkowitz S S The superficial lymphatic capillary network of the skin *Arch Dermat & Syph* 36 1176 1937
- Mallory F B and Parker F Reticulum *Am J Path* 3 515 1927
- Murray M and Stout A P The glomus tumor *Am J Path* 18 183 1912
- Nageotte J and Guyon L Reticulin *Am J Path* 6 631 1930

TABLE 1—SURVEY OF STAINING METHODS EMPLOYED AT THE MASSACHUSETTS  
GENERAL HOSPITAL

FIXATION AND EMBEDDING	STAIN	PURPOSE OF STAIN	RESULTS
Helly's—paraffin (Autotechnicon)	Hematoxylin and eosin <sup>1</sup>	Routine	Nuclei blue, collagen, nerves, muscle red
Helly's—paraffin (Autotechnicon)	Phosphotungstic acid hematox- ylin	Nuclear details and for distinction of collagen and muscle	Nuclei, nerves, muscle blue, collagen brown
Helly's—paraffin (Autotechnicon)	Aniline blue (Mallory)	Collagen	Nuclei, nerves, muscle bright red, collagen blue
Helly's—paraffin (Autotechnicon)	Mason's tri- chrome	Collagen	Nuclei, nerves, muscle dark red collagen green
Helly's—paraffin (Autotechnicon)	Van Gieson	Collagen	Collagen red, nuclei, nerves, muscle yellow
Helly's—paraffin (Autotechnicon)	Verhoeff	Elastic tissue	Elastic tissue black
Helly's—paraffin (Autotechnicon)	Foot	Reticulum	Reticulum and neurofibrils black
Helly's—paraffin (Autotechnicon)	Potassium ferro- cyanide	Iron	Iron blue
Helly's—paraffin (Autotechnicon)	Giemsa	Eosinophils, bacteria, Donovan bodies, <i>Leishmania</i> , Frisch bacilli, <i>Histoplasma</i> <i>capsulatum</i>	Bacteria blue, granules of eo- sinophils, Donovan bodies, <i>Leishmania</i> , Frisch bacilli, <i>Histoplasma capsulatum</i> brilliant red
Helly's—paraffin (Autotechnicon)	Gram	Bacteria	Gram positive bacteria blue, Gram-negative bacteria red
Helly's—paraffin (Autotechnicon)	Ziehl-Neelsen Fite	Acid fast bacteria	Acid fast bacteria red Acid fast bacteria black
10% Formalin— paraffin	Levaditi	Spirochetes	Spirochetes black
10% Formalin— paraffin	Bodian	Nerves	Neurofibrils black
10% Formalin— paraffin			" "
10% Formalin— frozen section			" "
10% Formalin— frozen section	Crystal violet	Amyloid	cholesterol brownish red Amyloid purple
Absolute alcohol— celloidin	Mucicarmine	Mucin	Mucin red
80% alcohol— paraffin	Van Kossa	Calcium	Calcium black
Fresh tissue— frozen section	Dopa	Melanin producing cells	Melanin producing cells show dark granules



# 4

## Laboratory Methods

### FIXATION, EMBEDDING AND STAINING

Fixation. Helly's fixative is used routinely in the Pathology Laboratory of the Massachusetts General Hospital. For its preparation two solutions are required: (1) Zenker's solution

Potassium dichromate	25 Gm
Mercury bichloride	80 Gm
Water	1000 cc

and (2) Formalin (which is a 40 per cent aqueous solution of formaldehyde). Formalin 1 cc. is added to each 20 cc. of Zenker's solution just before use. The specimen should remain not less than 5 hours and not longer than 24 hours in Helly's fixative. It is then washed in running water for at least 2 hours and transferred into 80 per cent alcohol, in which it can be left indefinitely.

For specimens which have to be mailed, the fixative of choice is 10 per cent Formalin because specimens may remain in this fixative indefinitely. One cc. of Formalin is added to each 9 cc. of water just before use.

Helly's fixative is contraindicated if one desires to demonstrate lipids, nerve fibers, amyloid, granules of mast cells, spirochetes, calcium or mucin, or if one wishes to perform a Hopastain. For the demonstration of lipids, nerve fibers, amyloid, granules of mast cells and spirochetes, fixation in 10 per cent Formalin is required, to demonstrate calcium, 80 per cent alcohol is used, and for demonstration of mucin, absolute alcohol is the approved fixative. For the Hopastain, no fixative should be employed; the fresh tissue should merely be wrapped in moist gauze and sent to the laboratory.

**Dehydrating and Embedding.** Following fixation, all routine specimens are placed successively into several beakers containing dioxane in increasing concentrations for dehydration and several beakers of hot paraffin for embedding. This may be done by hand or with the aid of an Autotechnicon machine.

In the Pathology Laboratory of the Massachusetts General Hospital, as in most other such laboratories, all routine specimens,

TABLE 1—SURVEY OF STAINING METHODS EMPLOYED AT THE MASSACHUSETTS GENERAL HOSPITAL

FIXATION AND EMBEDDING	STAIN	PURPOSE OF STAIN	RESULTS
Helly's—paraffin (Autotechnon)	Hematoxylin and eosin	Routine	Nuclei blue collagen nerves muscle red
Helly's—paraffin (Autotechnon)	Phosphotungstic acid hematoxylin	Nuclear details and for distention of collagen and muscle	Nuclei nerves muscle blue collagen brown
Helly's—paraffin (Autotechnon)	Aniline blue (Mallory)	Collagen	Nuclei nerves muscle bright red collagen blue
Helly's—paraffin (Autotechnon)	Mason's trichrome	Collagen	Nuclei nerves muscle dark red collagen green
Helly's—paraffin (Autotechnon)	Van Gieson	Collagen	Collagen red nuclei nerves muscle yellow
Helly's—paraffin (Autotechnon)	Verhoeff	Elastic tissue	Elastic tissue black
Helly's—paraffin (Autotechnon)	Foot	Reticulum	Reticulum and neurofibrils black
Helly's—paraffin (Autotechnon)	Gram	Bacteria	Gram positive bacteria blue Gram negative bacteria red
Helly's—paraffin (Autotechnon)	Ziehl-Neelsen	Acid fast bacteria	Acid fast bacteria red Acid fast bacteria black
10% Formalin—paraffin	Levadina	Sporochetes	Sporochetes black
10% Formalin—paraffin	Bodian	Nerves	Neurofibrils black
10% Formalin—paraffin	Methylene blue Giemsa	Granules of mast cells	Mast cell granules purple
10% Formalin—frozen section	Scarlet red	Lipids	Neutral fat bright orange cholesterol brownish red
10% Formalin—frozen section	Crystal violet	Amyloid	Amyloid purple
Absolute alcohol celloidin	Mucicarmine	Mucin	Mucin red
80% alcohol paraffin	Von Kossa	Calcium	Calcium black
Fresh tissue—frozen section	Dopa	Melanin producing cells	Melanin producing cells show dark granules

whether unfixed or already placed in Helly's fixative or in 10 per cent Formalin, are sent through the Autotechnicon. This machine, which is controlled by an electric clock, accomplishes fixation, dehydration and embedding automatically overnight. The specimens are placed in small, perforated cassettes, which are then put into a perforated metal basket suspended from an arm of the machine. The metal basket is automatically lowered into and raised out of a succession of beakers containing Helly's fixative, water, 95 per cent alcohol, dioxane and warm liquid paraffin.

Specimens that are to be stained for lipids, nerve fibers, amyloid, calcium, mucin, spirochetes or granules of mast cells, or are to be treated with the dopa stain, are not sent through the Autotechnicon, in order to avoid exposure to Helly's fixative. Specimens to be stained for nerve fibers, calcium, granules of mast cells or spirochetes, after fixation, are carried by hand through dioxane and embedded in paraffin. Specimens to be stained for mucin, after fixation are embedded in celloidin. Specimens to be stained for lipids or amyloid, after fixation, are cut on the freezing microtome and then stained. Specimens to be stained with dopa are cut on the freezing microtome without previous fixation.

**Staining.** All routine specimens, after having been cut on a rotary microtome into sections from 5 to 8 microns thick, are stained with hematoxylin and eosin. With this method all nuclei stain blue, and collagen, muscle and nerves stain red. Special stains are employed for the demonstration of particular structures (see Table I).

## HISTOCHEMICAL STAINING

Brief mention may be made of two histochemical stains which have attained considerable practical importance—the Feulgen reaction and the Hotchkiss-McManus stain (DeLamater, Mescon and Barger). Both stains can be carried out on material fixed in 10 per cent Formalin.

The Feulgen reaction results in red staining of desoxyribonucleic acid (DNA) which is present in nuclei and in many virus inclusion bodies. On the other hand, ribonucleic acid (RNA), present in nucleoli, cytoplasm and keratohyalin, does not stain. This reaction is important as a stain for viruses since it often allows their differentiation from nucleoli and keratohyaline granules.

The Hotchkiss-McManus stain, or periodic acid Schiff (PAS) stain, demonstrates the presence of polysaccharides such as glycogen, hyaluronic acid and other mucoid substances by staining them red. It is also of value in the study of fibrinoid degeneration where there is depolymerization of collagen resulting in the formation of polysaccharides. For the demonstration of glycogen, it is necessary to

compare two serial sections one exposed to diastase prior to staining and the other not. Since the split products of glycogen resulting from the action of the diastase no longer are colored red by this stain only such substances represent glycogen which stain red without preliminary diastase digestion but do not stain after diastase digestion.

In addition the Hotchkiss McManus stain is of great value in the demonstration of fungi. Since the cell walls of fungi are composed of a mixture of cellulose and chitin they contain considerable amounts of polysaccharides. All fungi therefore stain bright red and thus are detected easily in histologic sections because other tissues stain very faintly or not at all (Kligman Mescon and DeLamater).

### POLARISCOPIC EXAMINATION

Polariscopic examination is the examination of histologic sections under the microscope with polarized light—i.e. with light from which all rays except those vibrating in one plane are excluded.

For the polariscopic examination two disks made of polarizing plastics are inserted into the microscope. One disk is placed below the condenser of the microscope and acts as the polarizer. The second disk is placed into the eyepiece of the microscope and acts as the analyzer. When the eyepiece containing the analyzing disk is rotated so that the path of the light through the two disks is broken at a right angle the field is dark. When however doubly refractile lipids are introduced between the two disks they break the polarization and are visible as bright white bodies in the dark field.

Cholesterol and cholesterol esters are doubly refractile while neutral fats are not. Doubly refractile fats are called anisotropic the others isotropic. Doubly refractile lipids are present regularly in the cutaneous lesions of xanthomatosis and hyperlipemia in xanthelasma palpebrarum and in extracellular cholesterosis. They are present occasionally in Hand Schüller Christian disease in foreign body granulomas and in histiocytoma. The lipid material in lipoid proteinosis and necrobiosis lipoidica diabetorum however is not doubly refractile as a rule.

### BIBLIOGRAPHY

- DeLamater E. D. Mescon H. and Barger J. D. The chemistry of the Feulgen reaction and related histo- and cytochemical methods, *J. Invest. Dermat.* 14:133 1950.
- Foot N. C. Pathology in Surgery pp 417 Philadelphia Lippincott 1945.
- Kligman A. M. Mescon H. and DeLamater E. D. The Hotchkiss McManus stain for the histopathologic diagnosis of fungus diseases. *Am. J. Clin. Path.* 21:86 1951.
- Mallory F. B. Pathologic Technique Philadelphia Saunders 1938.

## 5

# Morphology of the Mesodermal Cells

Various types of mesodermal cells infiltrate the dermis and occasionally the epidermis in the inflammatory dermatoses, in the granulomas and in the lymphomas. It is important for diagnostic purposes to identify the cell types. Three groups of mesodermal cells are recognized generally—the myeloid group, the lymphoid group and the reticular (or histiocytic) group. In addition, plasma cells and mast cells occur. It is likely that both originate from the reticular group of cells.

### MYELOID GROUP

Polymorphonuclear leukocytes and eosinophilic leukocytes may occur in the skin in various dermatoses. Basophilic leukocytes, however, do not occur. In myelosis (myeloid leukemia), in addition to polymorphonuclear leukocytes and eosinophilic leukocytes, one finds immature myeloid cells—namely, myeloblasts and myelocytes.

**Myeloblast and Myelocyte** The myeloblast is a large cell with non-granular cytoplasm and a round or oval, vesicular nucleus. The myelocyte differs from the myeloblast mainly by having in its cytoplasm either neutrophilic or eosinophilic granules. Myelocytes are always oxidase positive, while myeloblasts may be oxidase positive or oxidase negative. (For a more detailed discussion, see page 191.)

**Polymorphonuclear Leukocyte** The polymorphonuclear leukocyte, or neutrophil, has a lobated nucleus and contains fine neutrophilic granules in its cytoplasm. This cell occasionally is referred to as a microphage because it is a small phagocytic cell with the ability to phagocytize bacteria only. In contrast, the macrophage, a large phagocytic, histiocytic cell, also can take up particulate matter such as hemosiderin, melanin and lipid. Because of their ability to phagocytize bacteria, numerous polymorphonuclear leukocytes are present in the skin in acute bacterial infections, e.g., in erysipelas and folliculitis. In addition, they are found in conspicuous numbers in acute dermatitis, allergic vasculitis (anaphylactoid purpura), erythema

erythema nodosum and

illed polymorphonuclear eosinophil is characterized by the presence of coarse round brilliant eosinophilic granules in its cytoplasm. The granules are visible with routine stains but stand out more clearly when Giemsa's stain is used. Its nucleus is lobated and thus has the same appearance as that of the polymorphonuclear leukocyte. The cell is able to phagocytize bacteria. Polymorphonuclear eosinophils form as such in the bone marrow.


In addition to polymorphonuclear eosinophils monolobed eosinophils are observed not infrequently in pathologic conditions of the skin (Burkhart and Montgomery). Their nuclei are either oval kidney shaped or band shaped. It is possible that they represent histiocytic eosinophils and as such are formed in the skin.

Both the polymorphonuclear and the monolobed eosinophil occur in the tissue as a response to local anaphylaxis (Campbell). Tissue eosinophilia is apt to be prominent in eczematous drug eruptions atopic dermatitis dermatitis herpetiformis pemphigus vegetans allergic vasculitis (anaphylactoid purpura) eosinophilic granuloma granuloma faciale mycosis fungoides and Hodgkin's disease.

## LYMPHOID GROUP

It is widely accepted that lymphoid and reticular cells arise from a common cell—the lymphoid reticular stem cell (Chart 2).

CHART 2—DEVELOPMENT OF LYMPHOID AND RETICULAR CELLS



Stem Cell The stem cell can be seen in the

matin particles and one or two prominent nucleoli (see page 474 and Figure 260) This cell may develop into either a lymphoid cell (stem cell  $\rightarrow$  lymphoblast  $\rightarrow$  lymphocyte) or a reticular cell (stem cell  $\rightarrow$  reticulum cell  $\rightarrow$  histiocyte  $\rightarrow$  fibroblast) (Gall and Mallory)

**Lymphoblast.** Lymphoblasts do not occur in ordinary inflammation of the skin but only in lymphoma They are spherical cells and possess a large nucleus surrounded by a uniform, basophilic rim of cytoplasm The nucleus is round or slightly indented (see page 477 and Figure 263) It is much larger and lighter than that of the mature lymphocyte

**Lymphocyte.** Lymphocytes have much smaller nuclei and less cytoplasm than lymphoblasts The nuclei are spherical and appear very darkly stained since the chromatin forms a thick layer at the periphery and several dark staining particles in the interior

Lymphocytes are found in the skin in most forms of lymphoma and in acute, subacute and chronic inflammations—for instance, in acute, subacute and chronic dermatitis, in psoriasis in lichen planus and in lupus erythematosus They also form a large proportion of the cells in most granulomas

It is probable that the lymphocytes present in the skin are hematogenous in origin, except in malignant lymphoma, in which they may arise autochthonously It is assumed by some authors that lymphocytes may transform into histiocytes (Kolouch) Some believe even that they may transform into fibroblasts (Goldsmith)

## RETICULAR OR HISTIOCYTIC GROUP

The reticular or histiocytic group of cells belongs to the reticulo endothelial system The two outstanding properties of the reticular or histiocytic group of cells are their ability to absorb particulate matter and to produce reticulum fibers As the cells age they may change into fibroblasts (see Chart 2, page 33) As such, they no longer possess phagocytic powers and form collagen rather than reticulum

No generally accepted nomenclature exists for the reticular group of cells The following names are used by various authors as designation for the mature cell of this group: histiocyte, reticulum cell, clasmatocyte, perithelial cell, tissue monocyte and wandering connective tissue cell The term histiocyte at present is employed the most widely Under special conditions, the histiocyte may become a macrophage or an epithelioid cell

Immature cells of the reticular groups are observed in lymphoma and generally are referred to as reticulum cells Therefore, lymphomas in which these cells predominate are called reticulum cell

lymphoma. However, it should be kept in mind that many authors use the term reticulum cell also for mature reticular cells.

**Reticulum Cell (Immature Reticular Cell)** This cell, as seen in reticulum-cell lymphoma, possesses a nucleus which is smaller than that of a stem cell but larger than that of a histiocyte. The nucleus may be round but usually is oval or kidney shaped. It is pale staining and has a moderately heavy chromatin network and a distinct basophilic nuclear membrane (see page 174 and Figure 261). The cytoplasm is eosinophilic and abundant. It often is irregular in outline and may show pseudopodic protrusions. Because of their immaturity, the cells may form reticulum fibers only to a slight degree and have little or no phagocytic power.

**Histiocyte (Mature Reticular Cell)** Histiocytes are formed in the skin in contrast to lymphocytes which do not arise in the skin except in malignant lymphoma.

In the normal skin, histiocytes are present in small numbers around the capillaries (perithelial cells). Under pathologic conditions which include the inflammatory diseases and the granulomas, histiocytes tend to wander from the pericapillary regions into the dermis (tissue monocytes—wandering connective tissue cells).

Histiocytes are much smaller than reticulum cells but otherwise do not differ from them in appearance. Histiocytes form abundant reticulum fibers and possess the ability to phagocytize bacteria and particulate matter.

Histiocytes may resemble fibroblasts greatly and often it is not possible to state with certainty whether a given cell is a histiocyte or a fibroblast. As a rule, however, the nuclei of histiocytes are larger and stain slightly paler than those of fibroblasts. Although both may have oval nuclei, those of histiocytes tend to be kidney shaped, those of fibroblasts spindle shaped. While histiocytes possess the ability to form reticulum fibers, fibroblasts form collagen. It should be remembered that no sharp borderline exists between the histiocyte and the fibroblast, since the histiocyte presents the parent cell of the fibroblast and may develop into a fibroblast.

**Macrophage** Histiocytes possess the ability to phagocytize particulate matter and certain microorganisms. Those containing phagocytized material are called macrophages. Histiocytes migrate as wandering cells to areas where material digestible to them is present. They may ingest melanin or melanophor.

They may also ingest bacteria, fungi, etc. Examples of bacteria which may be ingested by histiocytes are lepra bacilli and Frisch bacilli, of



fungi, *Histoplasma capsulatum*; of viruses, Donovan bodies, and of protozoa, *Leishmania tropica*. When individual macrophages are unable to deal with particles to be removed, they tend to fuse together and to form multinucleated foreign body giant cells. Excellent examples of foreign body giant cells may be seen in paraffinoma, gout and Malherbe's calcifying epithelioma. The nuclei in foreign body giant cells usually are clumped together in an irregular arrangement, but they may lie regularly along the periphery of the cell, so that foreign body giant cells may be indistinguishable from Langhans giant cells.

**Epithelioid Cell.** Under certain conditions, histiocytes may change into epithelioid cells. Epithelioid cells possess a large, usually oval, pale, vesicular nucleus resembling the nucleus of epithelial cells and abundant, ill defined, slightly eosinophilic cytoplasm. Pseudopodic elongations of the cytoplasm usually can be seen. When lying in groups, the cytoplasm of neighboring cells often appears coalesced. Epithelioid cells may form giant cells, the so called Langhans type of giant cell. It is likely that this type of giant cell forms by amitotic nuclear division without cellular division. The nuclei are arranged in an arc along the periphery of the cell in horseshoe fashion. Epithelioid cells and Langhans giant cells have phagocytic power and have the ability to form reticulum fibers.

Epithelioid cells are found in a variety of granulomas, especially in tuberculosis, sarcoidosis, leprosy, syphilis, blastomycosis and coccidioidomycosis. In tuberculosis epithelioid cells form as a tissue response to the lipid fraction of the tubercle bacillus (Sabin). As a granulomatous lesion containing epithelioid cells heals, the epithelioid cells may mature into fibroblasts and their reticulum fibers into collagenous fibers. This process can be observed particularly well in healing lesions of sarcoidosis.

**Fibroblast.** The common (resting) connective tissue cells are called fibroblasts because they are instrumental in the elaboration of the collagenous fibers (see page 20). Their nuclei are elongated, often spindle shaped. Because of their pale staining and the presence of a fine nuclear membrane, the nuclei have a vesicular appearance. The cell body, which is spindle shaped, is not discerned easily. Fibroblasts usually are found adjacent to the surface of collagenous bundles.

## PLASMA CELLS

The plasma cell has abundant cytoplasm which is deeply basophilic, homogeneous and sharply defined. The nucleus is eccentrically placed and round, and along its membrane shows coarse, dark stain

lished. All stages of transition between reticulum cells and plasma cells can be observed in the spleen of rabbits after repeated intravenous injections of horse serum. Antibodies are formed in great quantities in reticulum cells during their development into plasma cells while mature plasma cells have already passed the stage of greatest activity (Fagraeus). It is assumed by some that not only reticulum cells but also all multipotent cells of the connective tissue as well as lymphocytes are capable of transforming into a plasma cell (Campbell and Good).

Plasma cells occur in small numbers in most chronic inflammatory diseases of the skin and in larger number in granulomas. They are particularly conspicuous in syphilis, granuloma inguinale and rhinoscleroma. In the presence of many plasma cells but especially in rhinoscleroma round hyaline acidophilic bodies so-called Russell bodies may be found within and outside of plasma cells. They form within plasma cells as a phenomenon of degeneration and finally are expelled (Pearse). They may attain a size twice that of a normal plasma cell (See also page 202).

## MAST CELLS

Mast cells and basophilic leukocytes have nothing in common aside from an identical basophilic metachromatic staining reaction of their granules. They are derived from a myeloid stem cell and have an oval nucleus and have a lobate nucleus.

The granules of mast cells are soluble in ordinary fixatives such as Helly's and Zenker's fluids and do not stain with hematoxylin-eosin. Ten per cent Formalin, absolute alcohol and saturated basic lead acetate are suitable as fixatives and methylene blue or Giemsa's stain is suitable for staining the granules. The granules are basophilic (i.e. they stain with basic aniline dyes) and metachromatic (i.e. they may stain in a color different from that possessed by the dye). In this sense the bluish dye thionine stains the mast-cell granules a reddish violet, toluidine blue stains them a purplish red and polychrome methylene blue stains them red (Michels).

Mast cells have two known functions: secretion of hyaluronic acid and secretion of heparin. That hyaluronic acid is present in the mast cell granules is proved by the observations that mast cell gran-

ules stain exactly like hyaluronic acid and no longer show metachromatic staining after the tissue containing them has been submitted to the action of hyaluronidase (Asboe Hansen). Evidence for the presence of heparin in the mast cell granules is the similarity of reaction of the two substances to metachromatic staining and the parallelism existing between the amount of extractable heparin and the mast cell content of certain organs (Jorpes).

Mast cells occur everywhere in the connective tissue of the body particularly in the vicinity of capillaries and in the walls of larger blood vessels. The normal skin contains relatively few mast cells which are small and spindle shaped. As a rule they are arranged in groups around the blood vessels but they occur also around the hair follicles and in the papillary layer of the dermis. Their number is increased in many different conditions (Asboe Hansen). For instance the granulation tissue in healing wounds contains more mast cells than normally are present. In most itching dermatoses such as atopic eczema, contact dermatitis and lichen planus an increased number of mast cells is present around the capillaries. In lupus erythematosus in which there is an increase in the amount of hyaluronic acid in the dermis the number of mast cells closely parallels the intensity of metachromatic staining. Also neurofibromas and the stroma of carcinomas of the skin contain numerous mast cells. Thus an increase in mast cells is of no diagnostic significance except in urticaria pigmentosa where they occur in tumor like aggregates especially in the upper dermis.

# BIBLIOGRAPHY

- Asboe Hansen C. A survey of the normal and pathological occurrence of mucinous substances and mast cells in the dermal connective tissue in normal tissue eosinophilia and in neurotropic virus diseases. *Ann. Allergy* 7:471, 1949.
- Campbell D. H. Relationship of the eosinophil response to factors involved in anaphylaxis. *J. Infect. Dis.* 72:42, 1943.
- Fagraeus A. Antibody production in relation to the development of plasma cells. *Acta Path. Microbiol. Scand.* 20:1, 1948.
- Jorpes E. J. Heparin. p. 30. London: Oxford Univ. Press, 1939.
- Kolouch F. Jr. The lymphocyte in acute inflammation. *Am. J. Path.* 15:113, 1939.
- Maximow A. A. and Bloom W. A Textbook of Histology, ed. 4, pp. 56-60. Philadelphia: Saunders, 1943.

- Michels N A The mast cells in Downey H Handbook of Hematology, vol I, p 235 New York Hoeber, 1938
- Pearse A G E The nature of Russell bodies and Karloff bodies J Clin Path 2 81, 1949
- Robb Smith A H T The reticular tissue and the skin Brit. J Dermat 56 151, 1944 (Instructive review)
- Sabin F R Cellular reactions to tuberculo proteins compared with reactions to tuberculo-lipids J Exper Med 68 837 1938
- Wislocki G B Bunting H, and Dempsey E W Metachromasia Am J Anat 81 1 1947

# 6

## Congenital Diseases (Genodermatoses)

### ICHTHYOSIS

Two forms of ichthyosis occur *ichthyosis vulgaris* and *ichthyosis congenita*. In *ichthyosis vulgaris*, the skin is dry and rough and shows



FIG. 12 *Ichthyosis vulgaris*. There are hyperkeratosis and absence of the granular layer. The stratum malpighi is thin, the rete ridges irregular. A large keratotic plug is located within a hair follicle ( $\times 100$ ).

scaling, often in the form of large lamellae resembling fish scales. Involvement is severest on the extensor surfaces of the extremities, while the flexures usually are spared. Follicular hyperkeratoses are frequently present.

*Ichthyosis congenita* is a more severe form of *ichthyosis* than

*ichthyosis vulgaris* The skin is represented by a thick horny cuirass with development of deep fissures

*Ichthyosis hystrix* is not related to *ichthyosis* but represents an extensive or systematized *nevus verrucosus* (see page 321)

**Histopathology** The characteristic lesion of *ichthyosis vulgaris* is hyperkeratosis with diminution or even complete absence of the granular layer (Fig 12) This represents an exception to the rule that an increase in thickness of the horny layer is accompanied by an increase in thickness also of the granular layer The probable reason for this exceptional finding is that in *ichthyosis vulgaris* there is an inadequate shedding rather than an overproduction of horny cells The hyperkeratosis in *ichthyosis vulgaris* is moderate and not associated with parakeratosis The stratum malpighii is thinner than normal and the rete ridges are diminished in number some are atrophic while others are slender elongated and branching (dove tailed) The hyperkeratosis usually affects also the upper portion of the hair follicles resulting in large follicular keratotic plugs The pressure of the keratotic plugs causes atrophy of the lower portion of the follicles and of the sebaceous glands

*Ichthyosis congenita* differs from *ichthyosis vulgaris* only by showing a much greater degree of hyperkeratosis A stratum granulosum is usually present but it is not prominent

## CONGENITAL ICHTHYOSIFORM ERYTHRODERMA

Like *ichthyosis* this disease is characterized by dryness roughness and scaling of the skin In addition generalized erythema is present and in many cases bullae occur In contrast to *ichthyosis* the flexural surfaces are involved most intensely

**Histopathology** In the erythematous areas of the skin one observes pronounced hyperkeratosis with occasional islands of parakeratosis A stratum granulosum is present Although it varies in thickness it is for the most part hypertrophic The stratum malpighii shows acanthosis with irregular elongation of the rete ridges The upper dermis shows a chronic inflammatory infiltrate

Examination of bullous lesions has revealed in some cases merely presence of a nonspecific subcorneal bulla (Mackee and Rosen) In others a pronounced ballooning of epidermal cells was noted either in the midportion of the epidermis (Ellis) or in the granular layer (Barker and Sachs) In the latter case the changes in the granular layer resembled those seen in *epidermodysplasia verruciformis* Ellis believes that the ballooning of epidermal cells is caused by a persistence of embryonal cells inasmuch as the cells of the embryonal stratum intermedium have a ballooned appearance (see page 4)

**Differential Diagnosis.** Congenital ichthyosiform erythroderma differs from ichthyosis by the presence of a usually hypertrophic granular layer, the presence of acanthosis and the presence of an inflammatory infiltrate in the dermis. In ichthyosis, the granular layer is diminished in thickness or absent, the stratum malpighii is thinner than normal and no infiltrate is present in the dermis (Layman and Murphy).

### KERATOSIS PALMARIS ET PLANTARIS

This condition, which often is hereditary, is characterized by diffuse thickening of the horny layer of the palms and the soles. Because of the formation of fissures, the condition causes great discomfort to the patient.

**Histopathology.** The histologic picture is nonspecific, showing considerable hyperkeratosis as the only constant change. Occasionally there also are acanthosis and a chronic inflammatory infiltrate in the upper dermis.

### KERATOSIS PUNCTATA PALMARIS ET PLANTARIS

Multiple, discrete, yellow to dark brown, firm, slightly elevated, conical keratotic plugs, 1 to 3 mm in diameter, are distributed symmetrically over the palms and the soles.

**Histopathology.** One observes a circumscribed hypertrophy of the stratum corneum consisting of a cone shaped plug which invaginates the subjacent structures. Beneath this plug the stratum malpighii is thinned. The dermis is free of any inflammatory infiltrate (Scott, Costello and Simuango).

### PACHYONYCHIA CONGENITA

This disease shows dystrophic changes of the nails, palmar and plantar hyperkeratosis, follicular keratosis and leukoplakia of the oral mucosa.

A variety of pachyonychia congenita is *dyskeratosis congenita*. It shares with pachyonychia congenita the dystrophic changes of the nails and the presence of leukoplakia. However, there are no hyperkeratotic changes of the skin. Instead the skin shows atrophy and patches of hyperpigmentation.

**Histopathology.** In pachyonychia congenita one observes hyperkeratosis with areas of parakeratosis. There is follicular plugging. Occasionally, horn plugs are present also in the sweat ducts (Andrews). The granular layer is hypertrophic. The stratum malpighii shows acanthosis with elongation of the rete ridges. A mild chronic inflammatory infiltrate is present in the upper dermis. Dyskeratotic

changes, similar to the corps ronds of Darier's disease, have been observed in the stratum malpighii of some cases (Andrews, Wright and Guequierre)

Dyskeratosis congenita shows thinning of the epidermis and almost complete absence of rete ridges. In areas in which the skin shows hyperpigmentation, the amount of melanin in the basal layer is increased and melanophores are present in the upper dermis (Cole, Rauschkolb and Toomey)

## POROKERATOSIS MIBELLI

One or several lesions may be present. They show an atrophic center surrounded by a raised keratotic wall. The wall has on its top a groove filled with keratotic material.



FIG. 15 Porokeratosis Mibelli. The section is taken from the keratotic wall. It shows the deep groove which forms the center of the keratotic wall. The groove is filled with a large horn plug (the 'cornoid lamella'). The cornoid lamella has a column of parakeratotic cells in its center. On the right is a normal sweat duct.

**Histopathology.** The atrophic center shows slight hyperkeratosis, atrophy of the stratum malpighii and fibrosis of the upper dermis. The keratotic wall shows considerable hyperkeratosis and acanthosis. In its center, the keratotic wall presents a deep groove filled by a large horn plug (the 'cornoid lamella'). This cornoid lamella has



in its center a column of parakeratotic cells. Beneath the parakeratotic column the granular layer is absent elsewhere it is well developed. The dermis underlying the cornoid lamella shows a chronic inflammatory infiltrate (Fig. 13).

Owing to the presence of the cornoid lamella the histologic picture of porokeratosis is diagnostic. The name porokeratosis is a misnomer since the cornoid lamella is not necessarily located in the opening of a sweat duct.

### **XERODERMA PIGMENTOSUM**

This disorder is associated with hypersensitivity to ultraviolet light. The lesions occur chiefly on exposed areas of the skin. An early and a late stage occur. In the early stage one observes slight diffuse erythema with scaling and small areas of hyperpigmentation resembling freckles. In the late stage atrophy of the skin, mottled pigmentation and telangiectases are present giving the skin an appearance similar to that of chronic radiodermatitis. Warty growths appear within the atrophic skin which may develop into carcinoma.

**Histopathology.** In the early stage the histologic appearance is not always characteristic; the diagnosis often can be made from the combination of the histologic changes, however. There are (1) hyperkeratosis, (2) thinning of the stratum malpighii with atrophy of some of the rete ridges and irregular proliferation and prolongation of others, (3) edema of the upper dermis, (4) a chronic inflammatory predominantly perivascular infiltrate in the upper dermis and (5) spotted melanin pigmentation of the basal layer with melanophores in the upper dermis.

In the late stage the epidermis shows atrophy in some areas and acanthosis in others. Atypical and multinucleated cells may be seen in the epidermis. The hyperkeratosis and hyperpigmentation already present in the early stage are more pronounced. The upper dermis shows degenerative changes of the collagen and of the elastic fibers of the same type as is seen in senile degeneration of the skin. Thus one observes basophilic degeneration of the collagen and senile elastosis (see page 157). In some areas the epidermis may show atypical downward growth so that the histologic picture in such areas is identical with that of senile keratosis.

Ultimately squamous-cell carcinomas and occasionally basal cell epitheliomas and sarcomas develop in some of the lesions.

### **CONGENITAL ECTODERMAL DEFECT**

This condition represents an incomplete development of the epidermis and its appendages. The skin is smooth and glossy. Hair

growth is sparse or completely absent. The facies is typical. It shows depressed nasal root and bridge, prominent frontal bosses and thick lips. In addition, there may be dental aplasia and dystrophy of the nails. Because of the diminution or the absence of sweat glands, the patient is unable to sweat adequately and therefore is intolerant to heat.

**Histopathology.** There is either a total absence of sweat glands or the presence of rudiments of nonfunctioning glands and ducts. There is usually a similar deficiency of pilosebaceous structures. The epidermis is thinner than normal and there may be likewise a reduction in the width of the dermis. The collagen, the elastic fibers and the blood vessels are normal in appearance, as a rule. Examination of the axillary skin in two cases revealed total absence of eccrine glands, while the apocrine glands were developed normally (Sunderman).

### ROTHMUND'S SYNDROME, WERNER'S SYNDROME (PROGERIA OF ADULTS) AND PROGERIA OF CHILDREN

These are three different, though related, genodermatoses in which

infancy with

the affected skin shows erythema, scaling, telangiectases and brownish pigmentation so that the appearance of the skin resembles that of poikiloderma atrophicans.

and the musculature of the extremities undergo atrophy leading on the legs to ulcerations. Cataracts develop early in adult life.

*Progeria of children* is not familial. It starts several months after birth. The patient develops into a dwarf with a large skull and bird-like features. The skin appears atrophic and wrinkled.

**Histopathology.** No characteristic histologic features are seen in the skin in any of the three diseases. In all of them, the skin merely shows atrophy of the epidermis, thinning of the collagen bundles in the dermis and atrophy or even disappearance of the cutaneous appendages (Reed). The areas of erythema in Rothmund's syndrome, in spite of their clinical resemblance to poikiloderma atrophicans vasculare, show no inflammatory infiltrate but merely atrophy and telangiectasis (Thannhauser).

in its center a column of parakeratotic cells. Beneath the parakeratotic column the granular layer is absent elsewhere it is well developed. The dermis underlying the cornoid lamella shows a chronic inflammatory infiltrate (Fig 13).

Owing to the presence of the cornoid lamella the histologic picture of porokeratosis is diagnostic. The name porokeratosis is a misnomer since the cornoid lamella is not necessarily located in the opening of a sweat duct.

### XERODERMA PIGMENTOSUM

This disorder is associated with hypersensitivity to ultraviolet light. The lesions occur chiefly on exposed areas of the skin. An early and a late stage occur. In the early stage one observes slight diffuse erythema with scaling and small areas of hyperpigmentation resembling freckles. In the late stage atrophy of the skin, mottled pigmentation and telangiectases are present giving the skin an appearance similar to that of chronic radiodermatitis. Warty growths appear within the atrophic skin which may develop into carcinoma.

**Histopathology** In the early stage the histologic appearance is not always characteristic the diagnosis often can be made from the combination of the histologic changes however. There are (1) hyperkeratosis (2) thinning of the stratum malpighii with atrophy of some of the rete ridges and irregular proliferation and prolongation of others (3) edema of the upper dermis (4) a chronic inflammatory predominantly perivascular infiltrate in the upper dermis and (5) spotted melanin pigmentation of the basal layer with melanophores in the upper dermis.

In the late stage the epidermis shows atrophy in some areas and acanthosis in others. Atypical and multinucleated cells may be seen in the epidermis. The hyperkeratosis and hyperpigmentation already present in the early stage are more pronounced. The upper dermis shows degenerative changes of the collagen and of the elastic fibers of the same type as is seen in senile degeneration of the skin. Thus one observes basophilic degeneration of the collagen and senile elastosis (see page 157). In some areas the epidermis may show atypical downward growth so that the histologic picture in such areas is identical with that of senile keratosis.

Ultimately squamous cell carcinomas and occasionally basal cell epitheliomas and sarcomas develop in some of the lesions.

### CONGENITAL ECTODERMAI DEFECT

This condition represents an incomplete development of the epidermis and its appendages. The skin is smooth and glossy. Hair

scarring results. In *hydroa aestivale* the changes in the dermis are less severe than in *hydroa vacciniforme*. Therefore healing takes place without scar formation.

## EPIDERMOLYSIS BULLOSA

In this condition vesicles or bullae form usually at points of trauma but sometimes without trauma. Three forms of the disease exist: *epidermolysis bullosa simplex*, *epidermolysis bullosa dystrophica* and *epidermolysis bullosa hereditaria letalis*. The simple form is inherited dominantly and the other two forms recessively. In the simple form the bullae heal without scarring, the mucous membranes and the nails are rarely affected and the disease improves or even subsides at puberty. In the dystrophic form the lesions heal with atrophic scars, oral lesions and dystrophic changes of the nails are frequently present and the disease persists throughout life. In the third form, *epidermolysis bullosa hereditaria letalis*, death usually occurs within the first 3 months of life. The bullae show little tendency to heal but if they heal no scars remain. Oral lesions and dystrophic changes of the nails are usually present.

**Histopathology.** The bullae are always in subepidermal location in the dystrophic form (Tulipan) and in the lethal form (Lamb and Halpert, Schaffer) whereas in the simple form they may be found in subepidermal location (Leoni), in intra epidermal or in subcorneal location (Johnson and Test). However it is likely that also in *epidermolysis bullosa simplex* the bullae always form subepidermally but because of the tendency of the epidermis to rapid regeneration the cleavage will be found intra epidermally or subcornerally in bullae that are a few days old. (This occurs not infrequently in *erythema multiforme* and in *bullous pemphigoid* see page 86.) No significant inflammatory reaction is observed in the dermis unless secondary infection has occurred. In the dystrophic form small milium like epidermal cysts may be found in the upper dermis (Tulipan).

Engman and Mook first described the absence of elastic fibers in the papillary and the subpapillary layers of involved as well as non-involved areas. They believed that the absence of elastic fibers was the cause of the disease. Although some authors have confirmed this finding others have found the elastic tissue to be normal (Allen). In general it appears that in the simple form the elastic tissue is normal (Johnson and Test) while in the dystrophic form the elastic tissue tends to be absent in the upper dermis of the involved areas (Tulipan, Lamb and Halpert). However it is probable that the absence of elastic tissue in the dystrophic form is not primary but secondary and is the result of its destruction by the disease process. In

## HYDROA VACCINIFORME AND HYDROA AESTIVALE

Hydroa is a recurring papulovesicular eruption occurring chiefly in the summer season, usually in boys, and solely on the exposed parts of the skin. Two forms exist: *hydroa aestivale*, the milder form, which does not produce scarring and ends at puberty, and *hydroa vacciniforme*, which produces scarring and persists as a rule through-




FIG. 14. *Hydroa vacciniforme*. There is a wedge-shaped area of necrosis involving the epidermis and the upper dermis. It is walled off by a chronic inflammatory infiltrate. ( $\times 100$ )

out life. In about one third of the cases, *hydroa vacciniforme* is associated with congenital porphyria, i.e., excretion of large amounts of uroporphyrin and coproporphyrin in the urine (see page 278).

**Histopathology.** In *hydroa vacciniforme*, a focal area of inflammation followed by necrosis forms in the upper dermis (Fig. 14). In the overlying epidermis, an intra-epidermal vesicle develops which at first is multilocular, but later, because of the degeneration of epithelial cells, becomes unilocular. The vesicle is filled with fibrin, leukocytes and the debris of epithelial cells. Within the area of necrosis located in the upper dermis, the blood vessels are thrombosed and foci of hemorrhage occur. The necrotic area is walled off with a chronic inflammatory infiltrate. On absorption of the necrotic area,

and grains (2) formation of lacunae and (3) irregular upward proliferation of papillae into the lacunae resulting in the formation of villi (Fig 15). There also are papillomatosis, acanthosis and hyperkeratosis. The dermis shows a chronic inflammatory infiltrate. In



the granular layer are several corps ronds (C.R.), characterized by large round homogeneous deep staining nuclei. In the lower third within a lacuna are grains (G.) characterized by small often grain shaped nuclei ( $\times 400$ ).

some cases in addition there is downward proliferation of epidermal cells into the dermis.

The corps ronds occur mainly in the stratum malpighii and the granular layer while the grains occur mainly in the horny layer. Both may be found within the lacunae. Corps ronds possess large, round homogeneous deeply basophilic nuclei and a homogeneous (hyalinized) deeply eosinophilic cytoplasm lined by a distinct membrane. They are much larger than normal squamous cells (Fig 16). Corps

any case, recent studies of normal elastic tissue indicate that it plays no part in the coherence between epidermis and dermis (see page 8).

**Differential Diagnosis.** Differentiation of epidermolysis bullosa from other bullous diseases often is impossible. The presence of small epidermal cysts and the absence of elastic tissue, however, may aid in the establishment of the diagnosis.

### DARIER'S DISEASE (KERATOSIS FOLLICULARIS)

This disease is characterized by a more or less extensive eruption consisting of hyperkeratotic or crusted papules which by confluence



Fig 15 Darier's disease Low magnification There are hyperkeratosis and papillomatosis. Numerous lacunae (L) are present. On the left are elongated papilli lined by only one layer of cells so called villi (V). Corps ronds (C.R.) are present in the granular layer and grains (G) in the horns layer and in some of the lacunae (X100)

may form verrucous, crusted areas. Occasionally, hypertrophic lesions are present with elevated, verrucous formations. The oral mucosa is commonly, and the vulva, the larynx and the pharynx occasionally, involved (Brunauer).

**Histopathology.** The characteristic changes in Darier's disease are (1) a peculiar form of dyskeratosis, namely, formation of corps ronds

tion, two rows of epithelial cells line the villi. Furthermore, no dyskeratosis occurs and large apocrine glands are present in the dermis

## FAMILIAL BENIGN CHRONIC PFMPHIGUS

(Hailey and Hailey)

This disease (which often, but not always, is familial) is characterized by a localized recurrent eruption of vesicles and bullae. By peripheral extension, the lesions may assume a circinate configuration.

**Histopathology.** Although early lesions may show like Darier's disease small suprabasal separations so-called lacunae fully devel-



FIG. 17. Familial benign chronic pemphigus (Hailey and Hailey). The bulla is suprabasal in position. One observes loss of intercellular bridges resulting in considerable acantholysis and formation of villi (upward growth of papillae). These features cause a surprising resemblance to the bulla of pemphigus vulgaris. ( $\times 200$ )

oped lesions show large separations, namely bullae in suprabasal position (Fig. 17). Villi i.e., elongated papillae lined by a single layer of basal cells protrude upward into the bullae and, in some cases, cords of basal cells proliferate downward into the dermis. The bullae con-



ronds develop on account of premature partial keratinization of the cell prior to reaching the horny layer, a process called benign dyskeratosis (Malignant dyskeratosis is observed in Bowen's disease and in squamous cell carcinoma)

The grains are small cells, considerably smaller than the corps ronds. They resemble parakeratotic cells except that their nuclei are more prominent. The nuclei are elongated, often grain shaped.

The lacunae represent small, slitlike, intra epidermal vesicles which are found most commonly directly above the basal layer. They contain desquamated, acantholytic epidermal cells which have lost their intercellular bridges due either to degenerative changes or to partial keratinization. Corps ronds and, especially, grains are present among these desquamated cells.

Elongated, often tortuous papillae lined usually with but a single layer of basal cells project into the lacunae. They often are referred to as villi.

The hyperkeratosis and the papillomatosis cause the formation of keratotic plugs. They often fill the pilosebaceous follicles but also are found outside of follicles. Darier's disease, therefore, is not, as Darier originally thought, primarily a follicular disease. Proof of this is the fact that areas devoid of follicles, such as palms, soles and oral mucosa, may be affected (Ellis). The term keratosis follicularis is, then, a misnomer.

In some cases of Darier's disease, but especially in those with hypertrophic lesions, one observes considerable downward proliferation of the epidermis, either as a proliferation of basal cells or as pseudoepitheliomatous hyperplasia (Beerman). The proliferations of basal cells consist of long narrow cords composed of two rows of basal cells between which there may or may not be a lacunar space. These proliferations may send out branches and may penetrate deep into the dermis. The pseudoepitheliomatous hyperplasia may suggest squamous cell carcinoma but, so far, no case of Darier's disease resulting in malignancy has been reported. (For a discussion of pseudoepitheliomatous hyperplasia see page 334.)

The lesions on the oral and other mucous membranes are analogous to those observed on the skin except that hyperkeratotic changes are mild or absent (Brunauer).

**Differential Diagnosis** For the differentiation of Darier's disease from familial benign chronic pemphigus see below. The villous proliferations into the lacunar spaces may resemble those of syringocystadenoma papilliferum (Beerman). However, in the latter condi-

asionally into basal cell epithelioma (Sullivan and Ellis) or squamous-cell carcinoma (Costa and Junqueira; Ormea).

In 1946, Lutz (one of the original describers), on the basis of successful implantation tests, stated that epidermodysplasia is "not an independent dermatosis, but rather a generalized eruption of warts with somewhat peculiar characteristics." Yet, some authors, though conceding that Lutz' case was one of verrucae planae, still regard epidermodysplasia verruciformis as an entity (Ormea; Teodorescu, Midana). Midana obtained negative results in auto inoculation experiments performed on two patients. Teodorescu found in two patients associated with the epidermodysplasia verruciformis, keratosis palmaris et plantaris hereditaria and regarded both diseases as genodermatoses.

### ACROKERATOSIS VERRUCIFORMIS

(Hopf)

Numerous hyperkeratotic and occasionally verrucous papules are present, predominantly on the dors of the hands and the feet.




FIG. 18. Acrokeratosis verruciformis (Hopf). The lesion consists of a circumscribed area of papillomatous, acanthosis and hyperkeratosis ( $\times 100$ ).

**Histopathology.** The papules show considerable hyperkeratosis, increase in thickness of the granular layer and acanthosis (Fig. 18). In addition, there are slight papillomatosis and some thickening of the rete ridges. The rete ridges all extend to a uniform level and their configuration is well preserved. There is no parakeratosis or

acantholytic cells some of which show evidence of keratinization. Occasionally there is evidence of dyskeratosis namely shrinking of some of the acantholytic cells which then assume the appearance of grains. However corpus ronds have been described in only a few cases (Ellis Winer and Leeb).

**Differential Diagnosis** Familial benign chronic pemphigus shares certain histologic features with both Darier's disease and pemphigus vulgaris. In all three diseases one finds suprabasal separations of the epidermis caused by acantholysis and upward proliferation of papillae as so called villi into the resulting lacunae or bullae. Familial benign chronic pemphigus however differs from Darier's disease by the larger size of the suprabasal separations (which thus appear as bullae rather than as lacunae) and the lesser degree or even the absence of dyskeratosis. If dyskeratosis is absent differentiation from pemphigus vulgaris may be impossible. Occasionally however one observes in familial benign chronic pemphigus even in the absence of dyskeratosis an early onset of keratinization in the lower layers of the detached epidermis. The presence of eosinophils in the bullae points toward a diagnosis of pemphigus vulgaris but their absence does not rule it out.

The nosologic position of the disease is at present uncertain. The presence of dyskeratotic changes in most cases of familial benign chronic pemphigus has led some observers to regard this disease as a bullous variant of Darier's disease (Ellis Finnerud and Szymanski Winer and Leeb). However until more is known about the cause of the disease it may be best to regard it as an independent entity (Hailey and Hailey Trunk and Rein).

## EPIDERMODYSPLASIA VERRUCIFORMIS

(Lewandowsky and Lutz)

In this dermatosis one finds an extensive eruption of flat topped hyperkeratotic lesions resembling verrucate plaques. By confluence lichenified plaques may form.

**Histopathology** The histologic changes consist of hyperkeratosis increase in the thickness of the granular layer acanthosis and a peculiar vacuolization of the cells in the upper layers of the stratum malpighii of the granular cells and of the horny cells. Because of the vacuolization the horny layer shows a loosely felted basket weave pattern. The histologic picture is like that of verrucate plaque (Fig. 122) with two exceptions (1) in epidermodysplasia verruciformis the nuclei of the vacuolated cells show more pronounced pyknosis and fragmentation than in verrucate plaque (Sullivan and Ellis Waisman and Montgomery) and (2) the lesions change oc-

casionally into basal cell epithelioma (Sullivan and Ellis) or squamous cell carcinoma (Costa and Junqueira, Ormea)

In 1946, Lutz (one of the original describers), on the basis of successful implantation tests, stated that epidermodysplasia is "not an independent dermatosis, but rather a generalized eruption of warts with somewhat peculiar characteristics." Yet, some authors, though conceding that Lutz' case was one of verrucar planae, still regard epidermodysplasia verruciformis as an entity (Ormea, Teodorescu, Midana). Midana obtained negative results in auto inoculation experiments performed on two patients. Teodorescu found in two patients, associated with the epidermodysplasia verruciformis, keratosis palmaris et plantaris hereditaria and regarded both diseases as genodermatoses.

## ACROKERATOSIS VERRUCIFORMIS

(Hopf)

Numerous hyperkeratotic and occasionally verrucous papules are present, predominantly on the dorsa of the hands and the feet.




FIG 18 *Acrokeratosis verruciformis* (Hopf). The lesion consists of a circumscribed area of papillomatous acanthosis and hyperkeratosis ( $\times 100$ )

**Histopathology.** The papules show considerable hyperkeratosis, increase in thickness of the granular layer and acanthosis (Fig 18). In addition, there are slight papillomatosis and some thickening of the rete ridges. The rete ridges all extend to a uniform level and their configuration is well preserved. There is no parakeratosis or

vacuolization of cells such as is seen in verruca vulgaris and verruca plana (Loveman; Niedelman).

### PSEUDOXANTHOMA ELASTICUM

This disorder represents a congenital defect of the elastic tissue which may be limited in extent or widespread. In addition to the elastic tissue in the dermis, the elastic membrane of the retina, called




FIG. 19 Pseudoxanthoma elasticum. Low magnification, Verhoeff stain. This illustration shows the usually observed late degenerative stage. In the lower dermis, the elastic fibers are increased in number, appear swollen and show signs of degeneration, such as fragmentation, disintegration and clumping (Patient's age, 31 years) ( $\times 50$ ).

Bruch's membrane, and the elastic fibers in the aorta, the arteries and the arterioles may be affected. The degeneration of the elastic fibers of Bruch's membrane causes small defects and breaks which manifest themselves clinically as angioid streaks (Urbach and Wolfram, Hagedoorn, Ebert). Involvement of the aorta causes diffuse dilatation (Urbach).

The cutaneous lesions consist of soft, yellowish papules and plaques. The papules frequently show a linear arrangement. The sides of the neck are the most common site of the lesions.

**Histopathology.** Histologic examination of the skin reveals the elastic tissue to be normal in the subepidermal portion of the dermis

In the middle and lower portions of the dermis, however, it is increased, usually in circumscribed areas. Within these areas, the elastic fibers are swollen and show disintegration and clu-  
 eration and imbibition with a basophilic staining and become visible with routine stains. In




FIG. 20 Pseudoxanthoma elasticum. High magnification of figure 19. Verhoeff stain. The elastic fibers are degenerated whereas the collagen shows no evidence of degeneration ( $\times 200$ ).

nerud and Nomland have shown by staining methods (von Kossa's stain for calcium) and Lobitz and Osterberg by micro incineration that the degenerated elastic fibers are richly infiltrated with calcium. The collagen fibers are unaltered.

In a few instances the elastic tissue of the arterioles and the small arteries in the deep dermis was found to have undergone a similar degeneration (Urbach, de Sá Penella and Esteves).

It appears that pseudoxanthoma elasticum occasionally is preceded by an early hyperplastic stage (Fig. 21). In this stage, which seems to be present only during infancy and childhood, the elastic fibers are increased in number and size but show no evidence of degeneration.

as they do in the more commonly observed late degenerative stage (Weidman, Anderson and Ayres, Pautrier)

**Differential Diagnosis.** Senile elastosis, like pseudoxanthoma elasticum, shows a great increase in material taking the elastic tissue stain. However, in senile elastosis this material is located in the upper third



FIG 21 *Pseudoxanthoma elasticum* Low magnification, Verhoeff stain. This illustration shows the early hyperplastic stage. The elastic fibers are large and swollen but show no evidence of degeneration. (Patient's age, 2 years) ( $\times 100$ )

of the dermis and is present as dense masses rather than as individual curls. Furthermore, staining of this material for calcium is always negative in senile elastosis.

### CUTIS HYPERELASTICA (EHLERS-DANLOS SYNDROME)

This syndrome consists of (1) hyperelasticity of the skin, (2) hyperextensibility of the joints, (3) fragility of the skin and the blood vessels with formation of atrophic scars and (4) development of raisin-like pseudotumors. The pseudotumors form at points of trauma and are soft and pigmented and present a wrinkled surface.

In many cases small, hard, subcutaneous nodules have been described. They also are the result of trauma.

The collagen bundles appear atrophic, split up and separated by edema. The elastic fibers are normal in some cases but in most cases show breaking up and clumping. The amount of elastic tissue often appears increased but it is likely that this increase is not a real one but is simulated by the atrophy of the collagen. In addition, the number of capillaries is increased and their lumina dilated. Large, round cystic spaces representing lymphangiectatic cavities may occur (Hortling and Gottron).

The raisinlike pseudotumors that are part of the syndrome form at areas of traumatic hemorrhage and consist either of accumulations of foreign body giant cells (Ronchese) or of proliferated connective tissue with large numbers of vessels in it. The hard subcutaneous nodules contain calcified necrotic fat or mucoid material enclosed in a thick fibrous capsule (Johnson and Falls).

### URTICARIA PIGMENTOSA

This disorder is characterized in most cases by the presence of a great number of brown macules scattered over the entire cutaneous surface. When rubbed with a blunt instrument, they become distinctly urticarial. In rare instances the lesions consist of one or several soft nodules or plaques.

**Histopathology.** The histologic picture shows an infiltrate composed chiefly of mast cells. Mast cells are characterized by the presence of basophilic metachromatic granules in their cytoplasm (see pages 87-88). These granules are not visible on staining with routine stains. Their demonstration requires special staining (see pages 28-29 and Plate I). Fixation in 10 per cent Formalin (without subsequent exposure to Kelly's solution) and staining with Giemsa's stain or with methylene blue are recommended.

In the macular type mast cells are present in the upper third of the dermis especially around the capillaries. Some appear to be round or oval but the majority are spindle-shaped. Since in sections stained with hematoxylin and eosin the mast cells resemble fibroblasts the diagnosis may be missed easily unless special staining is employed.

In the nodular type (Figs. 22-23) mast cells lie closely packed in tumorlike aggregates. The infiltrate may extend through the entire dermis into the subcutis. Wherever the mast cells lie in dense aggregates they are cuboidal rather than spindle shaped and show ample, slightly eosinophilic cytoplasm. Because of their shape and ample



cytoplasm, they look unlike any other cell and the diagnosis can be made without resorting to special staining

If a biopsy is performed shortly after the lesion has been stroked, the section will show edema, an influx of eosinophils and shrinking of the mast cells associated with a great decrease in the amount of




FIG 22 1 • may  
lovely  
hapt  
nification  
picked in  
Because of staining with hematoxylin eosin the granules  
in the mast cells are not visible (X200)

granules within them indicating that the granules have been expelled from the cells (Drennan). There may even be disintegration and a temporary disappearance of mast cells which may be the explanation for certain reports in the literature where in otherwise typical cases of urticaria pigmentosa mast cells have not been found (Drennan and Beare)

The pigmentation of urticaria pigmentosa is due not to the presence of the mast cells but to melanin which is present in increased amounts in the basal layer and occasionally also in melanophores

The presence of extracutaneous lesions in urticaria pigmentosa has been reported recently. However, the systemic lesions in Ellis'

case may well have been lymphoma, and the bone lesions in Sagher's and Clyman's cases were not examined histologically




FIG 23 Urticaria pigmentosa, nodular type. High magnification of Figure 22, hematoxylin-eosin stain. The mast cells appear as large cuboidal cells ( $\times 400$ )

### INCONTINENTIA PIGMENTI

The disease frequently begins immediately after birth with inflammatory lesions, particularly bullae, in linear and grouped arrangement. The bullae may recur for months. They finally give way to areas of pigmentation. There may be an intermediary stage of linear verrucous lesions. The pigmented areas are widely disseminated, located especially on the trunk, and have an irregular, bizarre outline.

**Histopathology.** In the early vesicular stage, one observes intra-epidermally located vesicles containing many eosinophils (Carney, Epstein, Vedder and Pinkus). In addition, there is spongiosis of the epidermis, and in the dermis an infiltrate composed of lymphocytes and eosinophils.

sp

The final stage shows diminution or absence of melanin in the basal layer of the epidermis and extensive deposits of melanin inside

cytoplasm, they look unlike any other cell and the diagnosis can be made without resorting to special staining

If a biopsy is performed shortly after the lesion has been stroked, the section will show edema, an influx of eosinophils and shrinking of the mast cells associated with a great decrease in the amount of




FIG 22 Urticaria pigmentosa, nodular type Low magnification hematoxylin eosin stain Mast cells lie closely packed in the upper dermis They are cuboidal in shape Because of staining with hematoxylin eosin the granules in the mast cells are not visible ( $\times 200$ )

granules within them indicating that the granules have been expelled from the cells (Drennan) There may even be disintegration and a temporary disappearance of mast cells which may be the explanation for certain reports in the literature where in otherwise typical cases of urticaria pigmentosa mast cells have not been found (Drennan and Beare)

The pigmentation of urticaria pigmentosa is due not to the presence of the mast cells but to melanin which is present in increased amounts in the basal layer and occasionally also in melanophores

The presence of extracutaneous lesions in urticaria pigmentosa has been reported recently However, the systemic lesions in Ellis

- Montgomery H and Reuter M J Xeroderma pigmentosum Arch Dermat & Syph 26 256 1932

### Congenital Ectodermal Defect

- Sunderman F W Persons lacking sweat glands Arch Int Med 67 816 1941  
 Upshaw B Y and Montgomery H Hereditary anhidrotic ectodermal dysplasia Arch Dermat & Syph 60 1170 1949

### Rothmund's Syndrome Werner's Syndrome and Progeria of Children

- Reed R Seville R H and Tattersall R N Werner's syndrome Brit J Dermat 65 165 1953  
 Jannhäuser S J Werner's syndrome (progeria of the adult) and Rothmund's syndrome two types of closely related hereditary familial atrophic dermatoses with juvenile cataracts and endocrine features Ann Int Med 23 559 1945

### Hydroa Vacciniforme and Hydroa Aestivale

- Senear F E and Fink H W Hydroa vacciniforme seu aestivale Arch Dermat & Syph 7 143 1923

### Epidermolysis Bullosa

- Allen A C Survey of pathologic studies of cutaneous diseases during World War II Arch Dermat & Syph 57 19 1943  
 Engman M F and Mook W H A study of some cases of epidermolysis bullosa with remarks upon the congenital absence of elastic tissue J Cutan Dis 24 30 1905  
 Johnson S A M and Fest A R Epidermolysis bullosa simplex of the hands and feet Arch Dermat & Syph 53 610 1916  
 Lamb J H and Halpert B Epidermolysis bullosa of the newborn Arch Dermat & Syph 55 369 1917  
 Leonis A Recherches sur le mécanisme de formation des bulles dans l'épidermolyse bulleuse simple Ann dermat et syph 10 501 1900  
 Schiffer G Two cases of epidermolysis bullosa hereditaria letalis Acta dermat venereol 31 701 19 1  
 Tulipan L Epidermolysis bullosa dystrophica (recessive type) Arch Dermat & Syph 37 70 1938

### Darier's Disease

- Beerman H Hypertrophic Darier's disease and nevus syringocystadenomatosus papilliferus Arch Dermat & Syph 60 500 1949  
 Brunauer S R Ueber Schleimhautveränderungen bei Morbus Darier und über die Pathogenese dieser Erkrankung Acta dermat venereol 6 131 1925  
 Ellis F A Keratosis follicularis is not primarily a follicular disease Arch Dermat & Syph 50 27 1944  
 Pels T R and Goodman M H Criteria for the histologic diagnosis of keratosis follicularis (Darier) Arch Dermat & Syph 39 438 1939

### Familial Benign Chronic Pemphigus

- Ellis F A Vesicular Darier's disease (so called benign familial pemphigus) Arch. Dermat. & Syph 61 715 1950  
 Finnerud C W and Seymanick F J Chronic benign familial pemphigus a possible vesicular variant of keratosis follicularis Arch. Dermat. & Syph 61 737 1950

and outside of melanophores in the upper dermis. In many cases the basal cells show degeneration and vacuolization. It is believed generally that the disease causes damage to the cells in the basal layer so that the melanocytes become incapable of holding and metabolizing melanin (Sulzberger, Doornink).

## BIBLIOGRAPHY

## Ichthyosis

Ingman A. Studien über Ichthyosis congenita s. foetalis. Acta dermat. venercol. 5:125 1924

Klinger L. B. Ichthyosis congenita. Arch. Dermat. & Syph. 13:91 1926

## Congenital Ichthyosiform Erythroderma

Barker L. P. and Sachs W. Bullous congenital ichthyosiform erythroderma. Arch. Dermat. & Syph. 67:443 1953

Filsh F. in discussion of Barker L. P. and Sachs W. (see above)

Lawless T. K. Congenital ichthyosiform erythroderma. Arch. Dermat. & Syph. 44:30 1941

Laymon C. W. and Murphy R. Congenital ichthyosiform erythroderma. Arch. Dermat. & Syph. 57:615 1948

## Keratosis Punctata Palmaris et Plantaris

Scott M. J., Costello M. J. and Simuangco S. Keratosis punctata palmaris et plantaris. Arch. Dermat. & Syph. 64:301 1951

## Pachyonychia Congenita

Andrews G. C. Pachyonychia congenita. Arch. Dermat. & Syph. 33:183 1936

Cole H. N., Rauschkopf J. E. and Toomey J. Dyskeratosis congenita with pigmentation, dystrophia unguium and leukoplakia oris. Arch. Dermat. & Syph. 21:71 1930

Diasio F. A. Pachyonychia congenita Jadassohn. Arch. Dermat. & Syph. 30:218 1934

Garb J. and Rubin G. Dyskeratosis congenita. Arch. Dermat. & Syph. 50:191 1944

Wright C. S. and Guequierre J. P. Pachyonychia congenita. Arch. Dermat. & Syph. 55:819 1917

## Porokeratosis Mibelli

Ambler J. V. and Stout R. L. Porokeratosis (Mibelli). Arch. Dermat. & Syph. 29:20 1934

Bloom D. and Abramowitz E. W. Porokeratosis Mibelli. Arch. Dermat. & Syph. 47:1 1943 (Good review)

Ritchie E. B. and Becker S. W. Porokeratosis (Mibelli). Arch. Dermat. & Syph. 26:1032 1932

## Xeroderma Pigmentosum

Becker S. W., Pautrier L. M. and Woringer F. A case of xeroderma pigmentosum. Arch. Dermat. & Syph. 25:915 1932

Bell E. T. and Rothnem T. P. Xeroderma pigmentosum with carcinoma of the lower lip in two brothers aged 16 and 13 years. Am. J. Cancer 30:574 1937

King H. and Hamilton C. M. Xeroderma pigmentosum in a Negress. Arch. Dermat. & Syph. 42:570 1940

- Johnson S A M and Falls H F Ehlers-Danlos syndrome Arch Dermat & Syph 60 82 1949  
 Korting G W and Gottron E Cutis laxa Arch f Dermat u Syph 193 14 1951 (Good review)  
 Ronchese F Dermatorthexis Am J Dis Child 51 1403 1936

### Urticaria Pigmentosa

(See also Chapter 5 Mast Cells page 38)

- Clyman S G and Rein C R Urticaria pigmentosa associated with bone lesions J Invest Dermat 19 179 1952  
 Drennan J M The mast cells in urticaria pigmentosa J Path & Bact 63 513 1951  
 — and Beare J M Investigations in a case of urticaria pigmentosa Brit J Dermat 63 257 1951  
 Ellis J M Urticaria pigmentosa Arch Path 48 126 1949  
 Finnerud C W Urticaria pigmentosa (nodular type) with a summary of the literature Arch Dermat & Syph 8 341 1923  
 Fraser J F and Richter M N Urticaria pigmentosa Arch Dermat. & Syph 17 489 1923  
 Sagher F Cohen C and Schorr S Concomitant bone changes in urticaria pigmentosa J Invest Dermat 18 425 1952

### Incontinentia Pigmenti

- Carney R G Incontinentia pigmenti Arch Dermat & Syph 64 126, 1951  
 Doornink F J Über Incontinentia pigmenti und über die Siemens Bloch'sche Pigmentdermatose Dermatol 102 63 1951  
 Epstein S Vedder J S and Pinkus H Bullous variety of incontinentia pigmenti (Bloch Sulzberger) Arch Dermat & Syph 65 557 1950  
 Sulzberger M B Ueber eine bisher nicht beschriebene „congenital“ Pigmentanomalie (Incontinentia pigmenti) Arch f Dermat u Syph 154 19 1928  
 — — Incontinentia pigmenti (Bloch Sulzberger) Arch Dermat & Syph 38 57 1938

## 62 Congenital Diseases (*Genodermatoses*)

- Frank S B and Rein C R Dyskeratoid dermatosis Arch Dermat & Syph 45 129 1912
- Hailey H and Hailey H Familial benign chronic pemphigus Arch Dermat & Syph 39 679 1939
- Winer L H and Leeb A J Benign familial pemphigus Arch Dermat & Syph 67 77, 1953 (Good discussion)

### Epidermodysplasia Verruciformis

- Costa O G and Junqueira M A Epidermodysplasia verruciformis (Lewandowsky and Lutz) Arch Dermat & Syph 46 469 1912
- Lutz W A A propos de l'épidermodysplasie verruciforme Dermatol 92 30 1916
- Midani A Sulla questione dei rapporti tra epidermodysplasia verruciformis e verrucosi generalizzata Dermatol 99 1 1919
- Ormea F Epidermodysplasia verruciformis und Hautcarcinom Arch f Dermat u Syph 188 278 1919
- Sullivan M and Ellis F A Epidermodysplasia verruciformis Arch Dermat & Syph 40 122 1939
- Tcodorescu S Teller M and Conu A Zieci f die von Epidermodysplasia verruciformis Lewandowsky Lutz Arch f Dermat u Syph 188 123 1919
- Waisman M and Montgomery H Verruca plana and epithelial nevus Arch Dermat & Syph 45 259 1912

### Acrokeratosis Verruciformis

- Lovenheim A B Acrokeratosis verruciformis (Hopf) Arch Dermat & Syph 43 971 1911
- Niedelman M L Acrokeratosis verruciformis (Hopf) Arch Dermat & Syph 56 18 1917 (Lit)

### Pseudoxanthoma Elasticum

- Libert R H Angioid streaks and pseudoxanthoma elasticum Arch Dermat & Syph 48 75 1913
- Finnerud C W and Nomland R Pseudoxanthoma elasticum Arch Dermat & Syph 35 653 1937
- Hagedoorn A Angioid streaks Arch Ophth 21 716 935 1939
- Iobitz W and Osterberg A E Pseudoxanthoma elasticum microincineration J Invest Dermat 15 297 1950
- Pautrier L M A propos de deux affections d'élastine du revêtement cutané témoins d'affections généralisées de l'élastine de tout l'organisme Arch belg de dermat et de syph 4 259 1918
- de Sá Penella L and Esteves J Pseudoxanthoma elasticum Arch Dermat & Syph 45 283 1912
- Urbach E and Wolfram S Ueber Veränderungen des elastischen Gewebes bei einem autopsisch untersuchten Falle von Groenblad Strandbergschem Syndrom Arch f Dermat u Syph 176 167 1937
- Weidman F D Anderson N P and Ayres S Juvenile elastoma Arch Dermat & Syph 28 182 1933

### Ehlers Danlos Syndrome

- Burrows A Cutis hyperelastica (Ehlers Danlos syndrome) Brit J Dermat 50 648 1938

TABLE 2—CLASSIFICATION OF VESICLES AND BULLAE

TYPE OF VESICLE OR BULLA	MODE OF FORMATION	<i>Site</i> SITE OF FORMATION	DISEASES
Subcorneal bulla	Detachment of horny layer	Subcorneal	Impetigo
Spongione bulla	Cellular necrosis followed by spongiosis and, occasionally, by reticular degeneration	Intra epidermal	Dermatitis-Eczema Pompholyx Congenital ichthyosisiform erythroderma Hydroa
Milium bulla	Retention of sweat	Intracorneal Intra-epidermal or subepidermal	Milium crystallina Milium rubra
Acantholytic bulla	Acantholysis	Intra epidermal a suprabasal	Pemphigus vulgaris Pemphigus vegetans Familial benign chronic pemphigus Darier's disease Senile keratosis
		b upper epidermis	Pemphigus foliaceus Pemphigus erythematosus
5 Viral bulla	Reticular and ballooning degeneration with acantholysis	Intra-epidermal	Varicella Herpes simplex Herpes zoster Varicella
6 Pressure bulla	Detachment of entire epidermis	Subepidermal	Bullous pemphigoid Benign mucous membrane pemphigoid Dermatitis herpetiformis Erythema multiforme Epidermolysis bullosa Porphyria cutanea tarda
7 Bulla due to basal cell degeneration	Degeneration of basal cells	Subepidermal	Incontinentia pigmenti Lichen planus Lichen sclerosus et atrophicus Lupus erythematosus
8 Bulla due to reticulum degeneration	Degeneration of sub-epidermal reticulum fibers	Subepidermal	Burns



# 7

## Noninfectious Vesicular and Bullous Diseases

Several new concepts have been introduced in recent years concerning the histologic structure of vesicles and bullae and their mechanism of formation. It appears appropriate therefore to present a classification of the different types of vesicles and bullae and to outline briefly their mode of formation before discussing individual diseases. (Since from a histologic point of view it is immaterial whether a lesion is a vesicle or a bulla only the latter term will be used in the following presentation.)

Eight distinct types of bullae can be recognized (Table 2)

1 *Subcorneal bulla* Detachment of the horny layer from the stratum malpighii occurs

2 *Spongiotic bulla* Liquefaction necrosis of a few epidermal cells causes formation of a minute cavity (*vesiculette primordiale*) which subsequently due to intercellular edema (spongiosis) and intracellular edema (*alteration cavitaire*) in the surrounding epidermis enlarges into an intra epidermal bulla. In the case of pronounced intracellular edema reticular degeneration of the epidermis may occur (For a detailed description see page 69)

3 *Miliarial bulla* In miliaria bullae form due to the escape of sweat from the sweat ducts. Two types occur *miliaria crystallina* in which the bulla is located within the stratum corneum and *miliaria rubra* in which the bulla forms either within the stratum malpighii or beneath the epidermis. In severe cases of miliaria rubra the sweat duct ruptures (For a detailed description see page 76)

4 *Acantholytic bulla* Degeneration of intercellular bridges causes loss of coherence between epidermal cells and formation of rifts which enlarge into bullae. Detached (acantholytic) epidermal cells are present in the bulla cavity. Acantholysis may take place within the lower epidermis predominantly right above the basal layer or within the upper epidermis predominantly within the granular layer (For a detailed description see pages

TABLE 2—CLASSIFICATION OF VESICLES AND BULLAE

TYPE OF VESICLE OR BULLA	MODE OF FORMATION	<i>Site</i> SITE OF FORMATION	DISEASES
Subcorneal bulla	Detachment of horny layer	Subcorneal	Impetigo
Spongiotic bulla	Cellular necrosis followed by spongiosis and, occasionally, by reticular degeneration	Intra-epidermal	Dermatitis-Eczema Pempholyx Congenital ichthyosiform erythroderma Hydroa
Miliarial bulla	Retention of sweat	Intracorneal Intra epidermal or subepidermal	Miliaria crystallina Miliaria rubra
Acantholytic bulla	Acantholysis	Intra-epidermal & suprabasal	Pemphigus vulgaris Pemphigus vegetans Familial benign chronic pemphigus Darnier's disease Senile keratosis
		to upper epidermis	Pemphigus foliaceus Pemphigus erythematosus
1 Viral bulla	Reticular and ballooning degeneration with acantholysis	Intra epidermal	Varicella Herpes simplex Herpes zoster Varicella
5 Pressure bulla	Detachment of entire epidermis	Subepidermal	Bullous pemphigoid Benign mucous-membrane pemphigoid Dermatitis herpetiformis Erythema multiforme Epidermolysis bullosa Porphyria cutanea tarda
7 Bulla due to basal cell degeneration	Degeneration of basal cells	Subepidermal	Incontinentia pigmenti Lichen planus Lichen sclerosus et atrophicus Lupus erythematosus
8 Bulla due to reticulum degeneration	Degeneration of sub-epidermal reticulum fibers	Subepidermal	Burns

5 *Irral bulla* Invasion of epidermal cells by certain viruses causes two types of degenerative changes in epidermal cells ballooning and reticular degeneration Ballooning degeneration leads to extensive acantholysis Although the bullae form within the epidermis progression of the ballooning degeneration leads to a subepidermal location in older bullae (For a detailed description see page 239)

6 *Pressure bulla* Detachment of the entire epidermis from the dermis occurs Pressure of the accumulating tissue fluid often causes in early bullae *rounding of the lateral walls and stretching of the* epidermal cells located there Older bullae due to regeneration of the epidermis at the floor of the bulla may have an intraepidermal location If in addition disintegration of the detached stratum malpighii occurs the bulla may be subcorneal in location (For a detailed description see page 86)

7 *Bulla due to basal cell degeneration* Several diseases (see Table 2) cause hydropic degeneration of the basal cells This may result in damage to the cytoplasmic processes which effect the coherence of the basal cell layer with the dermis Thus a subepidermal bulla may form (For a detailed description see page 296)

8 *Bulla due to reticulum degeneration* Damage to the subepidermal feltwork of reticulum fibers causes the cytoplasmic processes of basal cells to pull out of this feltwork A subepidermal bulla forms into which numerous cytoplasmic processes extend from the basal cells at the roof of the bulla (For a detailed description see page 90)

## DERMATITIS ECZEMA

The terms dermatitis and eczema now are used generally as synonyms They refer to an inflammation of the skin based on an allergic response of the skin to a variety of agents such as chemicals proteins bacteria and fungi The exciting allergen may act on the skin either from the outside or from the inside

Dermatitis or eczema may be acute subacute or chronic The clinical picture is characterized by polymorphism of the eruption Among the primary lesions that may be observed are macules papules and vesicles the macules and papules tend to coalesce to form areas of diffuse erythema Among the secondary lesions are scaling crusting lichenification and fissuring The lesions of dermatitis usually are not demarcated sharply but merge gradually into the surrounding normal skin Moderately severe to severe itching is present in most forms of dermatitis

No generally accepted classification of dermatitis exists and many cases defy assignment to any definite type In this section the following types of dermatitis will be discussed (1) *contact dermatitis*

(2) nummular eczema (3) atopic dermatitis (or neurodermatitis disseminata) (4) lichen simplex chronicus (or neurodermatitis circumscripta) (5) exudative discoid and lichenoid chronic dermatosis (6) seborrheic dermatitis including Lerner's disease (7) stasis dermatitis and (8) generalized exfoliative dermatitis. In addition lesions of dermatitis may occur in superficial fungus infections as drug eruptions and in lymphoma. The latter three forms of dermatitis will be discussed when the respective diseases are described.

*Contact dermatitis* is caused by contact of the skin with an agent that acts either as a specific allergic sensitizer or as a primary irritant. Contact dermatitis may be acute, subacute or chronic. In acute and subacute contact dermatitis diffuse erythema, edema, oozing and crusting predominate. In addition vesicles are often present if a specific allergic sensitizer is the cause. In chronic contact dermatitis erythema, scaling and lichenification prevail.

*Nummular eczema* probably caused by a temporary loss of resistance in the skin to the ordinary bacterial flora of the skin presents fairly sharply demarcated patches of erythema studded with discrete pinpoint vesicles or pinpoint erosions.

*Atopic dermatitis* a constitutional familial dermatosis of unknown cause which often is aggravated by emotional tension or allergic factors shows lichenified and scaling erythematous areas which when active show also oozing and crusting but no vesicles.

*Lichen simplex chronicus* shows one or several lichenified plaques with little scaling. Oozing and vesiculation are absent. On the lower legs especially lichen simplex chronicus may become hypertrophic and assume a verrucous nodular appearance (lichenificatio gigantea lichen cornuus hypertrophicus).

*Exudative discoid and lichenoid chronic dermatosis* shows small lesions with central crusting and later sent an erythematous border.

*Seborrheic dermatitis* shows fairly sharply demarcated brownish red areas which show only little infiltration and are covered with fine greasy scales. Oozing may be present but no vesiculation is found. Generalized seborrheic dermatitis in infants often is referred to as Lerner's disease.

*Stasis dermatitis* presents erythema, edema, scaling and occasionally oozing and crusting. It differs from other forms of dermatitis first by showing brownish pigmentation due to hemosiderin deposits and second by its location on the lower legs.

skin

... scaling and in severe cases

oozing. It represents a peak reaction to which several forms of dermatitis may lead—for instance, contact dermatitis, atopic dermatitis, seborrheic dermatitis, stasis dermatitis, drug dermatitis and lymphoma dermatitis. However, it may occur as an idiopathic disease.

**Histopathology.** The various types of dermatitis rarely present a histologic picture sufficiently diagnostic to allow their differentiation



FIG. 24 Acute dermatitis contact dermatitis due to poison ivy. Numerous intraepidermally located vesicles and marked intracellular edema are present. The vesicles are separated by thin septa formed by the resisting walls of edematous epidermal cells (reticular degeneration) and thus form a multilocular bulla ( $\times 100$ )

because the same histologic reactions occur in all forms of dermatitis: exudation leading to vesiculation in the acute stage, proliferation leading to acanthosis in the chronic stage, and a combination of these two reactions in the subacute stage. Since, as a rule, no more specific diagnosis than acute, subacute or chronic dermatitis can be made, the histologic picture as presented by an acute, subacute and chronic dermatitis will be described first. Thereafter, the distinctive features occasionally presented by the various members of the dermatitis eczema group will be discussed.

IN ACUTE DERMATITIS, intra epidermally located vesicles or bullae predominate the histologic picture. Considerable intercellular edema (spongiosis) and intracellular edema (altération cavitaire) may be present in the epidermis surrounding the vesicles. If the number of

vesicles is great and the intracellular edema pronounced the vesicles due to reticular degeneration of the epidermis will be separated from one another only by thin septa formed by the resisting walls of edematous epidermal cells and will thus form a multilocular bulla (Fig 24) The vesicles and the bullae contain a few lymphocytes eosinophils and neutrophils and disintegrated epidermal cells Migrating lymphocytes and neutrophils are present in the epidermis The cells of the stratum corneum may be parakeratotic and intermingled with fibrin and numerous neutrophils (Winer and Lap-  
 ————— vascular dilatation edema and a

neutrophils  
 ————— formation of the vesicles and bullae

" had been  
 the vesicles  
 necrosis is

the primary factor and that in 1925 Civatte described as the primary lesion the vesiculette primordiale formed by the lysis of two or three squamous cells through cyto-  
 ————— alteration Spongiosis followed this and caused enlargement  
 ————— expressed the same  
 support vesicles tend

to form in areas where the rete ridges are deepified vesicles often lie in areas formerly occupied by epidermal cells (Fig 25) and spongiosis may be entirely absent in the vicinity of vesicles Polak and Mom studying the formation of vesicles in experimental eczema with the aid of silver impregnation found that at the time the vesiculette primordiale formed the intercellular bridges were still intact and only broke later by mechanical force when the vesicle had increased in size

✓ Differentiation of acute vesicular or bullous dermatitis from bullous erythema multiforme and dermatitis herpetiformis is not always possible Although in the latter two diseases the bullae form subepi-  
 dermally they may be located intra epidermally during the stage of healing due to regeneration of the epidermis In that case secondary findings such as the presence of marked spongiosis in contact der-  
 matitis and its absence in the other two diseases and the number of  
 ↑ eosinophils may aid in the decision It should be stressed as of utmost importance that for the diagnosis of all vesicular and bullous dis-  
 eases an early lesion must be chosen for histologic examination be-  
 cause secondary factors such as regeneration and pyogenic infection  
 may obscure the diagnostic features

oozing. It represents a peak reaction to which several forms of dermatitis may lead—for instance contact dermatitis atopic dermatitis seborrheic dermatitis stasis dermatitis drug dermatitis and lymphoma dermatitis. However it may occur as an idiopathic disease.

**Histopathology** The various types of dermatitis rarely present a histologic picture sufficiently diagnostic to allow their differentiation.




Fig. 24 Acute dermatitis contact dermatitis due to poison ivy. Numerous intraepidermally located vesicles and marked intracellular edema are present. The vesicles are separated by thin septa formed by the resisting walls of edematous epidermal cells (reticular degeneration) and thus form a multilocular bulla ( $\times 100$ ).

because the same histologic reactions occur in all forms of dermatitis exudation leading to vesiculation in the acute stage proliferation leading to xanthosis in the chronic stage and a combination of these two reactions in the subacute stage. Since, as a rule no more specific diagnosis than acute subacute or chronic dermatitis can be made the histologic picture is presented by an acute subacute and chronic dermatitis will be described first. Thereafter the distinctive features occasionally presented by the various members of the dermatitis eczema group will be discussed.

In ACUTE DERMATITIS intraepidermally located vesicles or bullae predominate the histologic picture. Considerable intercellular edema (spongiosis) and intracellular edema (alteration cavitaire) may be present in the epidermis surrounding the vesicles. If the number of

Sachs, Muller and Gray (1916) to the histologic picture as described above for chronic dermatitis because it is characteristically present in neurodermatitis disseminata (atopic dermatitis) and neurodermatitis circumscripta (lichen simplex chronicus). However the same histologic picture may occur in any dermatosis belonging to the dermatitis eczema group.




Fig. 96 Chronic dermatitis neurodermatitis circumscripta. There are hyperkeratosis, acanthosis, elongation of the rete ridges and elongation and broadening of the papillae. The dermis shows a chronic inflammatory infiltrate and fibrosis ( $\times 50$ ).

Many diseases not members of the dermatitis eczema group show either regularly or occasionally a histologic picture allowing no more specific diagnosis than chronic dermatitis. Diseases which regularly show the nonspecific histologic picture of chronic dermatitis include psoriasis, psoriasis rosea, parapsoriasis and pellagra. Many other diseases such as psoriasis, lichen planus and lupus erythematosus to name but a few show a diagnostic histologic picture in clinically typical cases but may show a nonspecific histologic picture that of chronic dermatitis in clinically atypical cases.

Trichomycosis fungoides must always be kept in mind as a possible diagnosis when a section showing chronic dermatitis is examined. It often is a very difficult task to establish or rule out early mycosis



IN SUBACUTE DERMATITIS, one sees spongiosis, intracellular edema and, usually, also vesicle formation. However, the vesicles are smaller than in acute dermatitis (Fig 25). Moderate acanthosis and varying degrees of parakeratosis are present. The inflammatory infiltrate in the dermis usually is pronounced and is composed of a multiplicity of cells. Lymphocytes predominate, but neutrophils, eosinophils and




FIG 25 Subacute dermatitis nummular eczema. There is intraepidermal vesicle formation. The vesicles lie in areas formerly occupied by epidermal cells. The epidermis shows parakeratosis and moderate acanthosis. The dermis shows a perivascular infiltrate ( $\times 100$ ).

histiocytes are also seen. There may be considerable migration of neutrophils and lymphocytes through the epidermis.

IN CHRONIC DERMATITIS, there is often marked acanthosis with elongation of the rete ridges. There is hyperkeratosis intermingled with areas of parakeratosis. Slight intercellular edema may be present in the epidermis, but vesicle formation is absent. In the upper dermis, one sees a moderate amount of predominantly perivascular infiltration composed of various types of cells. Lymphocytes prevail, but the number of eosinophils, histiocytes and fibroblasts may be considerable. Neutrophils are absent. The number of capillaries is increased and the walls of the arterioles and the small arteries may be thickened. The term neurodermatitic reaction has been given by

that of psoriasis which shares with lichen simplex chronicus the tendency to elongation of the rete ridges. However, psoriasis shows predominantly parakeratosis rather than hyperkeratosis thinning of the suprapapillary portions of the stratum malpighii edema of the upper portions of the papillae and not infrequently Munro micro abscesses. Furthermore the papillary capillaries in psoriasis are dilated and tortuous whereas in lichen simplex chronicus they appear normal. According to Stoughton and Wells the Hetchkiss McManus stain demonstrates the capillary changes in psoriasis very clearly and thus aids in its differentiation from lichen simplex chronicus.

**EXUDATIVE DISCOID AND LICHENOID CHRONIC DERMATOSIS** (Sulzberger Garbe) The epidermis may show spongiosis and vesiculation as in subacute dermatitis or acanthosis with little or no edema as in chronic dermatitis. The vessels of the upper dermis and the mid dermis are dilated and their walls thickened. About them is a mantle of lymphocytes histiocytes polymorphonuclear neutrophils eosinophils and numerous plasma cells. Sachs and Kirsch state that the presence of many plasma cells gives the histologic picture a distinctive appearance so that the diagnosis can be established on microscopic findings alone.

**SEBORRHEIC DERMATITIS** The histologic picture is not diagnostic. It may be said to be halfway between psoriasis and chronic dermatitis. The horny layer because of the tendency to desquamation is only poorly developed and most of its cells are parakeratotic. The epidermis shows slight to moderate acanthosis with elongation of the rete ridges and slight intracellular edema and spongiosis. The dermis shows a mild chronic inflammatory infiltrate. Munro micro abscesses and neutrophils migrating through the epidermis as seen in psoriasis occasionally are observed. In cases in which the histologic picture resembles that of psoriasis the presence of spongiotic areas

in venous Older lesions may show numerous newly formed capillaries embedded in a fibrotic dermis. Whereas it has been assumed generally that venous and capillary stasis is responsible for the clinical manifestations of stasis dermatitis Kulwin and Hines in a study of the vessels at the dermo-subcutaneous junction found changes more often and more severely in the arterioles than in the venules. The changes in both arterioles and venules consisted of intimal proliferation endothelial hyperplasia and medial hypertrophy. Complete obliteration of arterioles

fungoides One should search for atypical histiocytes (so-called mycosis cells) mitotic figures clumping of nuclei karyorrhexis (disintegration of nuclei into nuclear dust) and Pautrier micro abscesses (For further details see page 485) However it should be realized that some atypicality of the histiocytes and an occasional mitotic figure sometimes may be seen in chronic dermatitis If in doubt, it is always best to request another specimen for histologic examination

A few words about the histologic aspects of the various members of the dermatitis eczema group are now in order

**CONTACT DERMATITIS** Contact dermatitis may be acute subacute or chronic The histologic descriptions given above for acute subacute and chronic dermatitis apply in general to contact dermatitis Acute contact dermatitis presents numerous closely set large and small intra epidermal vesicles (Fig 24) Chronic contact dermatitis shows irregular acanthosis Even at that stage spongiosis and small intra epidermal vesicles are often present (Sachs Miller and Gray 1944 Miller)

**NUMMULAR ECZEMA** (infectious eczematoid dermatitis) This eruption characterized clinically by pinpoint vesicles usually shows histologically a picture of subacute dermatitis (Fig 25) In a moderately acanthotic epidermis one finds numerous scattered areas of intra epidermal vesiculation (Sachs Miller and Gray 1946) As a rule spongiosis about the vesicles is slight or absent

**ATOIC DERMATITIS** (neurodermatitis disseminata) The histologic picture is usually that of a chronic dermatitis showing acanthosis with varying degrees of spongiosis The number of eosinophils in the inflammatory infiltrate is often considerable (Burkhart and Montgomery)

**LICHEN SIMPLEX CHRONICUS** (neurodermatitis circumscripta) The microscopic appearance is essentially that of a chronic dermatitis (Fig 26) There are hyperkeratosis interspersed with small areas of parakeratosis acanthosis characterized by rather regular elongation of the rete ridges and elongation and broadening of the papillae There may be some spongiosis but vesiculation does not occur In addition to a chronic inflammatory infiltrate the dermis often shows a fair number of fibroblasts and some fibrosis even in the papillae

In the hypertrophic type of lichen simplex chronicus (lichenification gigantea or lichen cornuus hypertrophicus) the epidermis shows in addition to acanthosis with elongation of the rete ridges considerable hyperkeratosis and papillomatosis (Shaffer and Beerman Hyman and Erger)

The histologic picture of lichen simplex chronicus may resemble

tion at first shows no evidence of lymphomatous infiltration in the dermis it is advisable to perform further biopsies at intervals (See also page 490)

### DERMATOPATHIC LYMPHADENITIS

Any extensive dermatitis but particularly generalized exfoliative dermatitis whether due to lymphoma or not may cause a generalized lymphadenitis of the subcutaneous lymph nodes

**Histopathology** This lymphadenitis has certain characteristic histologic features not found in other types It was described first by Pautrier and Woringer as hypo-melanotic reticulosis Hurwitz intro-

duces the term lymphadenitis centers The pulp of the lymph node shows considerable hyperplasia of reticulum cells Since the reticulum cells possess abundant and faintly eosinophilic cytoplasm the areas of reticular hyperplasia appear as large pale patches (Lapilly) The reticulum cells show phagocytic activity and may contain hemosiderin melanin and, occasionally, fat The lymph follicles as well as the pulp of the lymph node are permeated with eosinophils neutrophils and plasma cells The intermediary sinuses are filled with reticulum cells (sinus catarrhi) The melanin and the fat occasionally present in the lymph nodes originate in the skin and are carried into the lymphatics by scratching It has been suggested that the fat may not be sebum but ointment base (Bettley)

**Differential Diagnosis** These histologic changes differ from those observed in mycosis fungoides Hodgkin's disease and follicular lymphoma by the absence of destruction of the basic architecture of the lymph node the absence of Sternberg Reed cells and the presence of phagocytic activity in the reticulum cells (Hurwitz) The large lymph follicles present in dermatopathic lymphadenitis differ from those of follicular lymphoma by greater uniformity in size smaller number and absence of fissures which frequently separate the lymph follicles from the stroma in follicular lymphoma

In recent years the relationship of dermatopathic lymphadenitis to lymphoma has been discussed by several authors Some have assumed that a dermatopathic lymphadenitis can develop into lymphoma (Bluefarb and Webster) and others have regarded cases of generalized erythroderma with dermatopathic lymphadenitis as Brill Symmers disease (i.e. follicular lymphoma) even in the absence of the histologic criteria of follicular lymphoma (Rost) Neither point of view is justified Dermatopathic lymphadenitis as such is an en-

was seen in specimens from ulcerated areas. These authors concluded that arteriolar changes may be an important etiologic factor in stasis dermatitis.

**GENERALIZED EXFOLIATIVE DERMATITIS** The histologic appearance may be that of a subacute or a chronic dermatitis. In the subacute type there is parakeratosis, marked intercellular and intracellular



FIG. 27 Generalized exfoliative dermatitis due to arsphenamine. There is parakeratosis, marked intercellular and intracellular edema in the upper stratum malpighii, acanthosis with elongation of the rete ridges and migration of inflammatory cells through the epidermis. The upper dermis shows edema and a considerable amount of inflammatory infiltrate ( $\times 100$ ).

edema, particularly in the upper stratum malpighii, acanthosis with elongation of the rete ridges and migration of cells through the epidermis. The upper dermis shows edema and a considerable amount of inflammatory infiltrate (Fig. 27). If the edema in the upper stratum malpighii is pronounced, the cells of the upper stratum may exfoliate together with the parakeratotic horny cells. In the chronic type of exfoliative dermatitis, the histologic picture is that of chronic dermatitis. Each such case requires thorough histologic investigation in order to rule out lymphoma. Montgomery has stated that 25 per cent of all cases of exfoliative dermatitis are proved on histologic examination to be associated with lymphoma. Even if the histologic examina-

Tranck A. Le cytodagnostic immédiat en dermatologie Ann dermat et syph 8 205 1948

#### **Dermatitis Herpetiformis**

Burkhardt R. J and Montgomery H Dermatologic significance of tissue eosinophilia Arch Dermat & Syph 49 19 1944

Civate A Diagnostic histopathologique de la dermatite polymorphe douloureuse ou maladie de Duhring Brocq Ann dermat et syph 3 1 1943

Lapière M A propos du diagnostic différentiel du pemphigus chronique vulgaire et de la dermatite polymorphe de Duhring Brocq Arch belges dermat et syph 1 216 1939

#### **Erythema Multiforme**

Costello M J Erythema multiforme exudativum J Invest Dermat 8 127 1947

Winer L H and Lipschultz C E Comparative study of histology and cytology in vesiculating eruptions Arch Dermat & Syph 65 270 1952

#### **Burn**

Moritz A R Studies of thermal injury III The pathology and pathogenesis of cutaneous burns Am J Path 23 915 1947

turely nonspecific reaction to an extensive dermatitis. Since lymphoma may manifest itself as an extensive dermatitis (see page 490), it can cause a dermatopathic lymphadenitis. Subsequently, the lymphoma may extend to lymph nodes previously affected by dermatopathic lymphadenitis, but in that case the latter is only chronologically—not etiologically—the forerunner of the lymphoma (Jarrett and Kelleit; Keller and Staemmler).

### MILIARIA

Miliaria occurs following excessive sweating in parts of the body covered by clothing. There are two types: *miliaria crystallina* and *miliaria rubra*. In *miliaria crystallina*, asymptomatic, small, superficial, noninflammatory vesicles are present. In *miliaria rubra*, the lesions consist of pruritic, discrete but closely aggregated papules, papulovesicles and vesicles surrounded by erythema.

The cause of miliaria lies in excessive hydration of the horny layer by sweat. This results in swelling of the keratin, closure of the narrow sweat pores by keratin and retention of sweat in the sweat ducts.

**Histopathology.** In *miliaria crystallina*, histologic examination reveals occlusion of the orifices of sweat ducts by keratin plugs and distention of the sweat ducts within the epidermis and, occasionally also in the dermis. Intracorneal vesicles are present. There is no inflammatory infiltrate. Evidence in favor of the assumption that the intracorneal vesicles contain sweat are the observations in experimental *miliaria crystallina* by Shelley and Horvath that the vesicles on serial sections proved to be in direct communication with sweat ducts and failed to form when sweating was inhibited by the local injection of atropine.

In *miliaria rubra*, an inflammatory infiltrate is present around the sweat ducts in the epidermis and the upper dermis. Either an intraepidermal or a subepidermal vesicle is seen. It appears that in severe cases of *miliaria rubra*, which are associated with thermogenic anhidrosis, the sweat duct ruptures either within the epidermis leading to an intraepidermal vesicle or at the epidermal dermal junction leading to a subepidermal vesicle (O'Brien, Sulzberger, Zimmerman and Emerson). On the other hand, in mild cases the sweat duct does not break, as a rule, but sweat merely escapes from the sweat duct into the epidermis leading to the formation of an intraepidermal "spongiotic" vesicle (Sulzberger and Zimmerman).

### POMPHOLYX (DYSHIDROTIC ERUPTION)

This is a recurrent eruption of numerous deep seated vesicles occurring singly and in groups on the palms and the soles. Inflammation

Tranck, A. Le cytodagnostic immédiat en dermatologie, Ann dermat. et syph. 8 205, 1948

### Dermatitis Herpetiformis

Burkhart, R. J., and Montgomery, H. Dermatologic significance of tissue eosinophilia Arch Dermat & Syph 68 10, 1954

Cava ..... dou

Lapi ..... vul  
gaite et de la dermatite polymorphe de Duhring Brocq Arch belges dermat et syph 1 216 1939

### Erythema Multiforme

Costello, M. J. Erythema multiforme exudativum J Invest Dermat 8 127, 1947  
Winer, L. H., and Lipschultz, C. E. Comparative study of histology and cytology in vesiculating eruptions Arch Dermat & Syph 65 270, 1952

### Burn

Moritz, A. R. Studies of thermal injury. III The pathology and pathogenesis of cutaneous burns Am J Path 23 915 1947



# 8

## Noninfectious Inflammatory Diseases

### URTICARIA

Urticaria is characterized by the presence of transient edematous macules or wheals, and is accompanied by considerable itching.

**Histopathology.** An urticarial wheal shows edema, particularly of the upper dermis. The collagen bundles as well as the individual fibers are separated by edema. The collagenous substance appears swollen and stains poorly. In early wheals, i.e., those only a few minutes old, one finds either no inflammatory reaction or merely a slight perivascular infiltrate composed of lymphocytes. Wheals an



FIG. 39 Prurigo nodularis. There are hyperplastic nodules of the epidermis approach

four old or older show a mild or even moderately severe lymphocytic infiltrate around the capillaries (Torok and Lehner)

## PRURIGO NODULARIS

usually chiefly

**Histopathology** One observes pronounced hyperkeratosis and acanthosis. There may be deep invaginations of the horny layer.

## ERYTHEMA NODOSUM

The lesions consist of tender red or livid red nodes which are slightly raised above the level of the skin. They vary from 1 to 5 cm in diameter and usually are limited to the anterior surfaces of the legs; they may however occur elsewhere. They involute within a few weeks without breaking down.

**Histopathology** The histologic changes are located mainly in the upper portion of the subcutaneous tissue. The dermis merely shows a moderate amount of perivascular infiltrate composed predominantly of lymphocytes.

In early lesions one observes in the upper portion of the subcutaneous tissue a scattered infiltrate consisting mainly of neutrophils and lymphocytes. Some histiocytes and occasionally eosinophils are present but plasma cells are absent. The inflammatory infiltrate extends as the disease progresses both upward toward the fatty tissue

irregular outline. No abscess formation or necrosis occur.

The blood vessels especially the veins may show severe involvement so that many authors believe that a vasculitis represents the primary and predominant lesion (Rotnes, Grzybowski). Other authors however state that the blood vessels are not necessarily affected severely and may show only mild involvement (Pautrier and Woringer, Lofgren and Wahlgren). In cases with vascular changes one observes especially in the larger veins invasion of the vascular

walls by the inflammatory infiltrate and marked endothelial proliferation (Fig 40). However, complete occlusion and thrombosis are rare.

Epithelioid and giant cells are absent in early lesions. However, occasionally one finds small nodules composed of histiocytes lying either in radial arrangement or in palisade like arrangement around




FIG. 40 Erythema nodosum. A nonspecific inflammatory infiltrate extends in small scattered patches through the subcutaneous fat. A large subcutaneous vein shows endothelial proliferation and invasion of its wall by inflammatory cells ( $\times 50$ ).

a small central fissure (Fig 41) (Miescher). Not infrequently these nodules are permeated by neutrophils. Since they are found in no other disease, they are of considerable diagnostic value (Nubé).

Older lesions show fewer neutrophils and more lymphocytes than young lesions. Often giant cells are present at this stage. The giant cells, usually of the foreign body type, may be found outside of or within foci of epithelioid cells. In the latter case, the arrangement simulates that found in tuberculosis (Rotnes). However, caseation is always absent.

**Differential Diagnosis.** For differentiation from erythema induratum, see page 186. In cases of erythema nodosum showing severe vascular involvement, periarteritis nodosa must be excluded. In the

latter disease however the arteries are predominantly involved one observes necrosis of vascular walls and the infiltrate usually contains a large percentage of eosinophils. Nodular vasculitis greatly resembles the late stage of erythema nodosum. It differs from erythema nodosum by showing a larger degree of vascular involvement including in




Fig. 41 Erythema nodosum. An older lesion shows two small nodules composed of histiocytes and to the right several giant cells of the foreign body type ( $\times 200$ )

volvement of vessels of large caliber. However it is possible that it represents a variant of erythema nodosum.

### NODULAR VASCULITIS

Clinically this disease is characterized by slightly painful nodules occurring chiefly on the legs. They generally persist for a

few weeks with iterative changes in both veins and arteries (Fig. 12). In addition one observes perivascular

walls by the inflammatory infiltrate and marked endothelial proliferation (Fig. 40). However, complete occlusion and thrombosis are rare.

Epithelioid and giant cells are absent in early lesions. However, occasionally one finds small nodules composed of histiocytes lying either in radial arrangement, or in palisade like arrangement around




FIG 40 Erythema nodosum. A nonspecific inflammatory infiltrate extends in small, scattered patches through the subcutaneous fat. A large subcutaneous vein shows endothelial proliferation and invasion of its wall by inflammatory cells ( $\times 50$ )

a small central fissure (Fig. 41) (Miescher). Not infrequently, these nodules are permeated by neutrophils. Since they are found in no other disease, they are of considerable diagnostic value (Nubé).

Older lesions show fewer neutrophils and more lymphocytes than young lesions. Often, giant cells are present at this stage. The giant cells, usually of the foreign-body type, may be found outside of, or within foci of, epithelioid cells. In the latter case, the arrangement simulates that found in tuberculosis (Rotnes). However, caseation is always absent.

**Differential Diagnosis.** For differentiation from erythema induratum, see page 186. In cases of erythema nodosum showing severe vascular involvement, periarteritis nodosa must be excluded. In the

is absent

The stratum malpighii is thinned above the papillae, often to only two or three layers of cells. The rete ridges show considerable elongation. They often are slender in their upper portion and thickened in



FIG 43 Psoriasis. Low magnification. There is parakeratosis. The rete ridges are elongated and thickened in their lower portion. There are edema and clubbing of the papillae. (X50)

their lower portion. They may show branching at their bases, causing

may be found scattered through the stratum malpighii.

Corresponding to the elongation and the branching of the rete ridges, the papillae are elongated and tortuous. The upper part of the papillae is edematous and club shaped. The capillaries in the papillae are dilated and tortuous and show slight thickening of their walls. These changes in the capillaries can be demonstrated best by the use of the Horchikiss McManus stain (Stoughton and Wells). A mild to moderately severe inflammatory infiltrate is present in the upper dermis, particularly in the papillae. It consists of lymphocytes

It is probable that nodular vasculitis is a variant of erythema nodosum. For their differential diagnosis see under erythema nodosum, see page 97.



FIG. 42. Nodular vasculitis. The wall of a large subcutaneous vein is greatly thickened and infiltrated with inflammatory cells. The lumen is obliterated ( $\times 50$ )

## PSORIASIS

*Psoriasis* is a chronic disorder characterized by dull red or brownish papules and plaques. The lesions are sharply demarcated, dry and usually covered with layers of fine silvery scales. As the scales are removed by gentle curettage, one frequently sees characteristic fine bleeding points. In severe cases, the disease may affect the entire skin and present the clinical picture of generalized exfoliative dermatitis (exfoliative psoriasis).

**Histopathology.** Histologically, psoriasis is characterized by (1) parakeratosis, (2) thinning of the suprapapillary portions of the stratum malpighii, (3) elongation of the rete ridges, (4) edema and clubbing of the papillae and (5) Munro micro-abscesses (Fig. 43).

Corresponding to the layered silvery scaling observed clinically, the horny layer is considerably thickened and consists predominantly of parakeratotic cells, arranged in lamellae with air spaces in between.


edema of the rete cells overlying the papillae and (4) capillary dilatation in the tips of the papillae

**EXFOLIATIVE PSORIASIS** The histologic picture of exfoliative psoriasis may still show enough of the characteristics of psoriasis to allow this diagnosis. Frequently however the appearance is indistinguishable from that of exfoliative dermatitis due to other causes

**Differential Diagnosis** The histologic picture of psoriasis may resemble that of lichen simplex chronicus. For their differentiation, see page 73. Differentiation from seborrheic dermatitis is not always possible. Seborrheic dermatitis may show all the features of psoriasis, though less pronounced than in psoriasis. In addition however, one finds a fair degree of spongiosis which in psoriasis is either very slight or absent

### PUSTULOSIS PALMARIS ET PLANTARIS (PUSTULAR PSORIASIS PUSTULAR BACTERID)

Two diseases have been described in the literature under the name of pustular psoriasis: generalized pustular psoriasis and pustular



PUSTULOSIS PALMARIS ET PLANTARIS (PUSTULAR BACTERID) A large intraepidermal unilocular pustule containing many neutrophils is present ( $\times 100$ )

psoriasis of the palms and the soles. Neither of them however, is a form of psoriasis. Actually, pustular lesions do not occur in psoriasis. So-called generalized pustular psoriasis is identical with the gen-



and histiocytes, except in early lesions which, in addition, show polymorphonuclear leukocytes in the infiltrate. Plasma cells are found only rarely and eosinophils hardly ever.

The Munro micro abscesses (Fig. 44) are located either in the stratum corneum or directly beneath it. They represent small ac-



FIG. 44 Psoriasis. High magnification of Figure 43. A Munro micro abscess (M) is located within the parakeratotic horny layer. The suprapapillary portions of the stratum malpighii are thinned and show intracellular edema. Intercellular edema, however, is absent. The capillaries in the tips of the papillae are dilated ( $\times 200$ ).

cumulations of neutrophils which have migrated there through the epidermis. Munro micro abscesses, as a rule, are found easily in early lesions. In older lesions, they are few in number or even absent. Thus, their absence does not rule out psoriasis. Neither does their presence establish a diagnosis of psoriasis, since Munro micro abscesses may occur also in seborrheic dermatitis, actiodermatitis continua (Hallopeau), keratosis blennorrhagica and parapsoriasis guttata.

The bleeding points which may be produced by gentle scraping of the skin correspond to the apices of papillae. They are due to the following histologic changes: (1) parakeratosis, (2) thinning of the stratum malpighii above the tips of the papillae, (3) intracellular

walls thus form a sponge like network in the interstices of which neutrophils continue to accumulate (Fig 46) As the pustule ages the cellular walls gradually break in the center of the pustule so that




FIG 46 *Acrodermatitis continua* of Hallopeau There is acanthosis with elongation of the rete ridges The upper stratum malpighii contains a spongiform pustule the cellular walls of the edematous squamous cells form a sponge like network in the interstices of which neutrophils have accumulated (X300)

a large cavity forms At the periphery of the pustule however, the network persists for a much longer time

**Differential Diagnosis.** The spongiform pustule of Kogoj represents a striking histologic lesion and is characteristic of *acrodermatitis continua* However it is not diagnostic of that disease since it occurs also in three other diseases *impetigo herpetiformis* *keratosis blennorrhagica* and *Reiter's disease* Differentiation of these four

eralized form of acrodermatitis continua (Lapierre, Bruck) and will be described there (see below)

Pustular psoriasis of the palms and the soles, a term introduced by Barber, is a misnomer. This disease is now referred to frequently as either pustular bacterid (Andrews) or acrodermatitis pustulosa (Sachs, MacKee and Rothstein). The term pustulosis palmaris et plantaris is preferable, however.

Pustulosis palmaris et plantaris is a chronic, indolent disorder limited to the palms and the soles and characterized by the appearance of crops of deep seated pustules within areas of erythema and scaling.

**Histopathology.** The histologic picture does not resemble that of psoriasis. A large, intra epidermal, unilocular pustule is present (Fig 45). It contains many neutrophils and disintegrated cellular elements. The epidermis surrounding the pustule shows slight acanthosis with little or no spongiosis. In the dermis underlying the pustule, a moderately severe inflammatory infiltrate is present, composed mainly of lymphocytes and histiocytes but containing also a few neutrophils. Histologic features of psoriasis, such as parakeratosis, elongation of the rete ridges, thinning of the suprapapillary portions of the stratum malpighii and dilatation of the papillary capillaries are always absent (Sachs and Scannone, Sachs, MacKee and Rothstein).

## ACRODERMATITIS CONTINUA

(Hallopeau)

Acrodermatitis continua of Hallopeau, a chronic disease having pustules as a primary lesion, usually is limited to the hands and the feet. The distal portions of the fingers and the toes are predominantly involved. Occasionally, however, the disease is generalized. The affected areas are dark red, dry, shiny and scaling and are studded with shallow pustules. Lesions located on the distal portions of the fingers and the toes may cause atrophy of the skin and loss of nails.

**Histopathology.** The epidermis shows parakeratosis and moderate acanthosis with elongation of the rete ridges. An inflammatory infiltrate containing many neutrophils but few or no eosinophils is present in the dermis. Many neutrophils are seen invading the epidermis.

However, the characteristic lesion consists of the so called spongiiform pustule of Kogoj. This type of pustule forms in the uppermost stratum malpighii through the migration of neutrophils into edematous squamous cells. This invasion causes disintegration of the cytoplasm and of the nucleus but not of the cellular walls. The cellular

walls thus form a sponge like network in the interstices of which neutrophils continue to accumulate (Fig 46) As the pustule ages, the cellular walls gradually break in the center of the pustule so that




FIG 46 Acrodermatitis continua of Hallopeau There is acanthosis with elongation of the rete ridges The upper stratum malpighi contains a spongiform pustule the cellular walls of the edematous squamous cells form a sponge like network in the interstices of which neutrophils have accumulated ( $\times 500$ )

a large cavity forms At the periphery of the pustule however, the network persists for a much longer time

**Differential Diagnosis** The spongiform pustule of Kogoj represents a striking histologic lesion and is characteristic of acrodermatitis continua However, it is not diagnostic of that disease since it occurs also in three other diseases impetigo herpetiformis, keratosis blennorrhagica and Reiter's disease Differentiation of these four

diseases may be impossible. However, in impetigo herpetiformis, the infiltrate contains, as a rule, a large number of eosinophils (see below). Older lesions of keratosis blennorrhagica and Reiter's disease often can be differentiated from acrodermatitis continua by the presence of a greatly thickened horny layer (see page 172).

### IMPETIGO HERPETIFORMIS

*This rare, usually fatal, disease is seen mainly in pregnant women and occasionally in hypoparathyroidism. It is characterized by the presence of pustules set in groups or in circinate arrangement on erythematous patches.*

**Histopathology.** The histologic picture, like that of acrodermatitis continua, is characterized by the presence of the spongiform pustule (Kogoj). (See "Acrodermatitis Continua" for a description of the spongiform pustule.) The dermal infiltrate as well as the pustule contain a conspicuous number of eosinophils (Rost).

**Differential Diagnosis.** The presence of many eosinophils aids in the differentiation of impetigo herpetiformis from the other three diseases with spongiform pustules: acrodermatitis continua, keratosis blennorrhagica and Reiter's disease. The latter two diseases, in addition, often show conspicuous hyperkeratosis in their older lesions.

### PARAPSORIASIS

Parapsoriasis comprises a group of rare dermatoses. Three of the four forms of parapsoriasis are characterized by an asymptomatic maculopapular eruption of slow evolution and marked chronicity. These forms are parapsoriasis guttata, parapsoriasis lichenoides (parakeratosis variegata) and parapsoriasis en plaques. In parapsoriasis guttata, one observes especially on the trunk numerous papules which may or may not be covered with fine scales. In parapsoriasis lichenoides, papules generally are distributed in a netlike arrangement. In parapsoriasis en plaques, well defined patches and plaques of various size and shape occur. The fourth form of parapsoriasis, parapsoriasis varioliformis of Habermann, also called pityriasis lichenoides et varioliformis acuta, differs from the previously mentioned forms by running an acute course terminating in from a few weeks to several months. The eruption consists of vesicles, papules and papulonecrotic lesions terminating in varioliform scars.

**Histopathology.** None of the four forms of parapsoriasis shows a diagnostic histologic picture. Parapsoriasis guttata, parapsoriasis lichenoides and parapsoriasis en plaques show the histologic picture of chronic dermatitis, while parapsoriasis varioliformis shows an acute inflammatory process with foci of necrosis.

the infiltrate in the manner of a claw clutching a ball

The cause of the disease is not known Ellis and Hill regard lichen nitidus as closely related to lichen planus because they found that the two diseases may occur together and histologic sections obtained from patients with lichen planus may show lichen nitidus like lesions

### LICHEN STRIATUS

Lichen striatus is an uncommon eruption which as a rule occurs in children It manifests itself usually on the extremities as a long band composed of small lichenoid papules The eruption appears suddenly and involutes within a few weeks or months Itching is absent

**Histopathology** The histologic changes are similar to those of neurodermatitis circumscripta In some cases the chronic inflammatory infiltrate is more markedly perivascular than in neurodermatitis circumscripta and surrounds the vessels of the upper dermis as densely packed maniles (Senear and Caro Pinkus)

### PITYRIASIS RUBRA PILARIS

The primary lesions are reddish follicular papules They gradually coalesce Ultimately most of the late stage however especially on the dorsa of the fingers

**Histopathology** The essential pathologic process is follicular hyperkeratosis In addition there is diffuse hyperkeratosis with spotted parakeratosis The epidermis shows irregular acanthosis usually of mild degree There often is liquefaction degeneration of the basal cells (Brunsting and Sheard) In the upper dermis a mild chronic inflammatory infiltrate is observed around the blood vessels

**Differential Diagnosis** For a differential diagnosis from phrynoderma (vitamin A deficiency) see page 286

### GRANULOMA FACIALE (EOSINOPHILIC GRANULOMA OF THE FACE)

This disorder only recently established as a disease entity consists of soft purplish slowly growing and asymptomatic patches limited to the face Except for prominence of the follicular openings the surface of the skin appears normal

**Histopathology.** A dense, granulomatous infiltrate is located mainly in the upper half of the dermis (Fig 52). Quite characteristically, the infiltrate does not invade the epidermis or the pilosebaceous appendages but is separated from them by a narrow zone of normal collagen. The pilosebaceous appendages are well preserved. Two stages can be differentiated in the development of the



FIG 52 Granuloma faciale. A dense infiltrate containing many eosinophils is present in the dermis. The infiltrate does not invade the epidermis or the sebaceous gland shown on the right but is separated from them by a zone of normal collagen ( $\times 100$ ).

infiltrate, an early "leukocytic" and a late "fibrotic" stage. The two stages may occur together in the same lesion.

In the early "leukocytic" stage, eosinophils predominate, but neutrophils and histiocytes are found in large number and, in addition, a few lymphocytes, plasma cells and mast cells. The infiltrate thus has a polymorphous appearance. The capillaries are dilated and surrounded by degenerated collagen (Fig 53). In some cases considerable amounts of hemosiderin are present in the upper dermis within and outside of histiocytes (Lever, Lane, Downing and Spangler). The presence of foam cells has been described in one case (Lever).

In the late "fibrotic" stage, the infiltrate is broken up into variously sized patches by strands of collagen. The number of eosinophils and neutrophils is greatly reduced, and plasma cells, lymphocytes and fibroblasts predominate. The capillaries often have fibrotic walls.




FIGURE 1. Infiltrate is densest around a dilated capillary. Two types of cells, eosinophils and histiocytes, predominate. (X400)

**Differential Diagnosis.** The arrangement and the composition of the infiltrate in granuloma faciale result in a diagnostic histologic picture with no resemblance to that of eosinophilic granuloma with which the disease was at first confused (Lever). Although eosinophilic granuloma, which represents an abortive form of Hand-Schüller-Christian disease (see page 265), also contains many eosinophils and histiocytes, its histologic appearance differs from that of granuloma faciale by the following: (1) the infiltrate is much larger and (2) the arrangement is in a well-defined patch.



the epidermis (Lever and Leeper) For a differentiation from *erythema elevatum diutinum*, see below

### ERYTHEMA ELEVATUM DIUTINUM

This rare disorder shows persistent, red to purple nodules with some tendency to grouping about the joints, especially on the dorsa of the hands, the wrists and the elbows. The nodules at first are soft but later, due to fibrous transformation, become hard.

**Histopathology.** Erythema elevatum diutinum has a fairly diagnostic histologic picture which was first described by Weidman and Besncon, and then by Ketron and by Weiss as well as by others. The disease described in 1891 by Crocker and Williams as erythema elevatum diutinum of the Bury type is now widely regarded as a variant or late stage of granuloma annulare (Haber and Russell).

Erythema elevatum diutinum, in its early stage, shows a dense, predominantly perivascular infiltrate composed largely of neutrophils intermingled with some lymphocytes and histiocytes. In most cases, a peculiar hyaline degeneration of the reticulum fibers around the capillaries has been observed. This degeneration has been referred to by Weidman and Besncon and by Ketron as formation of "toxic hyalin." This toxic hyalin has a smooth, glassy appearance and stains intensely pink with eosin.

In the late, fibrous stage, the cellular infiltrate is much less pronounced and extensive fibrosis is present. The capillaries may still show their mantle of toxic hyalin or may merely show fibrous thickening (Ketron).

**Differential Diagnosis.** The histologic appearance of erythema elevatum diutinum is distinguished from that of granuloma annulare by the absence of a zone of normal collagen beneath the epidermis and the presence of toxic hyalin about the capillaries.


### GRANULOMA ANNULARE

The lesions which are found most commonly on the hands and the feet, consist of small, firm, pale red nodules which tend to be symmetrical in distribution. The disease is chronic.

Granuloma annulare is characterized by focal degeneration of the collagen in the dermis, deposits of mucin between the degenerated collagenous bundles and reactive inflammation and fibrosis. The degeneration of collagen within the foci may be complete or incomplete. Some cases show foci of both complete and incomplete degeneration and others show only one



fo  
hu



d  
la  
tra Ne  
eral to  
Rammz

large focus of complete collagen degeneration. In the majority of cases, however, one finds multiple foci of incomplete degeneration without any foci of complete degeneration.

In foci of complete degeneration one finds a sharply demarcated area of coagulation necrosis surrounded by an infiltrate of histiocytes, fibroblasts and lymphocytes in a radial arrangement (Fig. 54).

In foci of incomplete degeneration one finds ill defined areas in which collagen fibers are found in various stages of degeneration ranging from mild homogenization and fragmentation to granular degeneration and coagulation necrosis. A considerable amount of the collagen in such areas may appear normal. Lymphocytes, histiocytes and fibroblasts infiltrate between the partially degenerated collagen fibers and new collagen is being laid down. Thus the affected areas of the dermis present a completely disorderly arrangement of the collagen bundles (Fig. 55).

In both types of collagen degeneration, though more frequently in areas of incomplete degeneration, fine threads and granules of mucin are deposited between the degenerated collagen bundles. The mucin stains light blue with routine stains and red when Best's mucicarmine stain is used (See Plate 1).

In areas of collagen degeneration the elastic tissue shows fragmentation and may be largely destroyed. In some areas there may even be greater destruction of elastic than of connective tissue. A moderate or even considerable amount of lymphocytic infiltration is found around the blood vessels outside the areas of degeneration. The walls of the blood vessels show no pathologic changes, however, except occasional mild endothelial proliferation. In some cases a few giant cells of the foreign body type are present. They usually are situated near the periphery of the infiltrate and are not associated with any zone of necrosis (Prunty and Montgomery).

**Differential Diagnosis.** The type of granuloma annulare showing multiple foci of incomplete collagen degeneration may greatly resemble necrobiosis lipoidica which shows this same pattern of collagen degeneration (see page 269). However, necrobiosis lipoidica differs from granuloma annulare by the presence of vascular changes of lipid material and of many foreign body giant cells and by the absence of mucin.

The type of granuloma annulare showing one or several large foci of complete collagen degeneration may be almost indistinguishable from the subcutaneous nodes of rheumatoid arthritis (see below). However, in the nodes of rheumatoid arthritis the areas of degeneration are in the dermis and are associated with increased cellular

eaction, mainly histiocytes and fibroblasts, while in granuloma annulare there are, in addition, many lymphocytes (Bowers)

### SUBCUTANEOUS NODULES OF RHEUMATIC FEVER AND RHEUMATOID ARTHRITIS

In both rheumatic fever and rheumatoid arthritis, small firm nodules may form in the subcutaneous tissue. The commonest sites



FIG. 56 Subcutaneous nodule of rheumatoid arthritis. There is a large central zone of necrosis, surrounded by histiocytes in palisade arrangement ( $\times 100$ )

are about the elbows, the knees and the ankles. Their size varies from a few millimeters to 2 cm.

**Histopathology** The subcutaneous nodules found in these two diseases are *similar in many of their pathologic features*. In both diseases they are *composed of three reasonably well-defined zones* (1) a *central zone, or zone of necrosis*, (2) an *intermediate zone*, comprising proliferating histiocytic cells and (3) a *peripheral zone*, consisting of chronic inflammatory cells (Bennett, Zeller and Bauer). However, *exudative changes prevail in the nodules of rheumatic fever*, *proliferative and necrotic changes in those of rheumatoid arthritis*.

large focus of complete collagen degeneration. In the majority of cases, however, one finds multiple foci of incomplete degeneration without any foci of complete degeneration.

In foci of complete degeneration, one finds a sharply demarcated area of coagulation necrosis surrounded by an infiltrate of histiocytes, fibroblasts and lymphocytes in a radial arrangement (Fig. 54).

In foci of incomplete degeneration, one finds ill defined areas in which collagen fibers are found in various stages of degeneration ranging from mild homogenization and fragmentation to granular degeneration and coagulation necrosis. A considerable amount of the collagen in such areas may appear normal. Lymphocytes, histiocytes and fibroblasts infiltrate between the partially degenerated collagen fibers and new collagen is being laid down. Thus, the affected areas of the dermis present a completely disorderly arrangement of the collagen bundles (Fig. 55).

In both types of collagen degeneration, though more frequently in areas of incomplete degeneration, fine threads and granules of mucin are deposited between the degenerated collagen bundles. The mucin stains light blue with routine stains and red when Best's mucicarmine stain is used (See Plate 1).

In areas of collagen degeneration, the elastic tissue shows fragmentation and may be largely destroyed. In some areas, there may even be greater destruction of elastic than of connective tissue. A moderate or even considerable amount of lymphocytic infiltration is found around the blood vessels outside the areas of degeneration. The walls of the blood vessels show no pathologic changes, however, except occasional mild endothelial proliferation. In some cases, a few giant cells of the foreign body type are present. They usually are situated near the periphery of the infiltrate and are not associated with any zone of necrosis (Prunty and Montgomery).

**Differential Diagnosis.** The type of granuloma annulare showing multiple foci of incomplete collagen degeneration may greatly resemble necrobiosis lipoidica which shows this same pattern of collagen degeneration (see page 269). However, necrobiosis lipoidica differs from granuloma annulare by the presence of vascular changes, of lipid material and of many foreign body giant cells and by the absence of mucin.

The type of granuloma annulare showing one or several large foci of complete collagen degeneration may be almost indistinguishable from the subcutaneous nodes of rheumatoid arthritis (see below). However, in the nodes of rheumatoid arthritis the areas of degeneration are usually larger, they are more deeply located in the dermis or even in the subcutis, and they are encircled by a less varied cellular

After many years' duration, an atrophic stage may be reached in which the findings are no longer diagnostic, since one sees merely an atrophic epidermis and an atrophic, fibrotic dermis without inflammatory infiltrate.



FIG 37. *Acrodermatitis chronica atrophicans*. There is atrophy of the stratum malpighii. A bandlike infiltrate is separated from the epidermis by a narrow zone of normal collagen. The dermis shows interstitial edema and atrophy of the collagen bundles. Because of this atrophy, the thickness of the dermis is markedly decreased and the sweat glands lie unusually close to the epidermis ( $\times 100$ ).

The bandlike areas of fibrosis resembling scleroderma and the

able from that of scleroderma

### RADIODERMATITIS

An early (acute) and late (chronic) stage of radiodermatitis are recognized. The early stage occurs within a few days after adminis-

In rheumatic fever, the central zone shows "fibrinoid degeneration" (For a discussion of fibrinoid degeneration, see "Acute Systemic Lupus Erythematosus," page 297) The intermediate zone is composed of edematous collagen infiltrated with inflammatory cells, including many large mononuclear cells (histiocytes) which resemble the predominant cells of myocardial Aschoff nodules

In rheumatoid arthritis, the central zone shows necrosis of all pre-existing collagen, and the intermediate zone distinct palisading of histiocytes and fibroblasts around the central zone of necrosis (Fig 56)

**Differential Diagnosis.** For differentiation of the nodules of rheumatoid arthritis from granuloma annulare, see 'Granuloma Annulare,' page 116

### ACRODERMATITIS CHRONICA ATROPHICANS

This condition affects the extremities. The skin presents a bluish red or a brownish, atrophic, wrinkled appearance. Fine scaling is usually present. Because of decrease in the amount of subcutaneous fat, the subcutaneous veins are clearly visible. In some cases, bandlike areas of fibrosis, resembling scleroderma, and fibrous nodules are observed, especially along the ulna as "ulnar bands" and along the tibia

**Histopathology.** The histologic picture is characteristic. One observes mild to moderate hyperkeratosis and atrophy of the epidermis, with absence of the rete ridges. Just beneath the epidermis, there is a narrow zone of connective tissue separating a dense band of inflammatory infiltrate from the epidermis. In addition to this bandlike infiltrate, one finds scattered areas of inflammatory infiltration throughout the dermis, particularly around the blood vessels. The infiltrate is composed predominantly of lymphocytes, but also contains histiocytes. Chromatophores laden with melanin or hemosiderin may be present. The entire dermis shows interstitial edema and atrophy of the collagenous bundles. The atrophy results in a gradual decrease of the dermis to a half or a quarter of its normal thickness (Fig 57)

The sebaceous glands and the hairs undergo atrophy early in the disease and usually are entirely absent. However, the sweat glands are, as a rule, preserved. Because of the thinness of the dermis they lie unusually close to the epidermis. The blood vessels are dilated and may show endothelial proliferation

The subcutaneous tissue shows decided atrophy. The fat cells vary in size and are irregular in shape. Foci of inflammatory infiltration and a varying degree of fibrosis are present

changes. An inflammatory infiltrate is seen throughout the dermis, especially about the appendages. The blood vessels are dilated and reveal edema of their walls and endothelial proliferation. The collagen bundles show edema and homogenization. In severe cases, the



FIG. 59 Late radiodermatitis. The vessels (V) show

epidermis and the upper dermis undergo necrosis. The area of necrosis is invaded and surrounded by polymorphonuclear leukocytes.

**Histopathology of Late (Chronic) Radiodermatitis.** The epidermis is irregular, showing atrophy in some areas and acanthosis with hyperkeratosis in others. The rete cells frequently show atypicality, such as disorderly arrangement, individual cell dyskeratosis and an increased number of mitotic figures. Thus the changes may resemble those of Bowen's disease (intra-epidermal squamous-cell carcinoma). In addition, there often is irregular downward growth of the epidermis. The



tration of a massive dose of roentgen rays or radium. At first, there is erythema, which is followed by desquamation and pigmentation. If the dose administered is sufficiently large, ulceration will ensue. The late (chronic) stage of radiodermatitis occurs from a few months



FIG. 58 Late radiodermatitis. The epidermis shows acanthosis and downward growth around a telangiectatic blood vessel. Many irregularly dilated lymphatics are located directly beneath the epidermis. The collagen shows degeneration ( $\times 100$ ).

to several years after the administration of large amounts of roentgen rays or radium. The skin shows atrophy with interspersed areas of hyperkeratosis, irregular hyperpigmentation, telangiectases and loss of hair. Ulceration may be present. The hyperkeratoses may develop into squamous cell carcinoma.

**Histopathology of Early (Acute) Radiodermatitis.** The cells of the epidermis, particularly those of the basal layer, are hydropic and their nuclei show pyknosis. The epithelial cells of the hair follicles, the sebaceous glands and the sweat glands may show similar degenerative

**Histopathology** The histologic appearance usually is that of a nonspecific chronic dermatitis (see page 70). However the erythematous lesions frequently present areas of liquefaction degeneration of the basal layer and edema together with a patchy predominantly lymphocytic infiltrate in the upper dermis. Differentiation from lupus erythematosus may then be impossible. However in some cases of solar dermatitis a moderate number of eosinophils is present in the infiltrate. Since eosinophils are generally not observed in lupus erythematosus their presence aids greatly in the differentiation of the two diseases (Lamb).

### HIDROCYSTOMA

This condition is characterized by the presence on the face of numerous clear tense deep-seated vesicles varying in size from a few millimeters to 1 centimeter.

**Histopathology** On histologic examination one observes in the dermis large cystic cavities lined by epithelial cells which usually are arranged in two layers. Small papillary projections into the lumina of the cysts have been described (Kenedy and Lehner).

Most authors regard the cysts as caused by retention of sweat in the dermal portion of eccrine sweat ducts. However Kenedy and Lehner believe that the cysts are not merely the result of retention of sweat and passive distention of the excretory ducts but are due to a proliferation of the excretory duct epithelium which leads to occlusion of the ducts.

### ACNE VULGARIS

Acne vulgaris a disease occurring predominantly in puberty affects the face the upper chest and the upper back. The primary lesion is the comedo. As a foreign body reaction to the comedo and due to secondary infection follicular papules and pustules and subcutaneous abscesses may occur.

**Histopathology** Histologically acne vulgaris represents a per folliculitis occurring around a comedo. Comedones are composed of sebum and keratotic as well as parakeratotic cells and are located in the pilosebaceous follicles.

In the early acne lesion one finds an infiltrate of lymphocytes and plasma cells around a comedo-filled pilosebaceous follicle together with atrophy and fragmentation of sebaceous glands. The atrophy and the fragmentation of sebaceous glands probably are due to the stasis and the pressure caused by the comedo within the follicle. The fragmentation of sebaceous glands in turn causes an inflammatory reaction (Lynch). Occasionally foreign body giant cells are found in

rete ridges may grow around telangiectatic vessels, which may thus become completely enclosed in the epidermis (Fig 58).

In the dermis, the collagenous fibers are swollen and sclerotic. In some areas, they are broken apart and stain basophilic. Formation of new, young connective tissue is found throughout the dermis, but especially around the blood vessels. Directly beneath the epidermis, one often sees numerous irregularly dilated lymphatics as well as lymphedema.

However, the fundamental pathologic change is that of fibrotic thickening of the walls of the vessels in the deeper portions of the dermis, leading to occlusive changes in the lumina (Fig 59). Some of the vessels show thrombosis and recanalization. The vessels nearest to the epidermis may show telangiectasia.

Hair follicles and sebaceous glands are absent, but the sweat glands usually are preserved, except in areas of third degree injury. The elastic tissue is, as a rule, less damaged than the collagenous tissue—at least in milder degrees of radiodermatitis. In severe reactions, it is destroyed. With Foot's reticulum stain, the reticulum fibers are seen to be increased throughout the dermis wherever there is formation of young connective tissue, but especially around blood vessels.

In severe cases of chronic radiodermatitis, ulceration occurs. The deep lying, large blood vessels beneath the ulcers, as a rule, show complete obliteration.

The above described epidermal changes of dyskeratosis and intra epidermal carcinoma frequently lead on to invasive squamous cell carcinoma. A fairly large percentage of them are of high (grade IV) malignancy and of the spindle celled variety (see page 338). In rare instances, basal cell epitheliomas develop in areas of radiodermatitis (Anderson and Anderson) (see under "Basal Cell Epithelioma," page 370). The occurrence of sarcoma, on the other hand is not fully established. Such instances have been reported (Blom Ides, Gentile), but always with the reservation that the diagnosis of spindle celled squamous-cell carcinoma could not be ruled out conclusively because this differentiation may be nearly impossible (see page 410). A point in favor of their being carcinoma is the fact that they are, in contrast to fibrosarcoma, radiosensitive (Blom Ides).

### SOLAR DERMATITIS

Solar dermatitis may occur as erythematous, urticarial, papular or vesicular lesions in areas exposed to the sun. The erythematous lesions may resemble lupus erythematosus in their clinical appearance by showing a livid red color and slight induration.

but he concedes that differentiation of the two diseases by histologic means may be impossible. Additional but not obligatory findings in the papular type of rosacea are superficial perifolliculitis with intrafollicular abscess formation ( pustules ) and dilatation of capillaries. There is no hypertrophy of the sebaceous glands.

In the glandular hyperplastic form the sebaceous glands are increased in size and number. The orifices of the sebaceous glands are dilated and filled with keratin and sebum. There is an associated hypertrophy of the connective tissue. The blood vessels are dilated. A chronic inflammatory infiltrate is present around the vessels. Perifolliculitis with intrafollicular abscess formation may be observed.

### FOX FORDYCE DISEASE

This disease occurs only in women. It is characterized by an eruption

apocrine glands is the limitation of the eruption to the areas where apocrine glands occur and the onset of the disease at puberty when the apocrine glands begin to function. Shelley and Hurley have expressed the opinion that Fox Fordyce disease may be basically a miliaria apocrina—in other words an apocrine sweat retention miliaria.

**Histopathology** Histologic examination reveals acanthosis with hyperkeratosis. The rete ridges are elongated. A moderate degree of round cell infiltration is present around the blood vessels in the upper dermis. The appearance thus is that of a chronic dermatitis.

The apocrine glands appear normal (Roxburgh). The statement frequently found in the literature (Nilén) that in Fox Fordyce disease the sweat glands are dilated is due to the fact that many authors have misinterpreted the apocrine glands as dilated sweat glands.

### ALOPECIA AREATA

Alopecia areata is characterized by loss of hair in one or more

in occasional instances the entire scalp is involved (alopecia totalis) in which case the loss of hair is usually permanent.

**Histopathology** The hair follicles and the hair bulbs are greatly reduced in size. The hair bulbs instead of being deep in the subcutis are located quite high up in the dermis. The hair follicles contain loose keratin and no hair. In long standing cases there is

the infiltrate but there is no evidence of tuberculoid reaction. At a later stage histiocytes and fibroblasts appear and fibrosis occurs.

In lesions in which secondary bacterial infection is present one finds disintegration of the follicle and abscess formation. Foreign body giant cells are then frequently found around the remnants of the follicle.

### ACNE VARIOLIFORMIS (ACNE NECROTICA MILIARIS)

Acne varioliformis is characterized by the presence of small indolent papules and pustules along the frontal hairline. The lesions undergo central necrosis and tend to heal with pitted scars.

Acne necrotica miliaris of the scalp is a diminutive variant of acne varioliformis. Because of the superficial location of the necrosis no hair loss ensues.

**Histopathology** The histologic changes of acne necrotica are similar to those of acne vulgaris except that the perifolliculitis is more limited in extent and almost invariably results in central necrosis. Blood vessels adjacent to the necrotic area may be thrombosed. Healing takes place with fibrosis and scar formation.

Acne necrotica miliaris of the scalp shows the same histologic picture as that of acne varioliformis (Montgomery).

### ACNE ROSACEA

Acne rosacea occurs in patients with seborrhea and affects the central portion of the face. Two forms of rosacea exist: the papular form and the glandular hyperplastic form. The two forms may occur together. The papular form is characterized by erythema, papules, pustules and telangiectases. The glandular hyperplastic form causes enlargement of the nose, called rhinophyma.

**Histopathology** In the papular variety the infiltrate is either diffusely distributed throughout the dermis or arranged about the hair follicles and the sebaceous glands. As a rule it is banal in character and composed largely of lymphocytes together with some histiocytes and plasma cells. However, a significant number of patients show in addition foci of epithelioid cells and Langhans giant cells, occasionally lying in true tubercle formation, so that the histologic picture is indistinguishable from that of cutaneous tuberculosis. Miescher, who found tuberculoid foci in 80 per cent of 59 patients with acne rosacea, concluded that the rosacea-like tuberculoid of Lewandowsky and the papular type of acne rosacea represent the same disease (see page 183). On the other hand, Laymon, who observed tuberculoid structures in only 11 per cent of 138 patients with acne rosacea, does not think that the two diseases are identical.

Purpura occurs as the result of either noninflammatory or inflammatory changes in the walls of small blood vessels

### NONINFLAMMATORY PURPURA

The following types of purpura are noninflammatory stasis purpura due to increased intracapillary pressure senile purpura due to degeneration of the dermal collagen in exposed areas of the skin scurvy due to reduction of the intercellular material between endothelial cells thrombocytopenic purpura due to inadequate formation of blood platelets and febrile purpura with platelet thrombosis a rare systemic fatal disease

**Histopathology** In stasis purpura scurvy and thrombocytopenic purpura no visible vascular changes are present and the only abnormality consists of the presence of extravasated red blood cells and at a later stage of hemosiderin In senile purpura one observes in addition severe degenerative changes of the dermal collagen as seen in senile elastosis (see page 157) (Tattersall and Seville) In febrile purpura with platelet thrombosis one observes in the capillaries of the skin as well as of many internal organs degeneration and proliferation of endothelial cells and obstructing thrombi composed of platelets (Trosbaugh)

### INFLAMMATORY PURPURA (VASCULITIS)

The following types of purpura are caused by inflammatory changes in the walls of small blood vessels (vasculitis) bacterial purpura due to meningococci *Streptococcus viridans* or other microorganisms anaphylactoid or Schoenlein Henoch purpura which is of unknown genesis and often associated with joint pains gastrointestinal bleeding and occasionally with glomerulonephritis purpura due to drug allergy and purpura associated with periarteritis nodosa

**Histopathology** There are vascular changes and an inflammatory infiltrate in addition to the extravasation of red blood cells

For the histologic changes in the purpuric lesions of meningococcemia and subacute bacterial endocarditis see pages 170 and 171 respectively For purpura associated with periarteritis nodosa see page 311 Purpura due to drug allergy has the same histologic picture as that of anaphylactoid purpura

**ANAPHYLACTOID PURPURA** Early lesions show prominent vascular changes in the upper dermis consisting of swelling and degeneration of endothelial cells and occasionally of focal necrosis of the vascular wall (Gardner Levinson) A rather severe inflammatory infiltrate is

gross thinning of the dermis, which may be reduced to half its normal thickness (Dillaha and Rothman). In cases of recent onset, a mild to moderately severe inflammatory infiltrate composed of lymphocytes is seen in the deeper dermis about the vessels, the sebaceous glands and the hair follicles, while, in cases of more than 1 year's duration, the inflammatory reaction has subsided (Laymon). The sebaceous and the sweat glands appear normal throughout the course of the disease.

### ALOPECIA CICATRISATA (PSEUDOPELADÉ BROCC)

In alopecia cicatrisata, one finds scattered through the scalp irregularly shaped patches of alopecia which in the early stage may show perifollicular erythema but in the late stage show smooth atrophy of the skin without any signs of inflammation. The loss of hair is permanent.

**Histopathology.** In the early stage, one finds a predominantly perifollicular infiltrate composed almost entirely of lymphocytes and a few histiocytes. The infiltrate is present around the upper and the middle thirds of the follicles and spares the lower third. It penetrates into the walls of the follicles and into the sebaceous glands, but not into the lumen of the follicles. Follicular hyperkeratosis may be present (Miescher and Lenggenhager). Gradually, the infiltrate destroys the follicles and the sebaceous glands.

In a late lesion, the epidermis is atrophic and the dermis shows fibrosis. The follicles and the sebaceous glands are absent, but arrectores pilorum and sweat glands are, at least in part, preserved.

**Differential Diagnosis.** In discoid lupus erythematosus, the inflammatory infiltrate not only is located around hairs and sebaceous glands but also is distributed in a patchy fashion throughout the dermis. In addition, the epidermis shows liquefaction degeneration of the basal cell layer, hyperkeratosis and more marked keratotic plugging not limited to the follicles. In the late cicatricial stage, a differentiation of the two diseases may be impossible.

**Folliculitis decalvans**, which represents a folliculitis of the scalp, differs from alopecia cicatrisata by showing formation of intrafollicular abscesses (pustules). The perifollicular infiltrate often contains a fair number of plasma cells which are absent in alopecia cicatrisata (Miescher and Lenggenhager).

### PURPURA

Purpura represents a hemorrhage into the skin. Lesions less than 3 mm in diameter are called petechiae. Larger lesions are called ecchymoses.

non which occurs in sensitized animals at the site of a subcutaneous injection of foreign protein. The presence of leukocytoclasia in anaphylactoid purpura would therefore support the view that this disease represents an allergic reaction (Rutter and Brandsma)

# PURPURA PIGMENTOSA PROGRESSIVA (Majocchi-Schamberg)

Four diseases are included under this term: purpura telangiectodes (Majocchi), progressive pigmentary dermatosis (Scham-



FIG. 61. Purpura pigmentosa progressiva (Majocchi-Schamberg). A group of capillaries located in a papilla show swelling, proliferation and degeneration of their endothelial cells. A small focus of extravasation of erythrocytes can be seen to the right of the group of capillaries ( $\times 400$ ).

berg), pigmented purpuric lichenoid dermatitis (Gougerot and Blum) and angioma serpiginosum. They are so closely related to one another that often they cannot be differentiated on clinical or histologic



present predominantly, but not exclusively around the damaged vessels (Fig 60). It consists largely of neutrophils and varying amounts of eosinophils with only a few lymphocytes. A characteristic feature is the presence of many scattered nuclear fragments the result of the disintegration of neutrophils (leukocytoclasis). Extravasation of



FIG 60 Anaphylactoid purpura (inflammatory purpura). The capillaries show swelling and degeneration of endothelial cells. A rather severe inflammatory infiltrate is present especially around the capillaries. It is composed largely of neutrophils many of which show disintegration of their nuclei (leukocytoclasis). This being an early lesion no extravasation of erythrocytes has as yet occurred ( $\times 200$ ).

erythrocytes is usually present although it may be absent in the early stage of the disease.

In older lesions extravasation of red cells usually is a prominent feature. In addition hemosiderin may be present due to the decomposition of red cells. The endothelial lining of the vessels may show proliferation. The infiltrate contains fewer neutrophils and eosinophils and consists predominantly of lymphocytes. However nuclear fragments are still present as a rule.

It may be pointed out that fragmentation of the nuclei of neutrophils is prominent in the histologic picture of the Arthus phenomenon.

non which occurs in sensitized animals at the site of a subcutaneous injection of foreign protein. The presence of leukocytoclasia in anaphylactoid purpura would therefore support the view that this disease represents an allergic reaction (Ruiter and Brandsma).

### PURPURA PIGMENTOSA PROGRESSIVA (Majocchi-Schamberg)

Four diseases are included under this term: purpura annularis telangiectodes (Majocchi), progressive pigmentary dermatosis (Scham-



FIG. 61. Purpura pigmentosa progressiva (Majocchi-Schamberg). A group of capillaries located in a papilla show swelling, proliferation and degeneration of their endothelial cells. A small focus of extravasation of erythrocytes can be seen to the right of the group of capillaries ( $\times 400$ ).

berg), pigmented purpuric lichenoid dermatitis (Gougerot and Blum) and angioma serpiginosum. They are so closely related to one another that often they cannot be differentiated on clinical or histologic

grounds. Their separation into different entities, therefore, is unwarranted (Randall, Kierland and Montgomery). The term purpura pigmentosa progressiva is suggested for this disease.

The basic process is a chronic capillaritis of unknown cause occurring in the upper dermis and leading to capillary fragility. Clinically, the primary lesion consists of purpuric puncta appearing in groups and slowly extending so that various sized patches form. Gradually, pigmentation, due to the deposition of hemosiderin, supervenes and in cases of long standing, may dominate the clinical picture. Inflammatory signs (such as erythema, scaling and papules) may be present or absent. In most instances, the disease is limited to the lower extremities but may be more or less generalized. The disease, though chronic, is harmless.

**Histopathology.** In early lesions, the capillaries of the upper dermis show swelling, degeneration and proliferation of their endothelial cells. Often the number of capillaries appears to be increased. Small amounts of extravasated red cells are usually found in the vicinity of some capillaries. A cellular infiltrate, consisting largely of lymphocytes together with some histiocytes and occasionally a few neutrophils, is present in the upper dermis, especially in the vicinity of the capillaries. The inflammatory infiltrate may invade the lower epidermis and provoke mild liquefaction degeneration of the basal cell layer, mild spongiosis of the stratum malpighii and patchy parakeratosis.

In older lesions, the number of capillaries is usually increased. Some show proliferation of their endothelium and others dilatation of their lumen. Extravasated red cells may no longer be present, but hemosiderin is almost always found, though in varying amounts. The inflammatory infiltrate is less pronounced than in the early stage. The epidermis may show slight atrophy with absence of the rete ridges.

**Differential Diagnosis.** It may be difficult to differentiate purpura pigmentosa progressiva histologically from stasis dermatitis because inflammation, extravasation of erythrocytes and deposits of hemosiderin occur in both. However, the process extends much deeper in stasis dermatitis and in addition, fibrosis of the dermis and fibrous thickening of the walls of medium sized vessels in the lower dermis is commonly present (see page 73). Anaphylactoid purpura differs from purpura pigmentosa progressiva by the predominance of neutrophils in the infiltrate and the presence of leukocytoclasia (see page 128).

## RELAPSING FEBRILE NODULAR NONSUPPURATIVE PANNICULITIS (WEBER CHRISTIAN DISEASE)

This disease is characterized by the appearance of crops of indurated, tender nodules and plaques in the subcutaneous fat. As the lesions involute, they often leave a depression in the skin. The overlying skin shows, as a rule, no involvement other than mild erythema.

**Histopathology.** Histologically, the disease can be divided into three stages. The first two stages occur while there is induration clinically. During the third stage, depression of the skin develops. The first stage is observed only rarely because it is of short duration. It is probable that it does not always occur. Most sections show a combination of the changes of the second and the third stage.

In the first stage (acute inflammatory stage), there is, between the fat cells, an inflammatory infiltrate composed of polymorphonuclear leukocytes, lymphocytes and histiocytes. Polymorphonuclear leuko-




FIG. 62 Relapsing febrile nodular nonsuppurative panniculitis. The inflammatory infiltrate is composed of polymorphonuclear leukocytes, lymphocytes and histiocytes.

cytes may predominate (Fig 62) (Ungar Lever) Abscess formation does not occur

In the second stage (macrophagic stage) the infiltrate consists predominantly of histiocytes A few lymphocytes and plasma cells are present The histiocytes are seen invading and digesting the fat cells Such histiocytes (macrophages) are large and have a foamy cytoplasm




FIG 63 Relapsing febrile nodular nonsuppurative panniculitis Second and third stages The left side and the center of the field show the second stage foam cells (macrophages) invading and digesting the fat cells The right side shows the third stage replacement by fibrotic connective tissue ( $\times 100$ )

(Fig 63) Some of them are multinucleated In some areas numerous macrophages with foamy cytoplasm completely replace the fat cells

In the third stage (fibroblastic stage) fibroblasts intermingled with lymphocytes replace the macrophages Collagen is laid down with resulting fibrosis

The epidermis and the dermis show no involvement In some cases the subcutaneous vessels show pathologic changes such as edema and thickening of their walls (Cummins and Lever Tilden Gotshalk and Avakian)

In cases of liquefying panniculitis (Shaffer Binkley) the second macrophagic stage instead of being followed by fibrosis is followed

by liquefaction of the foam cell infiltrate. One finds an amorphous foamy matrix in which the nuclei of foam cells as well as lymphocytes and some polymorphonuclear leukocytes are suspended.

It is now established that the disease may affect the internal adipose tissue. Four autopsies are on record. In three of them internal lesions of panniculitis were present which had the same histologic appearance as the cutaneous lesions. Involvement was slight in one of these cases (Spain and Foley) but extensive in two (Ungar, Mostofi and Engleman). However, only one of these patients (Mostofi and Engleman) had died definitely as a result of this disease.

**Differential Diagnosis.** The histologic appearance of nodular panniculitis is diagnostic in the second stage because there is no other condition in which one finds such preponderance of foam cells in the subcutaneous fat. In foreign body granuloma one usually finds besides foam cells a great variety of other cells including foreign body giant cells.

### SCLEREMA NEONATORUM

Two types occur: generalized sclerema neonatorum and nodular sclerema neonatorum (subcutaneous fat necrosis of the newborn; adiponecrosis subcutanea neonatorum; lipophagic granuloma).

In generalized sclerema neonatorum the skin of the entire body has a waxlike appearance and is hard, dry and cold. Death usually occurs within a week. In nodular sclerema neonatorum one observes deep-seated indurated areas in the subcutaneous fat. The process is localized and self-limited; the lesions requiring about 4 months to disappear. The general health is not affected.

**Histopathology.** The fundamental lesion is the same in generalized sclerema neonatorum and in nodular sclerema neonatorum (Zeek and Madden, Flory) and both may occur simultaneously (Eichenlaub and Sandler). One finds degeneration, necrosis and crystallization of the subcutaneous fat together with an inflammatory reaction and fibrosis. It is possible that the disease is due to a delay in the maturation of the fat and that on account of this there is an unusually low oleic acid content of the fat so that it solidifies more easily than normally (Noonin, Pace and Davis).

**GENERALIZED SCLEREMA NEONATORUM.** The fat cells of the subcutis vary greatly in size and shape. In some areas they are necrotic and have lost their outline. Needle-shaped empty clefts lie singly or in radial arrangement inside of fat cells (Fig. 64). In frozen sections these clefts are found to be occupied by crystals. The crystals fail to stain with fat stains. Some of them are doubly refractile in the polarizing microscope (Reich, Zeek and Madden). The chemical composi-

tion of the crystals is not yet established fully. Most investigators regard the crystals as those of fatty acids (McIntosh). A granulomatous infiltrate is present mainly at the periphery of the fat lobules but also in scattered foci throughout the lobules. The infiltrate is composed of lymphocytes, histiocytes, foreign-body giant cells and fibroblasts. The septa between the fat lobules are considerably thickened.




FIG 64 Sclerema neonatorum (subcutaneous fat necrosis of the newborn). Several fat cells contain needle shaped clefts in a radial arrangement. These clefts are indicative of fat crystals. The fat crystals themselves are not visible because of fixation in Helly's solution, which dissolves fat. A granulomatous infiltrate containing foreign body giant cells is present between the fat cells ( $\times 100$ )

In two cases of generalized sclerema neonatorum, autopsy revealed in the visceral fat lesions which were histologically identical with those of the subcutaneous fat. In one case (Zeek and Madden), the lesions were widely distributed. In the other case (Flory), they were limited to the perirenal fat and to the fat about the ribs.

**NODULAR SCLEREMA NEONATORUM (SUBCUTANEOUS FAT NECROSIS OF THE NEWBORN).** One observes, as a rule, a more marked inflammatory reaction and a much larger number of foreign body giant cells than in generalized sclerema neonatorum (Fig 64). The foreign-

body giant cells may contain fat crystals (Fox) Healing takes place with fibrosis

**Differential Diagnosis** The histologic picture of sclerema neonatorum differs from that of relapsing febrile nodular nonsuppurative panniculitis by the presence of fat crystals and the absence of foam cells

### HEREDITARY EDEMA OF THE LEGS (MILROY'S DISEASE)

This disease occurs as a solid white indolent and persistent edema of the lower extremities Usually several members of a family are affected

**Histopathology** The dermis and especially the subcutaneous fat show severe interstitial edema There is an increase in the amount of collagen in the dermis and segmentation of the subcutaneous fat by thick strands of collagen The capillaries and the lymphatics are dilated and increased in number and show slight perivascular lymphocytic infiltration The walls of the larger blood vessels may be thickened by fibrosis

### CHONDRODERMATITIS NODULARIS CHRONICA HELICIS

In this disorder one or several small well-defined hard painful nodules are found on the upper margin of the ear The surface of the nodules often is hyperkeratotic After removal of the keratotic layer a small ulcer may be visualized

**Histopathology** The epidermis shows hyperkeratosis irregular acanthosis and usually also central ulceration The dermis and frequently also the perichondrium are permeated by a chronic inflammatory granulomatous infiltrate composed of lymphocytes plasma

and severity There are giant cells but occasionally a calcification so that the cartilage instead of staining blue stains homogeneously pink with hematoxylin and eosin in the affected areas

The pathogenesis is not clear Whereas most authors regard degenerative changes in the cartilage as the primary change (Foerster Ehemus) Newcomer recently has pointed out that similar degenerative changes occur in the aural cartilage with advancing age It is probable that trauma such as frostbite or pressure causes focal degeneration in the dermis of the ear Due to poor vascularization



in the affected area, repair does not ensue and the lesion becomes a focus of chronic inflammation.

### SCABIES

Scabies, which is caused by the itch mite *Acarus scabiei*, presents burrows as its characteristic lesion. The burrows, produced by the female mite, occur on the palmar surfaces of the hands and the fingers,



FIG 65 Scabies (Norwegian scabies) Two female mites are located entirely within the horny layer of the epidermis ( $\times 200$ ) (Robert N Buchanan, Jr, MD)

on the interdigital skin, on the flexor surfaces of the wrists, on the nipples of women and on the genitals of men. They appear as fine, angular or tortuous blackish threads, a few millimeters long. Often a vesicle is visible near the blind end of the burrow. In addition to the burrows, scabies presents a papular eruption which is usually most pronounced on the abdomen, the lower portions of the buttocks and the anterior axillary folds.

In especially susceptible individuals, the clinical picture of so called Norwegian scabies may result, showing diffuse hyperkeratosis, scaling and crusting of the skin and burrows everywhere on the skin, even on the face and the scalp.

**Histopathology** Histologic examination reveals the burrow in almost the whole of its course limited to the horny layer. Only the extreme blind end of the burrow is either in contact with or extends into the stratum malpighii. The female mite is situated at the blind end of the burrow (Fig. 62). Its head is bored more or less deeply into the stratum malpighii. Intracellular and intercellular edema is present in the stratum malpighii beneath the mouth parts of the mite to such extent that vesicle formation often results. The mite thus takes its food in fluid form. The dermis beneath the burrow shows a chronic inflammatory infiltrate which is composed predominantly of lymphocytes.

Norwegian scabies shows a much greater number of burrows than ordinary scabies so that almost every section shows several of them (Ingram).

## INSECT BITES

Insect bites especially when caused by ticks may cause persistent lesions which pose a diagnostic problem clinically as well as histologically. The clinical appearance of the lesions usually is that of a firm papule or of a nodule with or without central ulceration.

**Histopathology** The epidermis frequently shows pseudo epitheliomatous hyperplasia (see page 331). The dermis presents a dense granulomatous infiltrate which often extends into the subcutaneous fat. It consists principally of eosinophils and plasma cells admixed with lymphocytes and histiocytes. Some of the histiocytes occasionally show nuclei that are hyperchromatic, show mitotic figures or are binucleated. In some lesions large lymphoid follicles with germinal centers are observed (Winer and Strakosch, Allen). In rare instances parts of the insect are found in the dermis either free in the tissue surrounded by a foreign body reaction or within an epidermal inclusion cyst (Allen).

**Differential Diagnosis** If parts of the insect are present in the section the diagnosis is obvious. In their absence the true nature of the lesion may be missed easily. The pseudo epitheliomatous hyperplasia must be differentiated from squamous-cell carcinoma (see page 129). The presence of mitotic figures in the histiocytes together with the large number of eosinophils may suggest mycosis fungoides; the presence of lymphoid follicles may suggest lymphocytoma cutis; and the binucleated histiocytes if present require differentiation from the Sternberg-Reed cells of Hodgkin's disease. However the abundance of plasma cells in association with the eosinophils usually rules out lymphoma.

### LICHEN URTICATUS (PAPULAR URTICARIA)

This is a recurrent, pruritic eruption occurring, especially in children, during the summer months. The lesions consist of edematous




FIG. 66 Foreign body granuloma caused by a silk suture. The silk suture is located in the center of the field. Around it there is a severe inflammatory infiltrate containing numerous foreign body giant cells ( $\times 200$ ).

papules. Sensitivity to flea and bedbug bites is now generally regarded as the usual cause (Goldman; Shaffer, Spencer and Blank).

**Histopathology.** The appearance is nonspecific. Acanthosis and, in some cases, hyperkeratosis are present. The stratum malpighii shows intercellular as well as intracellular edema. A rather pronounced chronic inflammatory infiltrate is present around the vessels of the upper dermis.

## FOREIGN BODY GRANULOMAS

Many foreign substances when injected or accidentally implanted into the skin may cause a foreign body reaction. In addition certain



FIG. 67 Lipid granuloma caused by mineral oil (paraffinoma). The many large and small ovoid or round cavities which give the section a Swiss cheese appearance represent spaces filled with mineral oil (paraffin). (X200)

substances formed within the body may produce a foreign body reaction when deposited in the dermis or in the subcutaneous tissue. Examples are:

epithelioma  
cysts

Histopathology A typical foreign body reaction shows around the foreign material histiocytes and foreign body giant cells (for

their description, see page 36) In addition, lymphocytes and plasma cells are present (Fig 66) The histiocytes and the giant cells often exhibit phagocytic activity and then contain some of the foreign material

Some foreign body granulomas have a rather specific appearance Among them are lipid, tattoo, silicon and beryllium granulomas

### LIPID GRANULOMA (PARAFFINOMA)

Granulomas following injections of oily substances such as mineral oil (paraffin), cotton oil, or camphor oil occur as irregular, hard, nodular subcutaneous swellings Ulceration may develop

**Histopathology.** Histologically, lipid granulomas have a 'Swiss cheese' appearance because of the presence of numerous ovoid or round cavities (Fig 67) These cavities represent spaces occupied by the oily substance (Conrad and Weiss) The spaces between the cavities are taken up by an infiltrate composed largely of lymphocytes and some plasma cells In addition, there are groups of histiocytes some of which, due to the ingestion of lipid, have a foamy cytoplasm Variable numbers of foreign body giant cells are present In older lesions, fibrosis is prominent

Under the term *sclerosing lipogranuloma*, Smetana and Beinhard have reported as a post traumatic process subcutaneous granulomas with the same histologic picture as just described However, Best, Mason, DeWeerd and Dahlin have thrown serious doubt on the existence of such an entity Investigating two similar cases, they found by chemical analysis that the 'lipid' material was not body fat but mineral oil The 'lipid' material in their cases stained with Sudan 4 but not with osmic acid

### TATTOO GRANULOMA

**Histopathology.** Ordinary tattoos show diffusely scattered granules in the upper half of the dermis within phagocytes and lying free in the tissue, without any inflammatory reaction

However, if an inflammatory reaction occurs, due to allergy, the inflammatory infiltrate shows aside from the phagocytes numerous lymphocytes with an admixture of many eosinophils and a few plasma cells (Rostenberg, Brown and Cro)

### SILICON GRANULOMA

Silicon may be introduced into the skin through the contamination of lacerations with particles of soil Such wounds heal at first and then, many months or years later indurated nodules develop in the skin or the subcutaneous tissue

**Histopathology.** The histologic picture is indistinguishable from that of sarcoidosis (see page 188) except for the presence in some of the giant cells of colorless, spiculated, crystalline particles varying in size from 1 to 100 microns in length. When examined

#### BERYLLIUM GRANULOMA

Beryllium granulomas of the skin may form in two different ways (Grier, Nash and Freeman). They may arise as a manifestation of



FIG. 69 Beryllium granuloma caused by laceration with a fluorescent light tube. There is a large area of necrosis surrounded by tuberculoid granuloma tissue. In the walls several sarcoïd-like epithelioid-cell islands are present (X100).

systemic berylliosis, in which case it must be assumed that particles of beryllium reached the skin through the circulation, or they may develop following a laceration of the skin through which beryllium entered the tissue. Such lacerations were observed some years ago from cuts with fluorescent light bulbs.

their description, see page 36) In addition, lymphocytes and plasma cells are present (Fig 66) The histiocytes and the giant cells often exhibit phagocytic activity and then contain some of the foreign material

Some foreign body granulomas have a rather specific appearance Among them are lipid, tattoo, silicon and beryllium granulomas

### LIPID GRANULOMA (PARAFFINOMA)

Granulomas following injections of oily substances such as mineral oil (paraffin), cotton oil, or camphor oil occur as irregular, hard nodular subcutaneous swellings Ulceration may develop

**Histopathology.** Histologically, lipid granulomas have a "Swiss cheese" appearance because of the presence of numerous ovoid or round cavities (Fig 67) These cavities represent spaces occupied by the oily substance (Conrad and Weiss) The spaces between the cavities are taken up by an infiltrate composed largely of lymphocytes and some plasma cells In addition, there are groups of histiocytes some of which, due to the ingestion of lipid, have a foamy cytoplasm Variable numbers of foreign body giant cells are present In older lesions, fibrosis is prominent

Under the term *sclerosing lipogranuloma*, Smetana and Bernhard have reported as a post traumatic process subcutaneous granulomas with the same histologic picture as just described However, Best, Mason, DeWeerd and Dahlin have thrown serious doubt on the existence of such an entity Investigating two similar cases they found by chemical analysis that the 'lipid' material was not body fat but mineral oil The 'lipid' material in their cases stained with Sudan 4 but not with osmic acid

### TATTOO GRANULOMA

**Histopathology.** Ordinary tattoos show diffusely scattered granules in the upper half of the dermis within phagocytes and lying free in the tissue, without any inflammatory reaction

However, if an inflammatory reaction occurs due to allergy, the inflammatory infiltrate shows aside from the phagocytes numerous lymphocytes with an admixture of many eosinophils and a few plasma cells (Rostenberg Brown and Ciro)

### SILICON GRANULOMA

Silicon may be introduced into the skin through the contamination of lacerations with particles of soil Such wounds heal at first and then, many months or years later indurated nodules develop in the skin or the subcutaneous tissue



## Prurigo Nodularis

- Shaffer B and Beerman H Lichen simplex chronicus and its variants. Arch Dermat & Syph 64 310 1951

## Erythema Nodosum

- Grybowski M Sur l'anatomie pathologique de l'érythème noueux Bull Soc. franç. de dermat et syph. 45 1073 1938
- Lufgren S and Wahlgren F On the histopathology of erythema nodosum Acta dermat venereol 29 1 1949
- Miescher G Zur Frage der Radiarknoten beim Erythema nodosum Arch f Dermat. u Syph 193 251 1951
- Nahé M J Miescher's granulomas in erythema nodosum Dermatolog 101 80 1950
- Pautrier L M and Woringer F Contribution a l'étude histologique de l'érythème noueux Bull Soc. franç. de dermat et syph. 45 1073 1938
- Rotz

## Nodular Vasculitis

- Irgang S Nodular vasculitis Arch Dermat & Syph 67 185 1953
- Woodburne A R and Philpotts O S Nodular vasculitis Arch Dermat & Syph 60 291 1947

## Psoriasis

- Burks J W and Montgomery H Histopathologic study of psoriasis Arch Dermat & Syph 48 473 1945 (Good review)
- Macher G W and Foster P D Histopathogenesis of psoriasis and its aberrant lesions Arch Dermat & Syph 34 35 1936
- Swoughton R and Wells G A histochemical study on polysaccharides in normal and diseased skin J Invest Dermat. 14 37 1950

## Pustulosis Palmatis et Plantaris

- Andrews L C and Machacek G F Pustular bacteriids of the hands and feet Arch Dermat & Syph 32 837 1935
- Baxter H W Acrodermatitis continua vel perstans (dermatitis repens) and psoriasis pustulosa Brit. J Dermat 42 500 1930
- Bruck Acta dermat venereol 21 275 1941
- Ebert M H Psoriasiform eruption with pustular exacerbations Arch Dermat & Syph 27 933 1933
- Ebraham T N Pustular psoriasis Arch Dermat. & Syph 70 690 1935
- Lapierre J P Pustular psoriasis
- Nach
- (psoriasis pustulosa per
- and Scannone F So-called pustular psoriasis J Invest Dermat. 6 349 1945

## Acrodermatitis Continua

- Kogoj F Die spungiforme (schwammartige) Pustel Dermat. Wchnschr 107 1485 1938
- Das klinische und histologische Bild der Acrodermatitis continua Arch f Dermat. u Syph. 193 417 1941



nodules over which the skin remains intact. They are always few in number. The most serious lesion in systemic berylliosis is the chronic progressive pneumonitis which is frequently fatal.

The cutaneous granulomas following laceration show as first sign incomplete healing of the laceration, followed by swelling, induration and tenderness and, finally, central ulceration.

**Histopathology.** The cutaneous granulomas of systemic berylliosis often are indistinguishable from *sarcoidosis* (Hardy and Tabershaw). Occasionally, however, a moderate amount of caseation necrosis is present in some of the epithelioid cell islands. The quantity of beryllium present in the lesions is too small to be demonstrable by spectrographic analysis.

The cutaneous granulomas following laceration show, as a rule, more caseation necrosis than the granulomas of systemic berylliosis. The caseation necrosis not only may be present within some of the epithelioid cell islands but also may affect the entire center of the lesion (Fig. 68) (Neave, Frank and Tolmach). A collar of lymphocytes often surrounds some of the epithelioid cell islands, giving them the appearance of true tubercles. Occasionally, *Schaumann bodies*, just like those of *sarcoidosis*, are present (Grier, Nash and Freiman). The epidermis shows acanthosis and may show ulceration. In some cases the presence of beryllium has been demonstrated by spectrographic analysis (Dutra).

### SWIMMING POOL GRANULOMA

Abrasions in swimming pools may lead to circumscribed areas of nodular infiltration on the face resembling *lupus vulgaris* (Hellerstrom) or to granulomatous, verrucous lesions on the knee or elsewhere resembling *tuberculosis verrucosa cutis* (Rees and Bennett). The lesions usually heal within 3 to 9 months.

**Histopathology.** The histologic picture is similar to that of *lupus vulgaris* (Hellerstrom) or of *tuberculosis verrucosa cutis* (Rees and Bennett). Hellerstrom regarded his cases as instances of inoculation *lupus vulgaris* but inoculations of tissue into guinea pigs have given negative results in all reported cases and Hellerstrom found acid fast bacilli in the tissue in but one of his six cases. Thus a tuberculous cause is not established. Silicon granuloma is also unlikely because no crystalline particles were ever observed. The cause remains obscure.

### BIBLIOGRAPHY

#### *Urticaria*

Torok, L., and Lehner, E. Zur Anatomie und Pathogenese der Urticaria. Arch f Dermat u Syph 132:401, 1921

## Lichen Striatus

- Pinkus H. Lichen striatus and lichen planus. *J Invest Dermat* 11:9 1948  
 Senear F. E. and Caro M. R. Lichen striatus. *Arch Dermat. & Syph* 43:116 1944

## Pityriasis Rubra Pilaris

- Brunving L. A. and Sheard C. Dark adaptation in pityriasis rubra pilaris. *Arch Dermat. & Syph* 43:42 1941

## Granuloma Faciale

- Lever W. F. Eosinophilic granuloma of the skin. Its relation to erythema elevatum diutinum and eosinophilic granuloma of the bone. *Arch Dermat. & Syph* 55:194 1947  
 ——— Lane C. G. Downing J. G. and Spangler A. S. Eosinophilic granuloma of the skin. *Arch Dermat. & Syph* 58:430 1948  
 ——— and Leeper R. W. Eosinophilic granuloma of the skin. *Arch Dermat. & Syph* 62:85 1950  
 Pfleger L. and Tappener S. Über das eosinophile Granulom des Gesichtes. *Arch f Dermat u Syph* 193:1 1951  
 Pinkus H. Granuloma faciale. *Dermatolog* 105:85 1952

## Erythema Elevatum Diutinum

- Crocker H. R. and Williams C. Erythema elevatum diutinum. *Brit J Dermat.* 6:1 33 1894  
 Haber H. and Russell B. Erythema multiforme perstans (erythema elevatum diutinum). *Brit J Dermat* 62:460 1950  
 Ketron L. W. Erythema elevatum diutinum. *Arch Dermat. & Syph* 50:363 1944  
 Weidman F. D. and Besancon J. H. Erythema elevatum diutinum. *Arch Dermat. & Syph* 20:93 1929  
 Weiss R. S. Cooper Z. L. and Gottschalk H. R. Erythema elevatum diutinum. *Arch Dermat. & Syph* 58:716 1948

## Granuloma Annulare

- Bowers R. E. The histology of granuloma annulare compared with that of the necrobiotic nodules of rheumatoid arthritis. *Brit. J Dermat* 61:247 1949  
 Coolman M. H. and Ketron L. W. Granuloma annulare. *Arch Dermat. & Syph* 33:173 1936  
 Frantz F. C. and Montgomery H. Granuloma annulare. *Arch Dermat. & Syph* 46:391 1912 (Good histologic discussion)

## Subcutaneous Nodules of Rheumatic Fever and Rheumatoid Arthritis

- Bennett G. A. Zeller J. W. and Bauer W. Subcutaneous nodules of rheumatoid arthritis and rheumatic fever. *Arch Path* 30:70 1940

## Acrodermatitis Chronica Atrophicans

- Montgomery H. and Sullivan R. R. Acrodermatitis atrophicans chronica. *Arch Dermat. & Syph* 51:32 1915  
 Sweitzer S. F. and Layman C. W. Acrodermatitis chronica atrophicans. *Arch Dermat. & Syph* 31:196 1935

Lapière van Runkelen and Dussart Etude de quatre cas d'acrodermatite continue pustuleuse d'Hallopeau Arch belges dermat et syph 23 1939

### *Impetigo Herpetiformis*

Kogoj F Das klinische und histologische Bild der Acrodermatitis continua Arch f Dermat u Syph 193 117 1951

Rost G A Ueber Impetigo herpetiformis Arch f Dermat u Syph 131 538 1921

### *Parapsoriasis*

Carol W L L Prætken J R and Stigler W Parakerioma variegata (Parapsoriasis variegata seu reticulata) Acta dermat venereol 24 1 1943

Keil H Parapsoriasis en plaques dissimines and incipient mycosis fungoides Arch Dermat & Syph 37 165 1938 38 245 1938

McCarthy I Differential diagnosis of parapsoriasis Arch Dermat & Syph 45 81 1942

Miescher G Ein Fall von ausgedehnter mikropapulöser und papulokrustöser Parapsoriasis guttata Dermatol 91 181 1947

Montgomery H and Barkhart R J Parapsoriasis its relation to mycosis fungoides and tuberculosis Arch Dermat & Syph 46 673 1942 (Good review)

Robinson S S Pityriasis varioliformis acuta Arch Dermat & Syph 36 971 1937

Senear F I and Oliver E A Pityriasis lichenoides et varioliformis acuta (Harbermann) Arch Dermat & Syph 23 12 1931

### *Pityriasis Rosea*

Percival G H Pityriasis rosea Brit J Dermat 44 211 1952

### *Lichen Planus*

Blair D S Lichen planus pemphigoides Arch Dermat & Syph 58 159 1948

Feldman S Lichen planus et acuminatus atrophicus Arch Dermat & Syph 34 378 1936

Fox H Lichen planus confined to the mouth Arch Dermat & Syph 24 1071 1931

Little E G Lichen planus J Cutan Dis 37 639 1919

Sachs W and DeOreo G Lichen planopilaris Arch Dermat & Syph 45 1031 1942 (Good review)

Silver H Chargin I and Sachs P M Follicular lichen planus (lichen planopilaris) Arch Dermat & Syph 67 346 1953

Strauss M J Vesicular and bullous lichen planus Acta dermat venereol 14 447 1933

Winer L H and Levitt H Lichen planus in the Negro Arch Dermat & Syph 56 437 1947

### *Lichen Nitidus*

Ellis F A and Hill W R Is lichen nitidus a variety of lichen planus Arch Dermat & Syph 38 569 1938

Michelson H E Lichen nitidus Arch Dermat & Syph 7 763 1923

Niles H D Lichen nitidus Arch Dermat & Syph 22 687 1930

Wien M S and Perlstein M O Lichen nitidus Arch Dermat & Syph 30 790 1934

## Lichen Striatus

- Pinkus H Lichen striatus and lichen planus *J Invest Dermat* 11 9 1948  
 Senear F E. and Caro M R Lichen striatus *Arch Dermat & Syph* 43 116  
 1941

## Pityriasis Rubra Pilaris

- Brunsting L A and Sheard C Dark adaptation in pityriasis rubra pilaris  
*Arch Dermat & Syph* 43 42 1941

## Granuloma Faciale

- Lever W F Eosinophilic granuloma of the skin Its relation to erythema ele-  
 vatum diutinum and eosinophilic granuloma of the bone *Arch. Dermat &*  
*Syph* 53 191 1947  
 --- Lane C G Downing J G and Spangler A S Eosinophilic granuloma  
 of the skin *Arch Dermat & Syph* 58 430 1948  
 --- and Leeper R W Eosinophilic granuloma of the skin *Arch Dermat &*  
*Syph* 62 85 1950  
 Pfleger L. and Tappeiner S Über das eosinophile Granulom des Gesichtes  
*Arch f Dermat u Syph* 193 1 1951  
 Linkus H Granuloma faciale *Dermatolog* 105 85 1952

## Erythema Elevatum Diutinum

- Crocker H R and Williams C Erythema elevatum diutinum *Brit J Dermat*  
 61 33 1894  
 Haber H and Russell B Erythema multiforme perstans (erythema elevatum  
 diutinum) *Brit J Dermat* 62 460 1950  
 Ketron L W Erythema elevatum diutinum *Arch Dermat & Syph* 50 363 1944  
 Weidman F D and Besancon J H Erythema elevatum diutinum *Arch Der-*  
*mat & Syph* 20 593 1929  
 Weiss R S Cooper Z K and Gottschalk H R Erythema elevatum diutinum  
*Arch Dermat & Syph* 58 716 1948

## Granuloma Annulare

- Bowers R E The histology of granuloma annulare compared with that of the  
 necrobiotic nodules of rheumatoid arthritis *Brit J Dermat* 61 247 1949  
 Croodman M H and Ketron L W Granuloma annulare *Arch Dermat &*  
*Syph* 53 475 1936  
 Trunty F C and Montgomery H Granuloma annulare *Arch Dermat & Syph*  
 46 391 1942 (Good histologic discussion)

## Subcutaneous Nodules of Rheumatic Fever and Rheumatoid Arthritis

- Bennett G J Zeller J W and Bauer W Subcutaneous nodules of rheuma-  
 toid arthritis and rheumatic fever *Arch Path* 30 70 1940

## Acrodermatitis Chronica Atrophicans

- Montgomery H and Sullivan R R Acrodermatitis atrophicans chronica *Arch*  
*Dermat & Syph* 51 52 1945  
 Sweitzer S F and Layman C W Acrodermatitis chronica atrophicans *Arch*  
*Dermat & Syph* 51 196 1955

## Radiodermatitis

- Anderson N P and Anderson H E Development of basal cell epithelioma as a consequence of radiodermatitis Arch Dermat & Syph 63 586 1951
- Blom Ides C Sarcoma in Röntgenodermatitis Acta dermat venereol 30 47 1950
- Gentile H Malignant fibroblastic tumors of the skin Acta dermat venereol 31, suppl 27 1951
- Montgomery H Pathologic histology of radiodermatitis in Meeke G M X ray and Radium in the Treatment of Diseases of the Skin ed 3 pp 333 362 Philadelphia Lea & Febiger 1938 (Good review)
- Sims C F and Kirsch N Spindle cell carcinoma in chronic radiodermatitis A
- Warren S Effect of radiation on norm Path 35 340 1943
- Wolbach S B A summary of the effects of repeated roentgen ray exposures upon the human skin antecedent to the formation of carcinoma Am J Roentgenol 13 139 1925

## Solar Dermatitis

- Lamb J H Shelmire B Cooper Z Morgan R J and Keaty C Solar dermatitis Arch Dermat & Syph 62 1 1950

## Hidrocystoma

- Dostrovsky A and Sigler F Experimentally induced disappearance and reappearance of lesions of hidrocystoma J Invest Dermat 5 167 1942
- Kenedy D and Lehner E Ein Fall von Hidrocystom Arch f Dermat u Syph 142 95 1923

## Acne

- Laymon C W Lupoid rosacea Arch Dermat & Syph 63 409 1951
- and Schoch E P Jr Micropapular tuberculid and rosacea Arch Dermat & Syph 58 286 1948
- Lynch F W Acne vulgaris Arch Dermat & Syph 42 593 1940
- Miescher G Rosacea und Rosacea ähnliche Tuberkulide Dermatol 88 150 1943
- Montgomery H Acne necrotica miliaris of the scalp Arch Dermat & Syph 36 40 1937
- Sams W M Rhinophyma Arch Dermat & Syph 26 834 1932

## Fox Fordyce Disease

- Nilzén A Two cases of Fox Fordyce disease Acta dermat venereol 28 518 1948
- Roxburgh A C A case of Fox Fordyce disease Brit J Dermat 55 121 1913
- Shelley W B and Hurley H J Jr Methods of exploring human apocrine sweat gland physiology Arch Dermat & Syph 66 156 1952

## Alopecia

- Dillaha C J and Rothman S Therapeutic experiments in alopecia areata with orally administered cortisone JAMA 150 516 1952
- Laymon C W The cicatricial alopecias J Invest Dermat 8 99 1947
- Miescher G and Lengggenhager R Ueber Pseudopelade Brocq Dermatol 91 122 1917



- Noojin R O Price B F and Davis H G Subcutaneous fat necrosis of the newborn certain etiologic considerations J Invest Dermat 12 331 1949  
 Reich N E Sclerema neonatorum Arch Dermat & Syph 45 342 1912  
 Zeek P and Madden E M Sclerema adiposum neonatorum of both internal and external adipose tissue Arch Path 41 166 1916

#### Hereditary Edema of the Legs

- McGuire J and Zeek P Pathogenesis of chronic hereditary edema of extremities (Milroy's disease) J A M A 98 870 1932  
 Rosenberg W A Hereditary edema of the legs (Milroy's disease) Arch Dermat & Syph 42 1113 1910

#### Chondrodermatitis Nodularis Chronica Helicis

- Ebenius B Chondrodermatitis nodularis chronica auricularis Acta radiol 22 563 1911  
 Foerster O H Painful nodular growth of the ear Arch Dermat & Syph 11 149 1925  
 Newcomer V D Steffen C G Sternberg T H and Lichtenstein L Chondrodermatitis nodularis chronica helicis Arch Dermat & Syph 68 241 1953

#### Scabies

- Heilesen B Studies on *Acarus scabiei* and scabies Acta dermat venereol 16 suppl XIV 1946  
 Ingram J T Ward epidemic from Norwegian scabies Brit J Dermat 63 311 1951

#### Insect Bites

- Allen A C Persistent insect bites (dermal eosinophilic granulomas) simulating lymphoblastoma histiocytoses and squamous cell carcinomas Am J Path 24 367 1918  
 Tobias N Tickbite granuloma J Invest Dermat 12 255 1949  
 Winer I H and Strakosch I A Tickbites—*Dermacentor variabilis* (Say) J Invest Dermat 4 249 1941

#### Lichen Urticatus

- Goldman L Lichen urticatus syndrome Arch Dermat & Syph 58 74 1918  
 Shaffer B Spencer M C and Blink H Papular urticaria J Invest Dermat 11 293 1948

#### Foreign Body Granulomas

- Ayres W W Ober W B and Hamilton P K Post traumatic subcutaneous granulomas associated with a crystalline material Am J Path 27 303 1951  
 Best C W Mason H I DeWeerd J W and Duhlin D C Sclerosing lipogranuloma of the male genitalia produced by mineral oil Proc Mayo Clin 28 623 1953  
 Conrad A H Conrad A H Jr and Weiss R S Sesame oil tumors J A M A 121 237 1943  
 Dutra F R Beryllium granulomas of the skin Arch Dermat & Syph 60 1140 1949  
 Grier R S Nash P and Freeman D J Skin lesions in persons exposed to beryllium compounds J Ind Hyg & Toxicol 30 228 1918  
 Hardy H L and Tabershaw I R Delayed chemical pneumonitis occurring in

- workers exposed to beryllium compounds *J Ind Hyg & Toxicol* 28 197, 1916
- Neave H J Frank S B and Tolmach J Cutaneous granulomas following laceration by fluorescent light bulbs *Arch Dermat & Syph* 61 401, 1950
- Rostenberg A Jr Brown R A and Caro M R Discussion of tattoo reactions with report of a case showing a reaction to a green color *Arch Dermat. & Syph* 62 510 1950
- Smetana H F and Bernhard W Sclerosing lipogranuloma *Arch Path* 50 296 1950
- Commerville J and Milne J A Pseudo-tuberculoma silicoticum *Brit J Dermat* 62 105 1950
- Weidman F D The danger of liquid petrolatum in parenteral injections, *JAMA* 80 1761 1923

#### Swimming Pool Granuloma

- Hullerstrum S Collected cases of inoculation lupus vulgaris *Acta dermat venerol* 31 191 1951
- Rees R B and Lennett J H Granuloma following swimming pool abrasions *JAMA* 152 1606 1953



## 9

# Eruptions Due to Drugs

Allergic reactions to drugs may cause various eruptions identical in their clinical appearances to cutaneous diseases occurring also as idiopathic entities. Drugs may cause, for instance, urticaria, erythema multiforme, erythema nodosum, dermatitis, including generalized exfoliative dermatitis, purpura, folliculitis and periarthritis nodosa. The histologic picture is the same in these diseases whether they are due to a drug or occur in their idiopathic form.

Only histologic changes more or less specific for eruptions due to drugs will be discussed.

### FIXED DRUG ERUPTION

Fixed drug eruptions are circumscribed lesions which persistently recur at the same site at each administration of the allergenic drug. The most common type of fixed drug eruption consists of one or several slightly elevated, erythematous plaques which on healing leave pigmented areas. Fixed drug eruptions may occur after the ingestion of phenolphthalein and coal tar derivatives such as amino pyrine and acetophenetidin.

**Histopathology.** There is an increase in melanin in the basal cells and in the dendritic cells of the epidermis. Melanin is present in the upper dermis. It is found predominantly in histiocytes (melanophores) but also free in the tissue. The superficial capillaries are surrounded by a mild inflammatory infiltrate composed of lymphocytes and histiocytes (Weiss and Kile, Chargin and Leifer).

### QUINACRINE HYDROCHLORIDE (ATABRINE) DERMATITIS

Quinacrine hydrochloride (Atabrine) may cause a subacute or a chronic dermatitis, lichen planus like lesions and exfoliative dermatitis (Bereston). In some patients the eruption is followed by anhidrosis in the involved areas.

**Histopathology.** The histologic picture is usually that of a non specific chronic dermatitis. In lesions that clinically resemble lichen planus, the histopathologic picture often resembles that of lichen

planus too. As a rule, however, the bandlike infiltrate is not so dense as in lichen planus and may contain eosinophils which are not present in true lichen planus (Wilson). In many cases one observes hyperkeratosis with keratotic plugging of pilosebaceous follicles. Numerous melanophores may be present in the dermis (Alden and Frank).

In patients in whom anhidrosis develops in areas of quinacrine hydrochloride dermatitis one may observe atrophy of the dermal portions of sweat ducts and dilatation and atrophy of the sweat glands. Focal inflammation may be present about the sweat glands (Sulzberger, Herrmann and Zak).

### EXFOLIATIVE DERMATITIS DUE TO DRUGS

The most common drugs to cause exfoliative dermatitis are arsenamine, gold salts, the sulfonamides and phenobarbital.

**Histopathology** The exfoliative dermatitis caused by these drugs has the same histologic appearance as exfoliative dermatitis due to other causes (see page 74 and Fig. 27).

Changes in the internal organs may be present, such as interstitial myocarditis (Brown and McNamara, French and Weller, Winer and Baer), interstitial nephritis (Winer and Baer) and fatty degeneration of the liver with inflammatory infiltration about the portal canals (Winer and Baer).

### BROMODERMA

Prolonged ingestion of bromides may cause the formation of granulomatous verrucous plaques which are called bromoderma. They occur usually on the lower extremities.

**Histopathology** There are papillomatosis and considerable downward proliferation of the epidermis, often of such degree as to produce the picture of pseudoepitheliomatous hyperplasia.

#### *Chronic bromoderma* (Bloch and Fenchio) (Fig. 69)

The dermis shows an extensive granulomatous infiltrate which may reach down into the subcutaneous layer. It is composed of a great variety of cells including lymphocytes, plasma cells and histiocytes. Neutrophils are usually numerous and abscesses may be found scattered through the infiltrate. Eosinophils are few or absent. The blood vessels are increased in number, are dilated and show proliferation of their endothelium. Small areas of hemorrhage are often seen within the infiltrate.

**Differential Diagnosis** The histologic picture of bromoderma is suggestive but not diagnostic. Pyoderma gangrenosum may show at

the margin of an ulcer, an identical histologic picture, including intra epidermal abscesses. Intra epidermal abscesses are also observed in older lesions of pemphigus vegetans and in blastomycosis. Pemphigus vegetans differs from bromoderma by the large percentage of eosinophils in the intra epidermal abscesses and in the granuloma




FIG 69 Bromoderma. There is downward proliferation of the epidermis. A large intra epidermal abscess is shown. The dermis contains a granulomatous infiltrate ( $\times 50$ )

tous infiltrate (see page 80). Blastomycosis is easily differentiated by its numerous giant cells and the presence of yeast cells in them.

### IODODERMA

Iododerma is characterized by granulomatous lesions which may have the same clinical appearance as bromoderma. As a rule, however, there is less verrucous proliferation and greater tendency to ulceration.

**Histopathology.** Histologically iododerma differs from bromoderma by showing less epithelial proliferation. The granulomatous infiltrate frequently destroys the epidermis, resulting in ulceration. The infiltrate may be composed predominantly of histiocytes, some of which may show mitotic figures and hyperchromatic nuclei, so that

distinction from lesions of mycosis fungoides in the tumor stage may be almost impossible (Eller and Fox, Hollander and Fetterman)

### ARGYRIA

This condition caused by prolonged ingestion of silver salts or their local application to the mucous membranes, is characterized by



Fig. 70. Argyria. Silver granules are present in the membrane propria of the sweat glands. In some places the granules are so dense that they form a solid black band ( $\times 100$ )

bluish gray slate colored discoloration of the entire skin most pronounced on the exposed portions of the skin

**Histopathology.** Silver is found in the dermis predominantly extracellularly, where small round brownish particles of uniform size. It is never seen in the epidermis or its appendages. The silver particles measure less than 1 micron in diameter and lie singly as well as in clumps. Although visible in routine stains, they appear more clearly in sections stained lightly with polychrome methylene blue. However, the best method for the demonstration of silver granules is by dark field illumination. If sections are placed under a dark field microscope the silver appears as brilliantly refractile, white granules

against a dark background. Many more granules can be seen than with direct illumination.

The silver granules are most numerous in the membrane propria surrounding the sweat glands (Fig. 70) and in the subepidermal dermis. In addition silver particles may be seen in the connective tissue sheaths about the hair follicles and the sebaceous glands in the walls of blood vessels particularly their intima and adventitia in the arterioles pilorum and the nerves and diffusely scattered through the middle and the lower dermis. Elastic tissue stains reveals a predilection of the granules of silver for elastic fibers. The location of silver in elastic fibers explains the presence of fingerlike chains of granules projecting into the papillary bodies (Hill and Montgomery). In many cases one finds increased amounts of melanin in the basal layer of the epidermis and melanin laden chromatophores in the dermis.

Silver is deposited not only in the skin but also in internal organs. It is found particularly in the intima of blood vessels and in the connective tissue of the internal organs. Analogous to the marked involvement of the basement membrane of the sweat glands the basement membranes around the acini of the testes and of the choroid plexus are particularly rich in granules (Harker and Hunter).

**Differential Diagnosis.** Histologic differentiation of argyria from other kinds of pigmentation is made readily. Melanin and hemosiderin possess larger granules which lie to a great extent intracellularly in chromatophores and are nonrefractile with dark field illumination. In pigmentation due to mercury whether from the use of creams or in tattoo marks the mercury is deposited in large coarse granules throughout the epidermis and the dermis without any special affinity for the membrana propria of the sweat glands. For differentiation from chrysiasis see below.

### CHRYSIASIS

In chrysiasis which may follow the parenteral use of gold salts the skin of the exposed parts shows an ash gray discoloration.

**Histopathology.** Gold granules are light refractile with dark field examination like silver granules but they are larger and more irregular in size than silver granules. In contrast to silver granules they are found predominantly within cells. They lie in the endothelial and the perithelial cells of capillaries and in macrophages throughout the upper dermis. Only occasionally do granules lie free in the tissue spaces (Schmidt). In some instances gold particles are found in the basal cells of the epidermis (Kochs).

## ARSENICAL KERATOSIS AND CARCINOMA

Prolonged ingestion of inorganic arsenic frequently produces multiple arsenical keratoses which may progress into squamous-cell carcinoma. Recently it has been suggested that not only multiple cutaneous cancers but also multiple internal cancers may be caused (Sommers and McManus). Occasionally multiple superficial basal cell epitheliomas form in addition to arsenical keratoses (Anderson, Montgomery and Waisman).

Arsenical keratoses resemble senile keratoses in their clinical appearance. They may occur anywhere on the skin but are found most frequently on the palms and the soles in contrast with senile keratoses which predominate on the face and on the dorsa of the hands.

**Histopathology.** In early arsenical keratoses one observes hyperkeratosis associated with acanthosis and irregular downward proliferation of the rete ridges. There usually is some degree of disorder of the squamous cells and pyknosis of some of the nuclei. The histologic picture thus resembles that of senile keratosis (see page 327).

In more advanced lesions in addition to the above mentioned

of cells is outlined than in Bowen's disease. Although vacuolization of cells occurs also in Bowen's disease this feature is not so prominent as in arsenical keratosis. The presence of numerous vacuolated cells may be regarded as diagnostic of arsenical keratosis (Montgomery and Waisman). The vacuolated cells are twice or three times as large as normal squamous cells and possess small irregular deeply staining nuclei. They resemble the Paget cells of Paget's disease of the nipple except that intercellular prickles are usually present.

Ultimately through invasion of the dermis frank squamous cell carcinoma may develop (see page 329). Even in the invading type of carcinoma vacuolization, clumping and dyskeratosis remain prominent.

## BIBLIOGRAPHY

## Fixed Drug Eruption

- Chargin L and Eisler W Fixed eruptions due to the arspenamines J Invest Dermat 3 413 1910  
 Weiss R S and Kile R I Unusual phenolphthalein eruption Arch Dermat & Syph 32 915 1935

## Quinacrine Hydrochloride (Atabrine) Dermatitis

- Alden H S and Frank I J Atypical lichenoid dermatitis Arch Dermat & Syph 56 13 1917  
 Bereston E S Lichenoid dermatitis J Invest Dermat 7 69 1916  
 Sulzberger M B Herrmann I and Zak F G Studies on seething J Invest Dermat 9 221 1917  
 Wilson D J Eczematous and pigmentary lichenoid dermatitis atypical lichen planus preliminary report Arch Dermat & Syph 51 377 1916

## Exfoliative Dermatitis Due to Drugs

- Brown C E and McNamara D H Acute interstitial myocarditis following administration of arspenamines Arch Dermat & Syph 12 312 1910  
 French A J and Weller C A Interstitial myocarditis following the clinical and experimental use of sulfonamide drugs Am J Path 18 109 1912  
 Winer N J and Brier R I Exfoliative dermatitis due to phenobarbital Arch Dermat & Syph 13 173 1911

## Bromoderma

- Bloch B and Tenschio F Zur Klinik und Pathogenese des Brom derma vegetans Arch f Dermat u Syph 165 93 1932

## Iododerma

- Eller J J and Fox E C Fatal iododerma Arch Dermat & Syph 21 715 1931  
 Hollander L and Feiterman G H Fatal iododerma Arch Dermat & Syph 34 228 1936

## Argyria

- Harker J M and Hunter D Occupational argyria Brit J Dermat 17 111 1935  
 Hill W R and Montgomery H Argyria Arch Dermat & Syph 11 588 1911

## Chrysiasis

- Kochs A G Zur Kenntnis der Chrysiasis Arch f Dermat u Syph 178 323 1938  
 Schmidt O E I Chrysiasis Arch Dermat & Syph 11 116 1911

## Arsenical Keratosis and Carcinoma

- Anderson N P Bowen's precancerous dermatosis and multiple benign superficial epithelioma evidence of arsenic is an etiologic agent Arch Dermat & Syph 26 1052 1932  
 Montgomery H and Waisman M Epithelioma attributable to arsenic J Invest Dermat 1 365 1911  
 Sommers S C and McManus R G Multiple arsenical cancers of skin and internal organs Cancer 6 317 1953

# Degenerative Diseases

## SENILE DEGENERATION

Senile changes in skin not regularly exposed to light manifest themselves clinically in thinning of the skin and atrophy of the subcutaneous fat. In exposed skin the changes usually are much more pronounced than in skin that is not exposed and include marked

skin after minor trauma ecchymoses may occur (senile purpura) exposed to light senile rete ridges and atrophy changes in the collagen or in the elastic tissue are absent. Mild obliterative changes may be found in some of the vessels (Hill and Montgomery).

In skin exposed to sunlight two types of degenerative changes are seen which however represent one and the same process: basophilic degeneration of the collagen and senile elastosis. These changes limited to the upper third of the dermis may start as early as in the third decade of life and progress with age.

In basophilic degeneration the collagenous fibers appear broken up into amorphous clumps and granules which stain for

are stained with orcein as for elastic tissue. One then observes in the upper dermis separated from a somewhat atrophic epidermis by a narrow band of normal collagen masses of twisted thick black staining fibers (Fig. 71). Because these fibers give the same staining reactions as elastic tissue Unna referred to them as collastin thinking they were the result of a merging of collagen with elastic fibers. However recent studies by x-ray diffraction and electron microscopy suggest that the essential change in senile elastosis is degeneration of collagen fibers. Furthermore the degenerated material can be removed—in contrast with elastic fibers—by treatment of the sections with trypsin and at the same time the



## BIBLIOGRAPHY

## Fixed Drug Eruption

- Chargm L. and Leifer W. Fixed eruptions due to the arspenamines *J Invest Dermat* 3 443 1940  
 Weiss R S. and Kile R L. Unusual phenolphthalein eruption *Arch Dermat & Syph* 32 915 1935

## Quinacrine Hydrochloride (Atabrine) Dermatitis

- Alden H S. and Frank L J. Atypical lichenoid dermatitis *Arch Dermat & Syph* 56 13 1917  
 Bereston E S. Lichenoid dermatitis *J Invest Dermat* 7 69 1916  
 Sulzberger M B. Herrmann J. and Zink F G. Studies on sweating *J Invest Dermat* 9 221 1917  
 Wilson D J. Eczematous and pigmentary lichenoid dermatitis atypical lichen planus preliminary report *Arch Dermat & Syph* 51 377 1916

## Exfoliative Dermatitis Due to Drugs

- Brown C E. and McNamara D H. Acute interstitial myocarditis following administration of arspenamines *Arch Dermat & Syph* 42 312 1910  
 French A J. and Weller C V. Interstitial myocarditis following the clinical and experimental use of sulfonamide drugs *Am J Path* 18 109 1912  
 Winer N J. and Bier R L. Exfoliative dermatitis due to phenobarbital *Arch Dermat & Syph* 43 173 1911

## Bromoderma

- Bloch B. and Tenchio F. Zur Klinik und Pathogenese des Bromoderma vegetans *Arch f Dermat u Syph* 165 93 1939

## Iododerma

- Eller J J. and Fox E C. Fatal iododerma *Arch Dermat & Syph* 21 745 1931  
 Hollander L. and Fetterman G H. Fatal iododerma *Arch Dermat & Syph* 34 228 1936

## Argyria

- Harker J M. and Hunter D. Occupational argyria *Brit J Dermat* 17 111 1935  
 Hill W R. and Montgomery H. Argyria *Arch Dermat & Syph* 41 588 1911

## Chrysiasis

- Kochs A G. Zur Kenntnis der Chrysiasis *Arch f Dermat u Syph* 178 393 1938  
 Schmidt O E L. Chrysiasis *Arch Dermat & Syph* 44 416 1911

## Arsenical keratosis and Carcinoma

- Anderson N P. Bowen's precancerous dermatosis and multiple benign superficial epithelioma: evidence of arsenic as an etiologic agent *Arch Dermat & Syph* 26 1052 1932  
 Montgomery H. and Wismann M. Epithelioma attributable to arsenic *J Invest Dermat* 4 365 1911  
 Sommers S C. and McManus R G. Multiple arsenical cancers of skin and internal organs *Cancer* 6 317 1953

xerotica obliterans formerly regarded as independent diseases probably are identical with lichen sclerosus et atrophicus (see below)

**Histopathology** In lichen sclerosus et atrophicus one observes (1) hyperkeratosis with keratotic plugging (2) atrophy of the stratum



FIG. 72 Lichen sclerosus et atrophicus. There are hyperkeratosis with follicular plugging, atrophy of the stratum malpighii, marked lymphedema of the upper dermis and an inflammatory infiltrate in the mid-dermis. The edema in the subepidermal dermis is so marked that a bulla has resulted. ( $\times 100$ )

The hyperkeratosis is so marked that often the horny layer is thicker than the atrophic stratum malpighii. The stratum malpighii is reduced to a few layers of flattened cells. The cells of the basal layer show hydropic degeneration. The rete ridges often are completely

elastic staining properties of the skin disappear (Tattersall and Seville) This suggests that the increase in orcein staining material is caused not by an increase in the amount of elastic tissue but by a false staining reaction of degenerated collagen Staining of frozen sections with fat stains occasionally reveals numerous fine lipid droplets in the areas of basophilic degeneration or senile elastosis (Weidman Percival Hannay and Duthie)



FIG 71 Senile degeneration (senile elastosis) In the upper derm is separated from the epidermis by a narrow band of normal collagen there are masses of twisted thick degenerated collagen fibers staining black with orcein—just like elastic tissue ( $\times 100$ )

**Differential Diagnosis** For differentiation from pseudoxanthoma elasticum see page 56

### LICHEN SCLEROSUS ET ATROPHICUS

This disorder is characterized by flat topped white papules which coalesce to form white patches without any infiltration The surface of the lesions often shows comedo like plugs Occasionally the patches become bullous due to accumulation of fluid beneath the epidermis Lichen sclerosus et atrophicus occurs not only on the skin but also on the vulva and the glans penis Kraurosis vulvae and balanitis

wiped out but they may persist in some areas and even show some irregular downward proliferation. In such proliferations, hydropic degeneration of the basal cells usually is pronounced.

Beneath the epidermis there is a broad zone of pronounced edema. Within this zone the collagenous fibers are swollen and homogeneous and contain only a few nuclei. They stain poorly with eosin and other connective tissue stains. This change does not represent sclerosis (as the name of the disease would imply) but lymphedema. The hydropic degeneration of the basal cells together with the edema of the subepidermal collagen may lead to the formation of subepidermal bullae (Gottschalk and Cooper). These bullae may thus be classified as being due to degeneration of the basal cells (see Classification of Bullae page 66). The elastic fibers are sparse or even absent in the zone of edema (Nomland).

In the mid-dermis beneath the area of edema there is an infiltrate which usually is perivascular but at times assumes a bandlike formation. It is composed almost exclusively of lymphocytes. In lesions of long duration the infiltrate may have almost disappeared.

### KRAUROSIS VULVAE

A satisfactory classification of the atrophic lesions of the vulva has not yet evolved and to a certain degree the vagueness of the term kraurosis vulvae is responsible for this. Although some authors still regard it as an entity (Wallace and Whimster) others regard it as synonymous with lichen sclerosus et atrophicus (Layman).

The simplest classification of the atrophic lesions of the vulva that can be offered at present is

1 SENILE OR PRESENILE ATROPHY. There is atrophy of the vulvar mucosa but no stenosis of the vaginal orifice. There may be itching and due to scratching, pyulvitis may result.

2 LICHEN SCLEROSUS ET ATROPHICUS (KRAUROSIS VULVAE). Considerable atrophy with stenosis of the vaginal orifice is present. The lesions are whitish in color sharply demarcated and may extend to the inguinal folds and the perineal region. Itching may be present. Because the lesions have a whitish color on casual inspection they resemble leukoplakia but they lack the induration observed in leukoplakia. There has been uncertainty as to whether lichen sclerosus et

FIG. 74 Kraurosis vulvae (lichen sclerosus et atrophicus). There are edema of the upper dermis and a bandlike inflammatory infiltrate beneath it. In addition there is irregular downward proliferation of the rete ridges with hydropic degeneration of the basal cells. The latter feature is typical of lichen sclerosus et atrophicus and rules out leukoplakia. (X100)



FIG 73 Kraurosis vulvae (lichen sclerosus et atrophicus). There are hyperkeratosis atrophy of the stratum malpighii and marked lymph edema of the upper dermis with homogenization of the collagen ( $\times 100$ )



FIG 74. See facing page for legend.

wiped out but they may persist in some areas and even show some irregular downward proliferation. In such proliferations, hydropic degeneration of the basal cells usually is pronounced.

Beneath the epidermis there is a broad zone of pronounced edema. Within this zone the collagenous fibers are swollen and homogeneous and contain only a few nuclei. They stain poorly with eosin and other connective tissue stains. This change does not represent sclerosis (as the name of the disease would imply) but lymphedema. The hydropic degeneration of the basal cells together with the edema of the subepidermal collagen may lead to the formation of subepidermal bullae (Gottschalk and Cooper). These bullae may thus be classified as being due to degeneration of the basal cells (see Classification of Bullae, page 66). The elastic fibers are sparse or even absent in the zone of edema (Nomland).

In the mid dermis beneath the area of edema there is an infiltrate which usually is perivascular but at times assumes a bandlike formation. It is composed almost exclusively of lymphocytes. In lesions of long duration the infiltrate may have almost disappeared.

### KRAUROSIS VULVAE

A satisfactory classification of the atrophic lesions of the vulva has not yet evolved and to a certain degree the vagueness of the term kraurosis vulvae is responsible for this. Although some authors still regard it as an entity (Wallace and Whimster) others regard it as synonymous with lichen sclerosus et atrophicus (Laymon).

The simplest classification of the atrophic lesions of the vulva that can be offered at present is

1 SEVILE OR PRESEVILE ATROPHY. There is atrophy of the vulvar mucosa but no stenosis of the vaginal orifice. There may be itching and due to scratching, vulvitis may result.

2 LICHEN SCLEROSUS ET ATROPHICUS (KRAUROSIS VULVAE). Considerable atrophy with stenosis of the vaginal orifice is present. The lesions are whitish in color, sharply demarcated and may extend to the inguinal folds and the perianal region. Itching may be present. Because the lesions have a whitish color on casual inspection they resemble leucoplakia.

FIG. 74 Kraurosis vulvae (lichen sclerosus et atrophicus). There are edema of the upper dermis and a bandlike inflammatory infiltrate beneath it. In addition there is irregular downward proliferation of the rete ridges with hydropic degeneration of the basal cells. The latter feature is typical of lichen sclerosus et atrophicus and rules out leukoplakia. (x100)

atrophicus, a perfectly harmless lesion elsewhere, may progress into leukoplakia and squamous cell carcinoma when occurring on the vulva. Such instances recently have been reported (Wallace and Whimster), but it seems to be rare. Nevertheless, as Wallace and Whimster state, any state of atrophy occurring on the vulva has malignant potentialities.

**3. LEUKOPLAKIA** One or several whitish indurated plaques are present. Leukoplakia may occur independently or secondary to senile atrophy of the vulva. Since leukoplakia is an early malignant lesion ("squamous cell carcinoma, grade  $\frac{1}{2}$ ," see page 328), development of squamous cell carcinoma is common.

**Histopathology.** The histologic picture of senile or presenile atrophy is nonspecific, consisting of atrophy of the epidermis and a mild to moderately severe chronic inflammatory infiltrate in the upper dermis.

**Lichen sclerosus et atrophicus of the vulva** (kraurosis vulvae) shows the same histologic picture as on the skin with the exception that lesions located on the mucous membranes show no keratotic plugs (Fig. 73). Irregular downward proliferation of the rete ridges in which the basal cells show considerable hydropic degeneration occur more often than on the skin (Fig. 71). They should not be confused with the early atypical proliferations of leukoplakia which do not contain such hydropic basal cells.

**Leukoplakia** shows atypicality of the epidermis with irregular downward proliferation of atypical cells (see page 328). The hydropic degeneration of the basal cells seen in lichen sclerosus et atrophicus is absent.

## BALANITIS XEROTICA OBLITERANS

Balanitis xerotica obliterans represents lichen sclerosus et atrophicus of the glans and prepuce (Leymon). It is a chronic, progressive atrophic process which frequently eventuates in urethral stenosis. In very rare instances carcinoma may supervene (Gruz).

**Histopathology** The histologic picture is that of lichen sclerosus et atrophicus. Because of the absence of follicles in the areas of involvement, no keratotic plugging occurs.

## STRIAE DISTENSAE

Striae distensae occur in pregnancy and obesity, and particularly in Cushing's disease. They represent linear areas of cutaneous atrophy.

**Histopathology.** In the early lesions, the elastic fibers are transformed into numerous faintly staining fibrillae due to fraying of the elastic fibers. In addition, there are a mild perivascular infiltrate and

a distortion of the collagen bundles. Old lesions show almost complete absence of elastic fibers in the center. At the margin coiled and clumped elastic fibers are interspersed with fine, poorly staining fibrillae.

On the basis of these findings Ebert has concluded that striae distensae are not due to mechanical tearing of elastic fibers alone but that degeneration of the elastic fibers precedes their disappearance in the center of the striae.

### MACULAR ATROPHY (ANETODERMA)

Macular atrophy of Schweninger and Buzzi or anetoderma is characterized by atrophic oval patches located particularly on the trunk. The skin of the patches is thin and bluish white and bulges slightly. The lesions may give to the palpating finger the same sensation as does a hernial orifice. Early lesions may show mild erythema.

In the literature a difference has been made between a primary idiopathic type of macular atrophy and a secondary type developing in patients with diseases such as syphilis, lupus erythematosus and acrodermatitis chronica atrophicans but in sites not clinically affected by these diseases (Scully and Nomland). It is likely that such association is merely coincidence.

**Pathology.** There is a gradual disappearance of the elastic tissue throughout the dermis. The collagen remains unaffected. Thus a diagnosis of this disease can be made only from sections stained for elastic tissue.

### ATROPHODERMA RETICULATUM (FOLLICULITIS ULERYTHEMATOSA RETICULATA)

The eruption which is limited to the sides of the face consists of numerous small closely set areas of atrophy separated by narrow ridges of normal skin resulting in a reticulated appearance.

**Histopathology.** The epidermis shows diminution of the number of rete ridges and follicular plugging. Horn cysts caused by keratinization of hair follicles are present in the dermis. Sebaceous glands are few in number and small in size. In addition one observes degeneration of the collagen and fragmentation of the elastic fibers.

### COLLOID MILIUM (COLLOID DEGENERATION OF THE SKIN)

Colloid milium is characterized by pinhead sized round sharply circumscribed yellowish nodules of the skin. The nodules have a



translucent appearance and on puncture give exit to a soft gelatinous mass. The forehead is the site of predilection.

In occasional instances instead of small nodules large plaque-like lesions are present. To such cases the term colloid degeneration of the skin has been applied (Reuter and Becker).

**Histopathology** Histologically colloid milium is characterized by the presence of circumscribed masses of colloid material in the sub




FIG. 75. Colloid milium. In the upper dermis separated from the epidermis by a narrow zone of normal collagen there are two large round spaces incompletely filled with a fine mass of homogeneous material. This material represents colloid. The colloid contains a moderate number of nuclei ( $\times 100$ ).

epidermal region. It is assumed by many that colloid represents a form of collagen degeneration. Pirakken has observed that collagen before becoming colloid passes through the stage of basophilic degeneration. However, it is possible that colloid like amyloid represents a deposit rather than a product of collagen degeneration (Percival and Dutline).

The epidermis shows hyperkeratosis and atrophy of the stratum malpighii. Large, newly formed spaces lie close to the epidermis separated from it by only a narrow zone of normal collagen. These spaces extend through the upper third of the dermis and are sharply demarcated by collagen bundles arranged circularly around them.

They are filled incompletely by a fissured mass of homogeneous appearing material containing a moderate number of nuclei (Fig 75). The material referred to as colloid usually stains eosinophilic with hematoxylin and eosin though to a lesser degree than normal collagen. Occasionally however it stains faintly basophilic. The fixing of the colloid is due to fixation and dehydration. The nuclei within the colloid are well preserved and represent fibroblasts. Elastic tissue stains reveal elastic fibers within the masses of colloid however they are fragmented and fewer in number than in normal collagen.

Amorphous degeneration of the skin colloid is present not in superficial circumscribed foci but diffusely throughout the dermis (Reiter and Becker).

Differential Diagnosis Differentiation of colloid milium from amyloidosis (see page 274) requires special staining since these two substances greatly resemble each other morphologically and usually stain alike a pale pink with hematoxylin and eosin. One may use van Gieson's stain which stains colloid yellow and amyloid pink or methyl violet with which only amyloid is stained.

## BIBLIOGRAPHY

### Senile Degeneration

- Hill W R and Montgomery H Regional changes and changes caused by age in normal skin J Invest Dermat 5 321 1940  
 Percival G H Hannay P W and Dutue D A Fibrous changes in the dermis with special reference to senile elastosis Brit J Dermat 61 69 1949  
 Tattersall R A and Seville R Senile purpura Quart J Med 19 151 1950  
 Linn F G Die Histologie der Hautkrankheiten pp 993 994 Berlin Hirschwald 1894  
 Weidman F D The pathology of the yellowing dermatoses Arch Dermat & Syph 21 94 1931

### Lichen Sclerosus et Atrophicus

- Cuttschall H R and Cooper Z k lesions and extensive involvement  
 Montgomery H and Hill W R L Arch Dermat & Syph 42 755 1940 (Good review)  
 Nomland R In discussion of Montgomery H and Hill W R Arch Dermat & Syph 42 755 1940

### Kraurosis Vulvae

- Layman C M Lichen sclerosus et atrophicus and related disorders Arch Dermat & Syph 64 690 1951  
 Seville F O and Pound A W Lichen sclerosus et atrophicus in relation to the diagnosis leukoplakia and kraurosis of the vulva Austral J Dermat 1 168 1952  
 Wallace E G and Nomland R Lichen sclerosus et atrophicus of the vulva Arch Dermat & Syph 57 940 1948

translucent appearance and on puncture give exit to a soft gelatinous mass. The forehead is the site of predilection.

In occasional instances instead of small nodules large plaque-like lesions are present. To such cases the term colloid degeneration of the skin has been applied (Reiter and Becker).

**Histopathology** Histologically colloid milium is characterized by the presence of circumscribed masses of colloid material in the sub




FIG. 75 Colloid milium. In the upper dermis separated from the epidermis by a narrow zone of normal collagen there are two large round spaces incompletely filled with a fused mass of homogeneous material. This material represents colloid. The colloid contains a moderate number of nuclei ( $\times 100$ ).

epidermal region. It is assumed by many that colloid represents a form of collagen degeneration. Prickren has observed that collagen before becoming colloid passes through the stage of basophilic degeneration. However it is possible that colloid like amyloid represents a deposit rather than a product of collagen degeneration (Percival and Dutlue).

The epidermis shows hyperkeratosis and atrophy of the stratum malpighii. Large nearly round spaces lie close to the epidermis separated from it by only a narrow zone of normal collagen. These spaces extend through the upper third of the dermis and are sharply demarcated by collagen bundles arranged circularly around them.

# 11

## Bacterial Diseases

### IMPETIGO

Impetigo may be caused by streptococci or by staphylococci. Bullae are the primary lesion in either case. In the streptococcal variety, the bullae soon rupture leaving sharply demarcated erosions which be-




FIG. 76 Impetigo. A subcorneal vesicle filled mainly with neutrophils is present. The epidermis shows spongiosis. Many neutrophils are seen migrating through the epidermis ( $\times 200$ ).

come covered with heavy, honey colored crusts. In the staphylococcal variety, the bullae are more durable and dominate the clinical picture. Staphylococcal impetigo occurs especially in the newborn (formerly also called pemphigus neonatorum).

**Histopathology** In both the streptococcal and the staphylococcal variety of impetigo, the bulla arises directly beneath the horny layer

Wallace H J and Whimster I W Vulval atrophy and leukoplakia Brit J Dermat 63 241 1951 (Good histologic discussion)

#### Balanitis Xerotica Obliterans

Grutz O Beitrage zur Klinik der Balanitis xerotica obliterans Dermat Wchenschr 105 1206 1937

Laymon C M Lichen sclerosus et atrophicus and related disorders Arch Dermat & Syph 61 620 1951

Welton D G and Nowlin P Balanitis xerotica obliterans Arch Dermat & Syph 59 636 1949

#### Striae Distensae

Ebert M H Hypertrophic striae distensae Arch Dermat & Syph 28 825 1933

#### Macular Atrophy (Aetoderma)

Butterworth T Multiple benign tumor like growths of the skin Arch Dermat & Syph 29 893 1931

Scull R H and Nomland R Secondary macular atrophy Arch Dermat & Syph 36 809 1937

#### Folliculitis Ulerythematosia Reticulata

MacKee G M and Paroungian M B Folliculitis ulerythematosia reticulata J Cutan Dis 36 339 1918

Wertheimer H G Folliculitis ulerythematosia reticulata Arch Dermat & Syph 25 1060 1932

Winer L H Atrophoderma reticulatum Arch Dermat & Syph 34 980 1936

#### Colloid Milium

Percival G H and Duthie D A Notes on a case of colloid pseudomilium Brit J Dermat 60 399 1948

Pricklen J R Colloid and senile degeneration of the skin Acta dermu venereol 31 713 1951

Reuter M J and Becker S W Colloid degeneration (collagen degeneration) of the skin Arch Dermat & Syph 46 695 1942

Wray S C Colloid milium Arch Dermat & Syph 45 1148 1942

**IMPETIGO BOCKHART** presents a subcorneal pustule situated in the opening of a hair follicle. The upper portion of the hair follicle is surrounded by a considerable inflammatory infiltrate containing varying numbers of polymorphonuclear leukocytes.

A **FURUNCLE** presents on histologic examination a perifollicular abscess composed of a dense mass of polymorphonuclear leukocytes with a few lymphocytes. The hair with its follicle and its sebaceous glands is destroyed in the process.

In **FOLLICULITIS BARBAE** the perifolliculitis is as a rule less acute than in furunculosis and does not necessarily result in a perifollicular abscess. Frequently the infiltrate around the follicle contains relatively few polymorphonuclear leukocytes but consists mainly of lymphocytes, plasma cells and histiocytes. Some of the sebaceous glands undergo destruction but the hair follicles may escape destruction. Foreign body giant cells frequently are present around the hair follicles and the remnants of sebaceous glands. In many cases instead of being limited to the vicinity of the follicles the infiltrate extends through the entire upper dermis.

### FOLLICULITIS KELOIDALIS (DERMATITIS PAPILLARIS CAPILLITIS)

Folliculitis keloidalis represents a chronic folliculitis resulting in keloidal scarring. It occurs on the nape of the neck in men. In early cases one observes follicular papules, pustules and occasionally abscesses. The lesions are replaced gradually by fibrous nodules.

**Histopathology.** Early lesions show the same histopathologic picture as that of a furuncle. Older lesions show chronic granulation tissue containing many fibrocytes and lymphocytes and eosinophils. In the later stages the histologic picture is the same as in keloid, showing bundles of hypertrophic and sclerotic collagen.

### HIDRADENITIS SUPPURATIVA

Hidradenitis suppurativa represents a chronic staphylococcal infection of the apocrine glands in the axillary or the pubic regions. Early lesions consist of red tender nodules which become fluctuating and finally discharge pus. Ulcers may develop and healing takes place with considerable scarring.

**Histopathology.** The infection enters by way of the follicles and excretory ducts of the apocrine glands. The earliest cellular reaction is encountered in the subcutaneous tissue within and around the

(see "Classification of Bullae," page 64) The bulla contains fibrin polymorphonuclear leukocytes and some lymphocytes (Fig 76) With Giemsa's or Gram's stain, groups of cocci can be recognized within the bulla. They may lie extracellularly or within neutrophils.

The stratum Malpighii underlying the bulla shows spongiosis. Many neutrophils can be seen migrating through it. The upper dermis contains a moderately severe inflammatory infiltrate of neutrophils and lymphocytes.

At a later stage, when the bulla has ruptured, the horny layer is absent, and a crust composed of fibrin and neutrophils may be found resting on the stratum malpighii.

### ERYSIPFLAS

Erysipelas is a localized acute inflammation of the skin caused by streptococci. It is characterized by the presence of well demarcated dusky red areas with or without edema and vesiculation. A typical sign is the advancing red, raised and indurated border.

**Histopathology.** The dermis shows marked edema and dilatation of lymphatics and capillaries. When the edema is intense subepidermal bullae may be present. There is a marked diffuse infiltrate extending throughout the dermis and, occasionally, into the subcutaneous fat. It is composed chiefly of polymorphonuclear leukocytes and some lymphocytes. If sections are stained with Giemsa's or Gram's stain, streptococci are found chiefly in the lymphatics, but also distributed in the tissue.

### FOLLICULITIS (IMPETIGO BOCKHART, FURUNCLE, FOLLICULITIS BARBAE)

Pustular folliculitis may occur in three forms: as impetigo Bockhart, as furuncle and as folliculitis barbae (sycosis vulgaris). All three are caused by staphylococci.

Impetigo Bockhart represents a superficial pustular folliculitis and is characterized by an eruption of pustules, many of which are pierced by a hair.

A furuncle is a deep-seated folliculitis leading to a perifollicular cellulitis and terminating in suppuration and necrosis.

Folliculitis barbae is a deep-seated folliculitis peculiar to the bearded region. At first there are follicular papules and pustules which are followed by diffuse erythema, crusting and infiltration of the skin. Abscesses may be present or absent.

**Histopathology.** These three diseases cannot always be differentiated on a histologic basis.

Osler nodes are erythematous slightly raised tender intracutaneous nodes averaging 5 mm in size. They occur most frequently on the fingertips and as a rule last 4 or 5 days.

Janeway lesions are small macular or papular lesions measuring from 1 to 4 mm in diameter. They are usually red but may be partially hemorrhagic. They occur most commonly on the palms and the soles. Unlike Osler nodes they are never tender.

**Histopathology** The petechiae show the histologic picture of an inflammatory purpura (see page 127) presenting severe vasculitis in addition to the extravasation of red cells. One observes marked endothelial proliferation of capillaries leading to narrowing or even obliteration of the lumina. An infiltrate composed of polymorphonuclear leukocytes, lymphocytes and histiocytes surrounds and invades the capillaries. Streptococci have never been found in the lesions (Merklen and Wolf, Cornil, Mosinger and Jouve).

The Osler nodes show involvement not only of the capillaries but also of the dermal arterioles and venules which show intense endothelial proliferation and not infrequently partial or complete occlusion of their lumina by secondary thrombosis. A dense infiltrate of polymorphonuclear leukocytes, lymphocytes and histiocytes is present in the walls of the vessels and in the perivascular areas. The center of the node may show necrosis. Extravasation of erythrocytes is absent (Cornil, Mosinger and Jouve).

The Janeway lesions resemble the petechiae in their histologic appearance except that extravasation of erythrocytes is less evident.

## MENINGOCOCCEMIA (WATERHOUSE FRIDERICHSEN SYNDROME)

In fulminating septicemic infections with *Neisseria meningitidis* purpura is common. Death may occur within from 12 to 24 hours with few or no signs of meningeal involvement. On autopsy hemorrhages are

Ma

(

Waterhouse Friderichsen syndrome occurs in septicemic infections with organisms other than *Neisseria meningitidis* such as *Streptococcus hemolyticus* and *Pseudomonas aeruginosa* B. pyocyaneus)

**Histopathology** The purpura of meningococemia is caused by degeneration and inflammatory invasion of the walls of blood vessels (vasculitis) (see page 127). The vessels are dilated, engorged with blood and frequently thrombosed. The endothelial cells are swollen and desquamating and the vessel walls show necrosis. Polymorphonuclear



lumina of apocrine glands. The infection extends through the subcutaneous fat by way of the lymphatics to other apocrine glands as well as to eccrine glands. The lymph vessels are distended and contain many leukocytes and clumps of cocci. In the early stage, the infiltrate is composed predominantly of neutrophils. Later on, lymphocytes and plasma cells predominate and foreign body giant cells may be present. The upper parts of the dermis and the epidermis are not involved until extensive destruction has occurred throughout the subcutis.

### PYODERMA GANGRINOSUM (CHRONIC UNDERMINING BURROWING ULCER)

Pyoderma gangrenosum begins with cutaneous abscesses which break down, forming ulcers. The ulcers spread peripherally. The advancing border of the ulcers is purplish red, edematous and undermined. The condition is associated not infrequently with ulcerative colitis.

**Histopathology.** The histologic appearance is not diagnostic. In the region of the ulcer, the epidermis is absent. The upper dermis shows necrosis and is permeated by an acute inflammatory infiltrate. Farther down, the infiltrate is chronic inflammatory, granulomatous in nature, consisting of lymphocytes, neutrophils, plasma cells, histiocytes and fibroblasts. Epithelioid cells and foreign body giant cells

1

The infiltrate may extend deeply into the subcutaneous layer. In areas of healing, fibrosis occurs.

The epidermis at the edge of the ulcer often shows considerable proliferation, so that the histologic picture of pseudoepitheliomatous hyperplasia (see page 334) may result. Intraepithelial abscesses may occur in this region.

**Differential Diagnosis.** Any ulcer, whatever its genesis, may present the histologic picture just described. If the biopsy specimen is taken from an area in which the epidermis shows marked hyperplasia, differentiation from bromoderma may be impossible, since bromoderma too shows marked epithelial hyperplasia and intraepithelial abscesses.

### SUBACUTE BACTERIAL ENDOCARDITIS

Three types of cutaneous lesions may occur in subacute bacterial endocarditis: petechiae, Osler nodes and Janeway lesions (Libman and Friedberg).

In old lesions, no pustulation remains and the greatly thickened horny layer consists largely of fully keratinized cells with only a few areas of parakeratosis (Herold and Smith)



FIG 77 Keratosus blennorrhagica late stage Low magnification There is a very thick parakeratotic horny layer (H L) permeated by numerous neutrophils The upper stratum malpighii is the seat of a spongiform pustule (SP) The spongiform appearance is caused by the preservation of the cellular membranes of epidermal cells within the pustule There are elongation of the rete ridges and edema of the papillae ( $\times 100$ )

Gonococci can be demonstrated only rarely in old lesions

... cases and in smears from the lesions in 5 Only one case is on record in which gonococci were recovered on direct culture of the lesions (Margolin) It is likely that many of the cases reported in the literature as keratosus blennor

clear leukocytes are found in the damaged vascular walls as well as perivascularly. Large and small areas of hemorrhage are present in the tissue.

Meningococci can be demonstrated in the cytoplasm of endothelial cells and polymorphonuclear leukocytes. They are also found free in the lumina of vessels and in the perivascular spaces. In addition the thrombi frequently contain meningococci. Although these organisms can be recognized in routine stains, they are best demonstrated by Giemsa stain.

### KERATOSIS BLENNORRAGICA (GONORRHEAL KERATOSIS)

Keratosis blennorrhagica occurs in chronic gonorrhea usually in association with urethral discharge and polyarthritis. The lesions have a predilection for the palms, the soles, and the glans penis. Early lesions are represented by pustules. Gradually, the lesions become covered with thick, horny crusts. Confluence of neighboring lesions leads to the formation of extensive horny excrescences, which have been compared with mountain ranges on a relief map.

Reiter's disease, which in typical cases consists of the triad of urethritis, arthritis and conjunctivitis, may present cutaneous lesions identical with those of keratosis blennorrhagica, but gonorrhea is not the cause of the disease.

**Histopathology.** The first histologic changes consist of the appearance of an acute inflammatory infiltrate in the dermis and the formation of pustules in the uppermost epidermis (Carr and Friedman, Epstein and Chambers). As a rule, the pustules are spongiform (Fig. 77) (Kogoj) as in acrodermatitis continua and in impetigo herpetiformis (see pages 102 and 104). The spongiform appearance of the pustules is caused by the presence of neutrophils inside of edematous degenerated epidermal cells whose cellular membranes traverse the pustule like the network of a sponge (Fig. 78).

Simultaneously with or shortly after the formation of the pustules the rete ridges become elongated. At this stage, the histologic picture is indistinguishable from that of acrodermatitis continua.

However, as the lesions age the parakeratotic horny layer thickens to a degree that is not observed in acrodermatitis continua and is typical of keratosis blennorrhagica. The greatly thickened horny layer is the anatomic substrate for the mountain relief appearance of the lesions observed clinically. The horny layer may measure several millimeters in thickness. It consists of parakeratotic cells intermingled with neutrophils (Fig. 77).

In old lesions no pustulation remains and the greatly thickened horny layer consists largely of fully keratinized cells with only a few areas of parakeratosis (Herold and Smith)



FIG. 11. KeratosiBlenorrhagica late stage. Low magnification. There is a very thick parakeratinic horny layer (HL) permeated by numerous neutrophils. The upper stratum malpighii is the seat of a spongiform pustule (SP). The spongiform appearance is caused by the preservation of the cellular membranes of epidermal cells within the pustule. There are elongation of the rete ridges and edema of the papillae ( $\times 100$ ).

Gonococci can be demonstrated only rarely in the lesions of keratosiBlenorrhagica. Male and Singletary found that out of a total of 115 cases reported in the literature Gram negative diplococci were demonstrated in tissue sections in 6 cases and in smears from the lesions in 5. Only one case is on record in which gonococci were recovered on direct culture of the lesions (Margolin). It is likely that many of the cases reported in the literature as keratosiBlenorrhagica

rhagica were nongonorrheal and represented instances of Reiter's disease (Kuske)

In REITER'S DISEASE the histologic picture is the same as in keratitis blennorrhagica (Lever and Crawford)




FIG. 78 Keratitis blennorrhagica. Late stage. High magnification of Figure 77. The spongiform nature of the pustule is well apparent ( $\times 400$ )

### TUBERCULOSIS

When a normal not previously infected guinea pig is inoculated intracutaneously with an adequate dose of tubercle bacilli a hard nodule develops at the site of inoculation after from 8 to 12 days. The nodule soon ulcerates. The regional lymph nodes become enlarged and sometimes ulcerate (Ghon complex). Histologic examination of the primary ulcer from 10 to 14 days after inoculation reveals a marked inflammatory response with many polymorphonuclear leukocytes and tubercle bacilli. During the next 2 weeks the histologic picture changes. Lymphocytes and epithelioid cells appear and replace the polymorphonuclear leukocytes. Distinct tubercles or tuberculoid structures develop at the site of inoculation, and in the regional lymph nodes as well within 3 to 4 weeks after the inoculation. Simultaneously with the appearance of epithelioid cells the number of tubercle bacilli decreases rapidly (Sulzberger)

Atypical tubercle consists of an accumulation of epithelioid cells surrounded by a wall of lymphocytes. Usually a few Langhans giant cells are present among the epithelioid cells. The epithelioid cell center of the tubercle may show various degrees of caseation. If such typical tubercles are present, one speaks of a tuberculous infiltrate. Frequently, however, in tuberculosis one does not find typical tubercles but only irregular accumulations of epithelioid cells within an inflammatory infiltrate, with or without caseation, and with or without Langhans giant cells. In that case, one speaks of a tuberculoid infiltrate.

It is important to realize that a tuberculous, and particularly a tuberculoid infiltrate does not necessarily mean tuberculosis. Either may occur in many other diseases, particularly syphilis, leprosy and several of the deep fungus infections. The Jadassohn-Lewandowsky law states that, wherever microorganisms or their products are being overcome or neutralized by the local immunobiologic reactions, tubercles or tuberculoid structures have a tendency to appear.

Of interest is the work of Sabin, who found, by intracutaneous injections of various fractions of tubercle bacilli, that the protein fraction evokes the necrotizing and lymphocytic response and the phospholipid fraction evokes the epithelioid cell response.

The histologic diagnosis of the various types of cutaneous tuberculosis is dependent on the correlation of the degree of caseation, necrosis, the amount of inflammatory infiltrate, the degree of vascular changes and the relationship of the tubercles to blood and lymph vessels.

Caseation necrosis is so called because of the cheesy macroscopic appearance of the affected tissue. Microscopically, areas of caseation necrosis show complete loss of their architectural outline. One observes eosinophilic granular material in which, unless the caseation necrosis is far advanced, some nuclei are still present. However, most of the nuclei show pyknosis (shrinkage) or karyorrhexis (fragmentation). In local tuberculous infections, caseation necrosis is caused by the action of bacterial toxins alone. In hematogenous infections, obliterative changes in the vessels are responsible in part. If any marked inflammatory infiltrate is absent in lesions of tuberculosis, it is indicative of either a relative resistance of the host or of an attenuated infection.

No generally accepted classification of tuberculosis of the skin exists. The classification presented in Table 3 is based on the classifications offered by Montgomery and by Layman and Michelson.

TABLE 3—CLASSIFICATION OF TUBERCULOSIS OF THE SKIN

	AMOUNT OF CASEATION
1 <i>Primary tuberculosis</i>	
a Localized infection tuberculous chancre	Considerable
b Hematogenous disseminate infection generalized miliary tuberculosis of the skin	Considerable
2 <i>Reinfection tuberculosis</i>	
a Localized infection	
(1) Lupus vulgaris	Slight
(2) Tuberculosis verrucosa cutis	Moderate
(3) Scrofuloderma	Considerable
(4) Tuberculosis cutis orificialis	Considerable
b Hematogenous disseminate infection tuberculids	
(1) Micropapular tuberculid either purely papular or rosea like (Lewandowsky)	None
(2) Lupus miliaris disseminatus faciei	Slight
(3) Papulonecrotic tuberculid	Considerable
(4) Iichen scrofulosorum	None
(5) Erythema induratum	Considerable

## 1 PRIMARY TUBERCULOSIS

Primary infection with tuberculosis usually takes place in childhood. It occurs only very rarely on the skin. In the vast majority of cases, it presents itself in the lung as the so called Ghon complex or primary complex. This consists of a small, caseous lesion at the periphery of one lung with caseation of the regional hilar lymph nodes. The Ghon complex does not become chronic. It either heals or extends rapidly. Extension may be by continuity or by hematogenous dissemination. In the latter case, miliary tuberculosis may develop.

## A TUBERCULOUS CHANCRE

Primary infection of the skin with tuberculosis is more apt to occur in children but may be seen in adults (Michelson, 1935). The cutaneous lesion usually consists of a crust covered ulcer. It is referred to as tuberculous chancre or primary inoculation tuberculosis of the skin. The regional lymph nodes are enlarged and may or may not suppurate and produce draining sinuses.

**Histopathology.** The histologic development of the lesion is very much like that observed in experimental cutaneous inoculation of the guinea pig with tubercle bacilli (see page 174). During the early

and at a later stage the histologic picture is that of a banal acute,

necrosis. After from 3 to 6 weeks, a more specific histologic picture develops. Epithelioid and Langhans giant cells are then present, though typical tubercles do not form, as a rule. Caseation necrosis remains a prominent feature. At this stage, the number of tubercle bacilli is so greatly decreased that it may be impossible to demonstrate them in histologic sections and the only proof of their presence is through positive animal inoculation experiments. Simultaneously with the decrease in the number of tubercle bacilli in the lesion, the tuberculin test, previously negative, becomes positive. The histologic picture in the regional lymph nodes is identical with that of the cutaneous lesion (O Leary and Harrison).

#### B GENERALIZED VILIARY TUBERCULOSIS OF THE SKIN

The cutaneous lesions are usually papules but may be vesicles or pustules. They tend to break down and form small ulcers.

**Histopathology.** In early lesions, one observes a nonspecific inflammatory infiltrate with foci of necrosis but without tuberculoid reaction. Numerous tubercle bacilli are found within the blood vessels and in the foci of necrosis. At a later stage, tuberculoid formations may be encountered (Wise).

### 2 REINFECTION TUBERCULOSIS

The immunity acquired by the primary infection almost always protects at least for several years. After such a latent period reinfection may occur. Reinfection tuberculosis usually is mono-organic, so that in cases in which the skin is affected other organs are, as a rule, free from active tuberculosis. As in primary tuberculosis, the infection of the skin may be localized or disseminate.

#### A LOCALIZED INFECTION

In localized infections one may find, depending on the virulence of the bacilli and the degree of resistance of the host, either a slight amount of caseation, as in lupus vulgaris, a moderate amount of caseation as in tuberculosis verrucosa cutis, or a considerable amount of caseation as in scrofuloderma and tuberculosis cutis orificialis.

##### (1) Lupus Vulgaris

In lupus vulgaris the lesions, which are found most commonly on the face consist of sharply demarcated, reddish brown patches containing pinhead sized, deep-seated nodules. If the blood is pressed



out of the skin by pressure with a glass slide (diascopy) the nodules stand out clearly as yellowish brown macules. Because of their yellowish brown color, the nodules are referred to as apple jelly nodules. In the course of time as a rule, the affected areas become atrophic with contraction of the tissue. However some areas may show ver




FIG 79 Lupus vulgaris. Low magnification. There are several tubercles. The large tubercle in the center shows slight caseation necrosis ( $\times 100$ ).

verrucous thickening (lupus vulgaris verrucosus) or superficial ulceration. Squamous cell carcinoma may develop at the margin of the ulcers.

**Histopathology** Typical tubercles with epithelioid cells, giant cells and a peripheral zone of lymphocytes are present. Caseation necrosis within the tubercles is slight and may be absent (Figs 79 and 80). The amount of inflammatory infiltrate composed of lymphocytes and plasma cells varies. In some cases the inflammatory infiltrate dominates the histologic picture so that one has to search for occasional tubercles, in other cases it is slight.

The infiltrate of lupus vulgaris is most pronounced in the upper dermis but in some areas may extend into the subcutaneous layer. It causes destruction of the cutaneous appendages. In areas of healing, extensive fibrosis occurs.

Secondary changes in the epidermis are common. In some areas the epidermis may show acanthosis, hyperkeratosis and even papill




FIG. 80. Lupus vulgaris. High magnification of Figure 79. A tubercle containing several Langhans giant cells is shown. The nuclei of the giant cells lie in horseshoe arrangement. (X400)

omatosis (lupus vulgaris verrucosus). In other areas, by the pressure of the infiltrate, atrophy and even destruction of the epidermis may occur, resulting in ulceration and secondary pyogenic infection. At the margin of such ulcers, pseudo-epitheliomatous hyperplasia and, in some instances, squamous-cell carcinoma may be found.

Tubercle bacilli are present in such small numbers that their presence can hardly ever be demonstrated by staining methods. Guinea pig inoculation on the other hand usually gives a positive result.

**Differential Diagnosis.** Differentiation from sarcoidosis may be very difficult and occasionally is impossible. No absolute histologic criterion exists by which the two diseases can be differentiated with certainty. As a rule, however, the infiltrate in sarcoidosis tends to

lie in scattered islands throughout the dermis while in lupus vulgaris the infiltrate tends to be compact and to lie mainly in the upper dermis. Furthermore, sarcoidosis usually shows much less lymphocytic reaction, more fibrosis around the nests of epithelioid cells, a smaller number of giant cells and complete absence of necrosis (Ronchese). The epidermis in sarcoidosis is either normal or shows pressure atrophy while in lupus vulgaris in addition to atrophy there may be areas of ulceration, acanthosis and pseudoepitheliomatous hyperplasia. However the only laboratory procedure by which the two diseases can be differentiated with certainty is guinea pig inoculation which usually is positive in lupus vulgaris, always negative in sarcoidosis.

For a discussion of swimming pool granuloma which is thought by some to be an inoculation lupus vulgaris see page 142.

## (2) *Tuberculosis verrucosa Cutis (Including Mycobacterium Necrogena)*

This form of tuberculosis represents an infection of a nearly immune skin with virulent tubercle bacilli. The lesions of which there may be one or several consist of verrucous hyperkeratotic areas surrounded by an inflammatory border. Crusts may be intermingled with the keratotic material. Frequently pus may be expressed from fissures within the verrucous lesion.

**Histopathology** The histologic picture shows acanthosis, hyperkeratosis and papillomatosis of the epidermis. Beneath the epidermis there is an acute inflammatory infiltrate of polymorphonuclear leukocytes and lymphocytes with abscess formation. In the middle portion of the dermis one usually finds typical tubercles with a moderate amount of caseation necrosis. At times however only a nonspecific inflammatory infiltrate is present. Tubercle bacilli are more numerous than in lupus vulgaris and therefore occasionally can be demonstrated histologically (Montgomery).

## (3) *Scrofuloderma (Tuberculosis Cutis Colligata)*

Scrofuloderma may originate in the subcutaneous tissue frequently however it represents a direct or lymphatic extension to the skin of an underlying tuberculous focus located usually in a lymph node or bone. The lesion becomes first manifest as a bluish red painless swelling, which suppurates and later breaks down to form an ulcer with irregular undermined bluish borders.

**Histopathology** The center of the lesion shows nonspecific abscess formation with ulceration of the epidermis. At the periphery how

ever one sees tubercle formation with marked caseation necrosis and a considerable amount of chronic inflammatory reaction (Fig 81). The number of tubercle bacilli is usually sufficient to enable one to find them in sections stained after Ziehl Neelsen (Michelson, 1924)




FIG 81 Scrofuloderma Margin of an ulcer On the right side of the photograph one observes necrosis of epidermis and dermis In the center are tuberculoid structures On the left the infiltrate is composed of lymphocytes and plasma cells (X200)

**Differential Diagnosis** For differentiation from erythema induratum see page 186 and from gummatous syphilis, see page 215

#### (4) Tuberculosis Cutis Orificialis

The lesions are shallow ulcers with a granulating base, occurring about the mucous orifices of patients with virulent internal tuberculosis.

**Histopathology** The histologic picture may show merely an ulcer surrounded by a nonspecific inflammatory infiltrate. In most instances however one finds tubercle formations with caseation necrosis deep in the dermis. The epidermis at the margin of the ulcer may show hyperplasia. Tubercle bacilli usually can be demonstrated in the sections even when the histologic appearance is nonspecific

## B HEMATOGENOUS DISSEMINATE INFECTION TUBERCULIDS

Tuberculids are caused by a hematogenous dissemination of tubercle bacilli in patients possessing a rather high immunity to tuberculosis. Because of the high immunity, the bacilli are rapidly destroyed in the skin and therefore are usually not demonstrable in sections or by animal inoculation. Although, naturally the hematogenous dissemination must take place from an internal focus, often this focus cannot be found. Clinically, as well as histologically, transitions between the various forms of tuberculids occur not infrequently.

(1) *Micropapular Tuberculid (Including Rosacea like Tuberculid of Lewandowsky)*

The eruption is limited to the face and consists of numerous small slightly indurated but not elevated papules of about pinhead size.



FIG. 82. Micropapular tuberculid (rosacea like tuberculid of Lewandowsky). In the center there is an island of epithelioid cells with only a slight admixture of lymphocytes. This picture is indistinguishable from that of sarcoid ( $\times 100$ ).

Some of the patients present in addition a diffuse erythema of the face. To these cases the term rosacea like tubercloid of Lewandowsky has been applied.

**Histopathology** Histologically the papules usually show islands of epithelioid cells with only a few lymphocytes and few or no giant cells they thus present a picture indistinguishable from that of sarcoidosis (Fig 82) (Laymon and Michelson). In other cases a moderate admixture of lymphocytes is present so that the appearance resembles lupus vulgaris more than sarcoid (Mackee and Sulzberger). Caseation necrosis is absent as a rule occasionally one finds slight central necrosis in the tubercles (Wile and Grauer).

Since the papular type of acne rosacea may show the same histologic picture as the rosacea like tuberculid of Lewandowsky (Miescher Laymon) (see page 124) the latter diagnosis should never be made on the basis of the histologic findings alone but only when the following additional criteria exist (1) presence clinically of minute lupoid nodules and (2) concomitant evidence of tuberculosis such as pulmonary tuberculosis or a high degree of tuberculin sensitivity (Laymon) The possibility exists that all cases of rosacea like tuberculid are instances of acne rosacea and that cases of micro papular tuberculid without rosacea like features represent military sarcoidosis

(2) *Lupus Miliaris Disseminatus Faciei*

Lupus miliaris disseminatus faciei occurs on the face only. The eruption consists of firm elevated discrete papules occurring singly or in groups.

less pronounced. *capus vulgaris* although in the latter caseation is

**ICHENOID TUBERCULID** A variant of lupus miliaris disseminatus faciei with the same histologic picture has been described by Okulu and Montgomery under the name Ichenoid tuberculid. There is a generalized eruption of discrete or grouped erythematous papules with predominate on the extremities and usually no lesions on the face.

(†) *Papulonecrotic Tuberculosis*

The eruption is not limited to the face as it is in the two preceding types. In addition to the face the extremities and the trunk may be affected. The lesions consist of indolent inflammatory papules which

come in crops and undergo central necrosis. Papulonecrotic tubercle occasionally occurs simultaneously with erythema induratum.

**Histopathology** One observes a small central area of necrosis involving the superficial dermis and the overlying epidermis. The area



FIG. 83. Lupus miliaris disseminatus faciei. A tubercle showing central necrosis is present in the dermis. It is surrounded by a moderately severe inflammatory infiltrate ( $\times 100$ ).

of necrosis is surrounded by a zone of inflammation which is largely nonspecific but may contain at its periphery tubercles showing a moderate amount of caseation necrosis. The infiltrate may extend into the subcutaneous fat. The blood vessels in the lower dermis show obliterative endarteritis and endophlebitis with thrombosis and canalization of their lumen. The walls of the vessels may be invaded by inflammatory cells. The vascular lesions are responsible for the

superficial necrosis. The histologic picture may be called a miniature erythema induratum.

#### (4) *Lichen Scrofulosorum*

The lesions which occur chiefly on the trunk consist of indolent pinhead sized papules. Their color varies from that of normal skin to a pale red.

**Histopathology** The infiltrate consists of almost pure epithelioid tubercles with only an occasional giant cell. A narrow zone of lymphocytes may or may not be present at the periphery of the tubercles. There is no caseation. The infiltrate is located in the upper dermis and frequently but not invariably is arranged about hair follicles (Montgomery).

#### (5) *Erythema Induratum*

Erythema induratum or tuberculosis cutis indurativa is a chronic recurrent eruption occurring on the calves of women. In contrast with the other tuberculids the presence of tubercle bacilli has been demonstrated repeatedly in histologic sections and by inoculation of guinea pigs (Montgomery, O Leary and Barker). The lesions consist at first of painless deep-seated subcutaneous infiltrations. Gradually the infiltrations extend to the surface forming bluish red

1.

• but, tuberculoid at least in some areas. Sometimes however it is necessary to cut deeper into the block of tissue to find such areas.

In areas in which the infiltrate is tuberculoid one finds epithelioid and giant cells occasionally in tubercle arrangement (Fig. 85). In areas of nonspecific infiltration one observes predominantly lymphocytes and plasma cells. Both types of infiltrates invade between the fat cells and gradually replace them (proliferation atrophy or Wucheratrophic of fat). Caseation necrosis is nearly always present and may be extensive. In areas of caseation necrosis the fat cells often are still preserved while the invading infiltrate between them has been supplanted by an amorphous finely granular eosinophilic material in which some pyknotic nuclei are present.

Vascular changes are extensive and usually severe. All sizes of vessels show proliferative changes. The changes are severest in the larger arteries and veins. Their walls become infiltrated with round cells and greatly thickened in all coats. Thrombosis and obliteration result



(Fig 86) The obliteration results in widespread necrosis and abscess formation. The necrosis may extend to the dermis and the epidermis and lead to ulceration.

**Differential Diagnosis** Differentiation from erythema nodosum rarely causes difficulties even though a few small foci of tuberculoid



FIG 84 Erythema induratum. Low magnification. The infiltrate is limited to the subcutaneous fat. It invades between the fat cells and gradually replaces them (proliferation atrophy of the fat) ( $\times 25$ ).

infiltrate may occur in erythema nodosum. In the first place the infiltrate is much more massive in erythema induratum than in erythema nodosum, where it usually consists of small scattered aggregates and most important one usually finds at least a few areas of caseation/necrosis which never occurs in erythema nodosum. Furthermore, extensive tuberculoid infiltration and abscess formation do not occur in erythema nodosum either and if present establish the diagnosis of erythema induratum beyond any doubt. Lesions showing a pronounced tuberculoid infiltrate, extensive caseation and ulcers



in the subcutis

proliferation of their walls (X=500)




FIG. 86 Erythema induratum. High magnification. A large vessel (V) in the subcutaneous fat is invaded by inflammatory cells and thrombosed. At the right of the vessel the inflammatory infiltrate is nonspecific; at the left epithelioid cells and a Langhans type of giant cell (G.C.) are located. (X200)

(Fig 86) The obliteration results in widespread necrosis and abscess formation. The necrosis may extend to the dermis and the epidermis and lead to ulceration.

**Differential Diagnosis** Differentiation from erythema nodosum rarely causes difficulties even though a few small foci of tuberculoid



FIG 81 Erythema induratum. Low magnification. The infiltrate is limited to the subcutaneous fat. It invades between the fat cells and gradually replaces them (proliferation atrophy of the fat). ( $\times 25$ )

infiltrate may occur in erythema nodosum. In the first place, the infiltrate is much more massive in erythema induratum than in erythema nodosum, where it usually consists of small scattered aggregates, and, most important, one usually finds at least a few areas of caseation/necrosis which never occurs in erythema nodosum. Furthermore, extensive tuberculoid infiltration and abscess formation do not occur in erythema nodosum either and if present, establish the diagnosis of erythema induratum beyond any doubt. Lesions showing a pronounced tuberculoid infiltrate, extensive caseation and ulcers

the center of the epithelioid cell islands. If a reticulum stain is employed, one sees a reticulum network surrounding and permeating the islands of epithelioid cells (Fig. 89). The network varies in density. In most areas, the reticulum fibers surround almost every epithelioid cell, but, in others, the fibers are concentrated at the periphery, leaving the central portions relatively free.



Fig. 89. Sarcoidosis active stage. Low magnification. There are large islands of epithelioid cells with only a slight admixture of lymphocytes (X50).

In the healing phase, one finds a gradual transformation of reticulum fibers into collagen. At the periphery of the islands of epithelioid cells and the lymphocytes, the reticulum fibers can be seen to be present in a regular pattern. In such lesions, the reticulum fibers can be seen to be present in a regular pattern. In such lesions, the reticulum fibers can be seen to be present in a regular pattern.

tion may resemble scrofuloderma. However scrofuloderma shows no significant vascular changes and usually shows many tubercle bacilli on staining with Ziehl Neelsen's stain.

For differentiation from gummatous syphilis see page 215

## SARCOIDOSIS

Sarcoidosis is a systemic disease which may affect many organs. The skin frequently is involved. Four types of cutaneous sarcoidosis are generally recognized: Boeck's sarcoid, subcutaneous sarcoid of Darier Roussy, lupus pernio of Besnier and erythrodermic sarcoid. Boeck's sarcoid, the most common type of cutaneous sarcoidosis, is characterized by soft brownish red intracutaneous papules, nodules or plaques. By central clearing annular lesions may result. The lesions only rarely ulcerate. Subcutaneous sarcoid of Darier Roussy shows subcutaneous nodules which often reach several centimeters in diameter and do not break down. The overlying skin is either of normal color or bluish red. Lupus pernio of Besnier presents soft nodules and infiltrated plaques of violaceous color, the surface of which shows numerous telangiectases. Erythrodermic sarcoid is characterized by large sharply demarcated slightly scaling brownish red patches showing little or no infiltration (Lever and Freiman, Wigley and Musso). It might be pointed out that Spiegler Fendt sarcoid is not sarcoidosis but a lymphoid hyperplasia possibly related to lymphoma (see page 491).

**Histopathology.** The histologic picture is essentially the same in all four types of cutaneous sarcoidosis. In the skin, as in other organs, it is characterized by the presence of circumscribed islands of epithelioid cells, so-called epithelioid cell tubercles.

In Boeck's sarcoid the islands of epithelioid cells are located predominantly in the dermis while in Darier Roussy's sarcoid they are found mainly in the subcutaneous tissue. In lupus pernio the histologic picture differs from that of Boeck's sarcoid only by the presence of greatly dilated capillaries in the upper portion of the dermis. In the erythrodermic form the infiltrate shows rather small foci of epithelioid cells intermingled with histiocytes and lymphocytes in superficial perivascular arrangement (Lever and Freiman, Wigley and Musso).

In active lesions of sarcoidosis the islands of epithelioid cells contain few Langhans giant cells or none at all (Fig. 87). A slight and is present particularly in the center (Fig. 88). Necrosis is almost always absent. Rarely some necrosis is found in

Schaumann, 1936) Numerous organs may be involved (Thomas, Ricker and Clark, Longcope and Freiman) The lymph nodes are frequently, and the tonsils occasionally, involved In the lungs, foci of sarcoidosis are often present and may, on roentgenologic examination, reveal a picture resembling that of miliary tuberculosis or Hodgkin's disease (McCort, Wood, Hamilton and Ehrlich). The



FIG 89 Sarcoidosis active stage Foot's reticulum stain Reticulum fibers surround and permeate the epithelioid-cell islands At the margin of the lesions one can observe the gradual transformation of reticulum fibers into collagen fibers ( $\times 100$ )

spleen may be considerably enlarged The phalanges of the fingers and the toes may show lesions of sarcoidosis, which in roentgenograms look like cysts and formerly were called *ostitis cystica* of Jungling Histologically, these lesions represent replacement of bone marrow and bone by sarcoid tissue (Ellis) Oveoparotitis of Heerfordt is also a manifestation of sarcoidosis (Pinner, Michelson) In occasional instances, the heart, the kidneys, the gastro-intestinal tract and the central nervous system are the site of lesions (Longcope and Freiman) Involvement of the lungs, if extensive, of the heart or of the central nervous system may cause death

Cause The cause of sarcoidosis is not yet known Some authors regard it as a noncaseating form of tuberculosis (Pinner). A few

nated and often calcified (Schaumann 1941) In other cases asteroid inclusion bodies have been found inside of giant cells (Fig 91) (Lever and Freeman) The significance of the Schaumann and the asteroid inclusion bodies is not known Neither is ~~is~~ specific for sarcoidosis since Schaumann bodies may occur also in berylliosis (see page 141) and asteroid bodies have been observed in other granulomas especially in foreign body granulomas



FIG 88 Sarcoidosis active stage High magnification of Figure 87 The island of epithelioid cells is sharply demarcated A slight lymphocytic infiltrate is present at the margin of the island Giant cells are absent ( $\times 200$ )

**Differential Diagnosis** The differentiation of lesions of sarcoidosis from those of lupus vulgaris may be very difficult in cases in which the islands show more than just a slight admixture of lymphocytes (see page 179) A histologic picture identical with that seen in sarcoidosis may occur in the cutaneous granulomas of systemic berylliosis (see page 141) and in tuberculoid leprosy (see page 195) In the latter however, epithelioid cell islands usually are found invading and destroying nerves in the dermis and the subcutaneous tissue

**Systemic Lesions** The systemic nature of sarcoidosis at first not recognized, has been established particularly by Schaumann's work

authors have reported the finding of tubercle bacilli in lesions of sarcoidosis (Kyrle Kalkoff and Mohr) It is possible however that in such cases either the diagnosis of sarcoidosis was erroneous or a concomitant tuberculosis existed

If sarcoidosis is due to tuberculosis it is necessary to assume that the tubercle bacilli are quickly killed and disintegrated under the influence of local immune forces Under these circumstances the lipids of the bacillus might be capable of inducing the epithelioid cell response whereas the protein of the bacillus in the absence of hypersensitivity produces no significant necrosis or constitutional reaction (see page 175) Thus the same immunologic situation responsible for the prompt killing of the bacilli might explain the morphologic character of the lesion (Freiman)

## LEPROSY

Leprosy is caused by the lepra bacillus *Mycobacterium leprae* an acid fast organism

Three types of leprosy exist lepromatous leprosy tuberculoid (or neural) leprosy and indeterminate leprosy (Canizares Arnold 1949) The lepromatous and the tuberculoid types are definite clinical pathologic bacteriologic and immunologic forms of leprosy They do not occur together in the same patient and transformation of the lepromatous type into the tuberculoid type or vice versa is very rare Indeterminate leprosy represents a transitional stage It may remain indeterminate for a long time and then heal or may develop into lepromatous or tuberculoid leprosy Lepromatous leprosy has a poor prognosis for arrest or recovery while tuberculoid leprosy has a good prognosis for arrest or recovery In lepromatous leprosy the lepromin test is negative because of absence of immunity in tuberculoid leprosy it is positive In indeterminate leprosy the lepromin test may be positive or negative Cases of indeterminate leprosy showing a positive test will probably remain indeterminate or will develop tuberculoid lesions those showing a negative lepromin test will probably

develop lesions of the skin called lepromas These granulomas occur not only in the skin but also in the mucous membranes of the upper respiratory tract the eyes the testes the superficially located nerves and the reticulo-endothelial structures such as the lymph nodes the liver the spleen and the bone marrow Neurologic changes such as anesthesia trophic disturbances and paralysis usually are present Tuberculoid leprosy also called neural or trculo anesthetic leprosy affects mainly the skin and the nerves






FIG 90 Sarcoidosis healing stage There is considerable fibrosis with progressive obliteration of the epithelioid cells. In contrast with early lesions giant cells are numerous ( $\times 200$ )



FIG 91 Sarcoidosis A large giant cell contains an asteroid inclusion body ( $\times 400$ )

authors have reported the finding of tubercle bacilli in lesions of sarcoidosis (Kyrle Kalkoff and Mohr) It is possible however that in such cases either the diagnosis of sarcoidosis was erroneous or a concomitant tuberculosis existed

If sarcoidosis is due to tuberculosis it is necessary to assume that the tubercle bacilli are quickly killed and disintegrated under the influence of local immune forces Under these circumstances the lipids of the bacillus might be capable of inducing the epithelioid cell response whereas the protein of the bacillus in the absence of hypersensitivity produces no significant necrosis or constitutional reaction (see page 175) Thus the same immunologic situation responsible for the prompt killing of the bacilli might explain the morphologic character of the lesion (Freiman)

## LEPROSY

Leprosy is caused by the lepra bacillus *Mycobacterium leprae* an acid fast organism

Three types of leprosy exist lepromatous leprosy tuberculoid (or neural) leprosy and indeterminate leprosy (Canizares Arnold 1949) The lepromatous and the tuberculoid types are definite clinical pathologic bacteriologic and immunologic forms of leprosy They do not occur together in the same patient and transformation of the lepromatous type into the tuberculoid type or vice versa is very rare Indeterminate leprosy represents a transitional stage It may remain indeterminate for a long time and then heal or may develop into lepromatous or tuberculoid leprosy Lepromatous leprosy has a poor prognosis for arrest or recovery while tuberculoid leprosy has a good prognosis for arrest or recovery In lepromatous leprosy the lepromin test is negative because of absence of immunity in tuberculoid leprosy it is positive In indeterminate leprosy the lepromin test may be positive or negative Cases of indeterminate leprosy showing a positive test will probably remain indeterminate or will develop tuberculoid lesions those showing a negative lepromin test will all progress to lepromatous leprosy

These granulomas occur not only in the skin but also in the mucous membranes of the upper respiratory tract the eyes the testes the superficially located nerves and the reticulo endothelial structures such as the lymph nodes the liver the spleen and the bone marrow Neurologic changes such as anesthesia trophic disturbances and paralysis usually are present Tuberculoid leprosy also called neural or maculo-mesthetic leprosy affects mainly the skin and the nerves

These granulomas occur not only in the skin but also in the mucous membranes of the upper respiratory tract the eyes the testes the superficially located nerves and the reticulo endothelial structures such as the lymph nodes the liver the spleen and the bone marrow Neurologic changes such as anesthesia trophic disturbances and paralysis usually are present Tuberculoid leprosy also called neural or maculo-mesthetic leprosy affects mainly the skin and the nerves

The cutaneous lesions consist of sharply circumscribed, erythematous, often hypopigmented patches or flat infiltrations. Annular configuration is common. The lesions usually are hypoesthetic. The peripheral nerves, especially the ulnar nerve are thickened and palpable. Anesthesia, trophic disturbances and paralysis occur just as




FIG. 92 Lepromatous leprosy, early stage. The granulomatous infiltrate consists predominantly of histiocytes and lepra cells with foamy cytoplasm ( $\times 200$ )

in the lepromatous type. Indeterminate leprosy resembles tubercloid leprosy in its clinical manifestations.

**Histopathology** Lepromatous leprosy shows a granulomatous infiltrate. It is massive in the upper dermis and focal around the arteries, the veins and the nerves in the lower dermis and the subcutaneous fat. Histiocytes and lepra cells predominate, but, in addition there are lymphocytes, plasma cells and in older lesions, fibroblasts (Fig. 92). Lepra cells, or Virchow cells develop from histiocytes. They are large, foamy cells resembling xanthoma cells. On staining with fat stains, they are shown to contain lipid which, in contrast to

that found in xanthoma is not doubly refractile (Tilden). On stain

stressed that *Mycobacterium leprae*, unlike *Mycobacterium tuberculosis* is not strongly acid fast. Therefore sections must be decolorized




FIG. 93 Lepromatous leprosy, early stage Ziehl-Neelsen stain. Numerous lepra bacilli are present. In the left center two bundles of bacilli are shown ( $\times 800$ ).

lepra bacilli are found in empty spaces so that they look like large lymphatics (Cowdry) (Fig. 94). On staining after Ziehl-Neelsen or Fite they may or may not contain large globi of partially degenerated lepra bacilli.

Tuberculoid leprosy shows a tuberculoid infiltrate. Patients with tuberculoid leprosy have fairly good immunity against the lepra bacillus. This explains in accordance with the Jadassohn-Lewandowsky law (see page 175) the scarcity or the absence of bacilli and the tuberculoid tissue response. Sections may show almost pure epitheli-

oid-cell tubercles so that differentiation from sarcoidosis may be difficult (Fig. 95). However, a thorough search usually will reveal invasion and destruction of nerves in the dermis or the subcutaneous tissue by epithelioid-cell tubercles, which never occurs in sarcoidosis. In some cases, the tubercles show a moderate admixture of lymphocytes and contain giant cells. However, caseation necrosis does not occur in the skin. Lepra bacilli may be absent in the lesions of tu




FIG. 91 Lepromatous leprosy, late stage. High magnification. There is some fibrosis. The lepra cells are very large in size and appear as round empty spaces ( $\times 200$ )

berculoid leprosy, but not infrequently they are found in small numbers after prolonged search (Arnold, 1915)

Indeterminate leprosy shows the vessels and the nerves of with difficulty (Canizares)

**NEURAL LESIONS** Lesions in the large peripheral nerves occur in almost every case of leprosy, regardless of type. In all three types, the histologic appearance of the neural lesions corresponds to that of the cutaneous lesions (Pardo Castello, Tiant and Piñeyro). The nerve lesions of lepromatous leprosy show large vacuolated Virchow cells and numerous lepra bacilli. In tuberculoid leprosy, one observes in the nerves an extensive infiltrate of epithelioid cell tubercles containing few or no bacilli. However, in contrast with the cutaneous lesions, caseation necrosis is common. In occasional instances, mas

sive caseation of the tuberculoid lesions with complete destruction of nerve tissue occurs ("colligative neuritis").



FIG 95 Tuberculoid leprosy. Low magnification. The infiltrate consists of epithelioid cell tubercles showing a slight admixture of lymphocytes particularly at their margins. That the histologic picture is very much like that of sarcoidosis ( $\times 100$ ).

### DIPHtheria OF THE SKIN

The clinical picture of diphtheria of the skin is varied. Most commonly encountered are one or several punched-out ulcers covered with a pseudomembrane the removal of which is difficult and followed by profuse bleeding. Next in frequency are moist, crusted eczematous lesions which occur particularly in the retro auricular region. Other types of lesions occasionally observed include impetiginous, vesicular and bullous lesions (Reiss).

**Histopathology.** The histologic picture of all forms of diphtheria is nonspecific. The ulcerative lesions show destruction of the epidermis and of the upper dermis within the region of the ulcer. At the margin of the ulcer, the epidermis is acanthotic. The dermis beneath the ulcer shows evidence of acute inflammation. The floor of the ulcer is covered with a layer composed of necrotic cells, fibrin and numerous neutrophils. In some instances, numerous Klebs-Loeffer bacilli can be seen in this necrotic layer (Allen). These bacilli have bipolar bodies and tend to lie in palisade like arrangement. The finding of such bacilli is suggestive of diphtheria. However, since diphtheria bacilli cannot safely be differentiated from pseudo diphtheria bacilli in stained sections, smears and cultures are necessary for the establishment of the diagnosis.

The eczematous lesions may show the histologic picture of either an acute, a subacute or a chronic dermatitis. The histologic changes thus are identical with those observed in an ordinary bacterial dermatitis (Robert).

### ANTHRAX

Anthrax, caused by *Bacillus anthracis*, is characterized by a carbuncle like lesion located usually on an exposed portion of the skin. The lesion starts as a papule, enlarges, ulcerates and then becomes covered with a black eschar. Frequently, a ring of vesicles is present at the margin of the eschar. Marked erythema and edema surround the lesion. Suppuration does not occur. Pain is characteristically slight or absent. Often there is regional lymphadenopathy, which may be painful.

**Histopathology.** At the site of the ulcer, the epidermis is destroyed and the ulcerated surface is covered with necrotic tissue. The adjoining epidermis shows spongiosis and occasionally intraepidermal vesicle formation and is invaded by neutrophils. There is marked edema of the dermis separating the bundles of collagen and loosening their fibrils. Numerous erythrocytes and neutrophils are present throughout the dermis and the subcutaneous tissue. However, abscess formation is absent. Only few histiocytes can be observed. The blood vessels are dilated and their walls show diffuse degenerative changes.

Anthrax bacilli are present in large numbers and can be recognized in sections stained with routine stains but are visualized best in sections stained with Gram's stain. The anthrax bacillus is a large, rod shaped, square ended Gram positive bacillus, from 6 to 10 microns long and from 1 to 2 microns thick. In the tissue, it is usually surrounded by a well defined capsule. Anthrax bacilli are found in countless numbers in the necrotic tissue at the surface of the ulcer. The dermis also contains numerous bacilli while the subcutaneous

tissue usually contains only a few. It is worth noting that phagocytosis of the bacilli by either neutrophils or histiocytes is absent (Lebovitch, McKillip and Conboy).

## TULAREMIA

Tularemia, an infectious disease caused by *Bacterium tularense*, occurs in four types: the ulceroglandular, the oculoglandular, the glandular and the typhoid types. Specific cutaneous lesions occur only in the ulceroglandular type. They are of two varieties: ulcers and nodes. One or several ulcers occur as primary lesions at the site of infection; they tend to have a punched-out appearance. Cutaneous or subcutaneous nodes form in the small lymph nodes located along the lymph vessels draining the primary lesion; they are hard and tender at first but may proceed to suppuration.

**Histopathology.** The primary ulcer shows a nonspecific granulation tissue into which are embedded granulomas composed of three zones: (1) a central zone of necrosis consisting of finely granular eosinophilic material, nuclear fragments and a few erythrocytes; (2) an intermediate zone composed of epithelioid cells with a few Langerhans giant cells and lymphocytes; and (3) an outer zone consisting largely of lymphocytes but containing also some histiocytes, plasma cells and extravasated erythrocytes. Vascular changes in the outer zone and in the surrounding granulation tissue are conspicuous; they consist of proliferation of the endothelial cells and infiltration of the vascular walls by inflammatory cells (Schuermann and Reich). Older lesions may show epithelioid-cell tubercles with only a slight inflammatory reaction and thus may have a sarcoid-like appearance. *Bacterium tularense* although present does not stain in the sections.

The cutaneous lymphatic nodes show like the primary ulcer of tularemia scattered granulomas in which three zones can be recognized.

**Differential Diagnosis.** The histologic picture differs from that of tuberculosis by the distinct arrangement in three zones which is rarely seen so clearly in tuberculosis; by the presence of vascular changes; and by the presence of erythrocytes in the central necrotic zone and the outer lymphocytic zone (Reich). Differentiation from sporotrichosis (see page 230) and lymphogranuloma venereum (see page 252) however may be impossible.

## CHANCROID

Chancroid caused by the *streptobacillus* of Ducrey (*Hemophilus ducreyi*) is a venereal disease causing one or several ulcers chiefly



in the genital region. The ulcers show little if any induration and tend to have an undermined border. Regional lymphadenitis is common and usually terminates in abscess formation (bubo).

**Histopathology.** The chancroidal ulcer presents a granulomatous reaction which is sufficiently distinct to permit a diagnosis of chancroid in many instances (Heyman, Beeson and Sheldon). The lesion consists of three zones (Sheldon and Heyman) and shows characteristic vascular changes (Pond, Greenblatt and Huie). The surface zone or base of the ulcer is rather narrow and is made up of polymorphonuclear leukocytes, fibrin, red blood cells and necrotic tissue. Below this is a fairly wide zone of edematous tissue composed mainly of endothelial cells in varying stages of proliferation. Newly formed blood vessels are numerous. There is marked endothelial proliferation within the vessels, frequently blocking their lumen and leading to thrombosis. In addition, there is degeneration of the walls of vessels. The third zone is composed of a dense infiltration of plasma cells and lymphocytes.

Demonstration of *Ducrey* bacilli in the tissue by staining with Giemsa's stain, Gram's stain or with polychrome methylene blue is only rarely possible. They are most apt to be found as Gram negative coccobacilli between the cells of the surface zone (Sheldon and Heyman).

### GRANULOMA INGUINALE

Granuloma inguinale is a venereal disease caused by *Donovania granulomatis*, which is found in the lesions in the form of intracytoplasmic inclusion bodies called Donovan bodies. The taxonomic position of *Donovania granulomatis* is as yet undecided but it is most likely a bacterium and not a virus. It grows well on the chorioallantoic membrane of chick embryos.

The lesions occur predominantly in the genital region and consist of sharply demarcated granulomatous areas which have a hard base and bleed easily. The border is elevated and often shows a serpiginous outline. The lesions spread very slowly by peripheral extension and may attain large size. After years the lesions heal with a thick, fibrous, contracted scar. In occasional instances squamous cell carcinoma supervenes (Beerman and Sonck, Alexander and Shields).

**Histopathology.** The epidermis may be thinned but more often shows acanthosis, which may reach the proportions of pseudoepitheliomatous hyperplasia (Beerman and Sonck). A granulomatous infiltrate is present in the dermis composed predominantly of histiocytes and plasma cells. Scattered throughout the otherwise mononuclear infiltrate one finds small abscesses composed of poly-

morphonuclear leukocytes (Allen) The number of lymphocytes is conspicuously small

Intracytoplasmic inclusion bodies called Donovan bodies are present within a variable number of histiocytes The parasitized histiocytes or macrophages possess abundant cytoplasm and may measure 20 microns and more in diameter Their cytoplasm has a cystic appearance Within the cystic compartments of the cytoplasm




Fig. 96. Granuloma inguinale. The granulation tissue is composed primarily of histiocytes and plasma cells. There is a diffuse sprinkling of polymorphonuclear leukocytes. Several of the histiocytes contain Donovan bodies within their cytoplasm. ( $\times 400$ )

one sees groups of small round or oval encapsulated bodies measuring 1 to 2 microns in diameter (Fig. 96). In the cross sections of large histiocytes several dozen of such bodies may be observed. The Donovan bodies are recognizable in stains with hematoxylin and eosin but are seen best in sections stained with Giemsa's stain. In such sections they appear bright red (Alexander and Schoch). When a silver stain is used the Donovan bodies appear black and have the shape of a closed safety pin because of their elongated ovoid shape and intense bipolar staining reaction (Torpin, Greenblatt and Pund). The capsule surrounding the Donovan bodies does not stain. In addition to their intracellular location a few Donovan bodies

easier to find them in tissue scrapings or tissue imprints stained with Giemsa's or Wright's stain than in fixed tissue sections

**Differential Diagnosis** The parasitism of histiocytes is strikingly similar to that observed in rhinoscleroma, histoplasmosis and leishmaniasis. However, the small size of the Donovan bodies and the presence of small abscesses in the infiltrate usually make a differentiation from these diseases possible (See Table 5, page 237)

A difficult problem may be posed by the marked epidermal proliferation present occasionally in granuloma inguinale (Beermann and Sonck). Several biopsies may be necessary to decide whether it represents merely pseudoepitheliomatous hyperplasia (see page 334) or squamous cell carcinoma which occasionally supervenes in granuloma inguinale.

### RHINOSCLEROMA

Rhinoscleroma is a chronic infectious disease in which the nose, the lip and the upper respiratory tract are infiltrated with hard plaques and masses.

**Histopathology.** The granulomatous infiltrate is strikingly rich in plasma cells and contains two peculiar types of cells, the Mikulicz cell and the Russell bodies. Because of their presence, the histologic picture of rhinoscleroma is diagnostic.

The Mikulicz cell is a very large, round histiocyte measuring up to 100, occasionally even 200, microns in diameter. It has a pale reticulated, ill defined cytoplasm and an eccentric nucleus. If a bacterial stain such as Giemsa's stain is used, one finds within the cytoplasm of many Mikulicz cells as well as in their vicinity many bacilli called *Klebsiella rhinoscleromatis* or Frisch bacilli (Fig 97). They are short, Gram negative rods measuring 2 by 5 microns in length and are surrounded by a narrow gelatinous capsule (called gloria). It is not certain that the disease is caused by this organism.

The Russell bodies are elliptic formations, measuring from 20 to 40 microns in diameter. Thus, they are smaller than a Mikulicz cell but still twice as large as a normal plasma cell. They have a homogeneous, brilliant red light refractile cytoplasm and no nucleus (Fig 98). They form within plasma cells as a result of their degeneration and finally are expelled (see page 37).

The amount of cellular infiltration varies with the age of the lesion. Early lesions show considerable infiltration by numerous plasma cells and lymphocytes with occasional histiocytes and eosinophils, many Russell bodies are present but only few Mikulicz cells. Gradually, the number of Mikulicz cells increases to such extent that they predominate the histologic picture giving the section a




FIG. 97 Rhinoscleroma. Giemsa stain. There are several Mikulicz cells; the cytoplasm of which is pale, reticulated and ill-defined. One Mikulicz cell contains in its cytoplasm many Frisch bacilli which appear here as small rod-shaped deeply staining bodies ( $\times 900$ ).




FIG. 98 Rhinoscleroma. The infiltrate contains many plasma cells and one Russell body. The Russell body is larger than the other cells in the infiltrate. It has a homogeneous brilliant red refractile cytoplasm ( $\times 400$ ).

easier to find them in tissue scrapings or tissue imprints stained with Giemsa's or Wright's stain than in fixed tissue sections

**Differential Diagnosis** The parasitism of histiocytes is strikingly similar to that observed in rhinoscleroma, histoplasmosis and leishmaniasis. However, the small size of the Donovan bodies and the presence of small abscesses in the infiltrate usually make a differentiation from these diseases possible (See Table 5, page 237)

A difficult problem may be posed by the marked epidermal proliferation present occasionally in granuloma inguinale (Beerman and Sonck). Several biopsies may be necessary to decide whether it represents merely pseudo epitheliomatous hyperplasia (see page 334) or squamous cell carcinoma which occasionally supervenes in granuloma inguinale

### RHINOSCLEROMA

Rhinoscleroma is a chronic infectious disease in which the nose, the lip and the upper respiratory tract are infiltrated with hard plaques and masses

**Histopathology.** The granulomatous infiltrate is strikingly rich in plasma cells and contains two peculiar types of cells, the Mikulicz cell and the Russell bodies. Because of their presence, the histologic picture of rhinoscleroma is diagnostic

The Mikulicz cell is a very large, round histiocyte measuring up to 100, occasionally even 200, microns in diameter. It has a pale, reticulated, ill defined cytoplasm and an eccentric nucleus. If a bacterial stain, such as Giemsa's stain, is used, one finds within the cytoplasm of many Mikulicz cells as well as in their vicinity, many bacilli, called *Klebsiella rhinoscleromatis* or Frisch bacilli (Fig 97). They are short, Gram negative rods, measuring 2 by 5 microns in length and are surrounded by a narrow gelatinous capsule (called gloer). It is not certain that the disease is caused by this organism.

The Russell bodies are elliptic formations, measuring from 20 to 40 microns in diameter. Thus they are smaller than a Mikulicz cell but still twice as large as a normal plasma cell. They have a homogeneous, brilliant red light refractile cytoplasm and no nucleus (Fig 98). They form within plasma cells as a result of their degeneration and finally are expelled (see page 37).

The amount of cellular infiltration varies with the age of the lesion. Early lesions show considerable infiltration by numerous plasma cells and lymphocytes with occasional histiocytes and eosinophils, many Russell bodies are present but only few Mikulicz cells. Gradually, the number of Mikulicz cells increases to such extent that they predominate the histologic picture giving the section a




FIG 97 Rhinoscleroma Giemsa stain There are several Mikulicz cells the cytoplasm of which is pale reticulated and ill-defined One Mikulicz cell contains in its cytoplasm many Frisch bacilli which appear here as small round deeply staining bodies ( $\times 900$ )




FIG 98 Rhinoscleroma The infiltrate contains many plasma cells and one Russell body The Russell body is larger than the other cells in the infiltrate It has a homogeneous brilliant red refractile cytoplasm ( $\times 400$ )

foamy, icellike appearance. In all, fibrosis replaces the Mikulicz cells (Kline and Brody)

**Differential Diagnosis** Parasitized histiocytes are observed also in cutaneous leishmaniasis histoplasmosis and granuloma inguinale. For their differentiation see cutaneous leishmaniasis (page 295 and Table 5 page 297). Russell bodies are not found in these three diseases and therefore are of considerable diagnostic value. However they are not specific for rhinoscleroma because they may occur in other diseases when an infiltrate rich in plasma cells is present—for instance in syphilis tuberculosis squamous cell carcinoma and mycosis fungoides.

### BIBLIOGRAPHY

#### Folliculitis Keloidalis

Adams H. G. Dermatitis papillaris capillitii (Koppe) Brit J Dermat 26 (9) 1911

#### Hidradenitis Suppurativa

Brunsting H. A. Hidradenitis suppurativa abscess of the apocrine sweat glands Arch Dermat & Syph 39 108 1939

#### Pyoderma Gangrenosum

Brunsting H. A. Goettkeman W. H. and O'Leary P. A. Pyoderma (ecthyma) gangrenosum Arch Dermat & Syph 22 605 1930

Dostrovsky A. and Sigher F. A study on ulcer phagedenicum cutis Arch Dermat & Syph 54 408 1946

Iane C. W. and Stroud C. M. Pyoderma gangrenosum Arch Dermat & Syph 27 460 1933

#### Subacute Bacterial Endocarditis

Cornil I. Mosinger M. and Joue A. A. Contribution à l'étude histologique du nodule d'Osler Ann dermat path 13 675 1936

Libman E. and Friedberg C. K. Subacute bacterial endocarditis. In Christian H. A. The Oxford Medicine vol 2 part 2 p 346 New York Oxford 1947

Merklen P. and Wolf M. Participation des endothélites artério-capillaires au syndrome de l'endocardite maligne lente Presse méd 36 97 1928

#### Meningococcemia

Brown C. L. The skin lesions in meningococcus septicemia Am J Dis Child 27 598 1924

Ferguson G. D. Immuring meningococcic infections and

syndrome Am J Path 24 763 1918

Hill J. Lesions in acute meningococcemia

#### Keratosis Blennorrhagica

Carr J. L. and Friedman M. Keratoderma blennorrhagicum Am J Path 20 709 1944

Epstein E. and Chambers S. O. Keratosis blennorrhagica with corneal lesions Arch Dermat & Syph 36 1044 1937

Herold W. C. and Smith D. C. Keratosis blennorrhagica Arch Dermat & Syph 44 398 1941

- Kogoj F. Das klinische und histologische Bild der Acrodermatitis continua Arch f Dermat u Syph 193 117 1951
- Kuske H. Ueber die Hauterscheinungen bei Morbus Reiter Arch f Dermat u Syph 179 58 1939
- Lever W F and Crawford G M. Keratosis blennorrhagica without gonorrhea (Reiter's disease) Arch Dermat & Syph 49 389 1914
- Margolin F S. Gonorrheal dermatitis as part of systemic gonorrhea Urol & Cutan Rev 47 512 1913
- Miale J B and Singletary W V. Cutaneous manifestations of gonococci in infection Arch Dermat & Syph 57 151 1918

## Tuberculosis

- Irgang S. Superficial papulonecrotic tuberculid in the Negro Arch Dermat & Syph 47 627 1913
- Micropapular tuberculid in the Negro Arch Dermat & Syph 53 372 1916
- Kinnear J. Bazin's disease Brit J Dermat 51 18 1939
- Laymon C W. Lupoid rosacea Arch Dermat & Syph 63 409 1951
- and Michelson H E. The micropapular tuberculid Arch Dermat & Syph 42 625 1940 (Good discussion of tuberculids)
- Mackee G M and Sulzberger M B. Rosacealike tuberculid of Lewandowsky Arch Dermat & Syph 51 159 1935
- Michelson H E. Scrofuloderma gummosa (tuberculosis colligativa) Arch Dermat & Syph 10 565 1924
- The primary complex of tuberculosis of the skin Arch Dermat & Syph 32 589 1935
- Miescher C. Rosacea und Rosacea ähnliche Tuberkulide Dermatolog 88 150 1913
- Montgomery H. Histopathology of the various types of cutaneous tuberculosis Arch Dermat & Syph 35 698 1937 (Good review)
- O'Leary P A and Barker N W. Nodular vascular diseases of the leg J A M A 128 335 1945
- Ockuly O F and Montgomery H. Lichenoid tuberculid J Invest Dermat 14 415 1950
- O'Leary P A and Harrison M W. Inoculation tuberculosis Arch Dermat & Syph 44 371 1911
- Rosenberg A
- Sch

Sul

- Wile U J and Crauer F H. Rosacea like tuberculosis Arch Dermat & Syph 51 174 1935
- Wise F. Miliary tuberculosis of the skin. Lichen scrofulosorum and the papulo-tuberculids J Cutan Dis 37 105 1919

## Sarcoidosis

- Ellis F A. Jungling's "ostitis tuberculosa multiplex cystoides" is not cystic tuberculosis ostitis Acta med Scandinav 101 21 1910
- Freeman D C. Sarcoidosis New England J Med 239 661 709 743 1948 (Good review)
- Kalkoff K W and Mohr H J. Zum Erregernachweis der Boeck'schen Krankheit Arch. f. Dermat. u. Syph 188 202 1949



- Kyrle J Die Anfangsstadien des Boeckschen Sarkoids Arch f Dermat u Syph 131 33 1921
- Lever W F and Freiman D G Sarcoidosis Report of a case with erythrodermic lesions subcutaneous nodes and asteroid inclusion bodies in giant cells Arch Dermat & Syph 57 639 1948
- Longcope W T and Freiman D G A study of sarcoidosis Medicine 31 1 1952 (Detailed monograph)
- McCort J J Wood R H Hamilton J B and Ehrlich D E Sarcoidosis Arch Int Med 80 293 1947
- Michelson H E Uveoparonychia Arch Dermat & Syph 39 329 1939
- Pinner M Noncaseating tuberculosis Am Rev Tuberc. 57 690 1938
- Ricker W and Clark M Sarcoidosis Am J Clin Path 19 725 1949
- Schaumann J Lymphogranulomatosis benigna in the light of prolonged clinical observations and autopsy findings Brit J Dermat 48 399 1936
- On the nature of certain peculiar corpuscles present in the tissue of lymphogranulomatosis benigna Acta med Scandinav 106 239 1941
- Thomas C Sarcoidosis Arch Dermat & Syph 47 58 1943
- Wigley J E M and Musso L A A case of sarcoidosis with erythrodermic lesions Brit J Dermat 63 398 1951

### Leprosy

- Allen A C Survey of pathologic studies of cutaneous diseases during World War II Arch Dermat & Syph 57 19 1948
- Arnold H L Differentiation of lepromatous from neural leprosy Arch Dermat & Syph 52 351 1945
- Macules of leprosy Arch Dermat & Syph 60 1148 1949
- Canizares O New official classification of leprosy Arch Dermat & Syph 59 584 1949
- Cowdry E V Cytological studies on globi in leprosy Am J Path 16 103 1940
- Pardo Castello V Tiant F R and Pineyro R Nerve lesions of leprosy Arch Dermat & Syph 55 783 1947
- Tilden I L Lepromatous leprosy a reticulo-endothelial disease Am J Clin Path 15 165 1945 (Good histologic description)

### Diphtheria of the Skin

- Allen A C Survey of pathologic studies of cutaneous diseases during World War II Arch Dermat & Syph 57 19 1948
- Reiss F Cutaneous diphtheria Arch Dermat & Syph 56 216 1947
- Robert P Les lésions histologiques de la diphtherie cutanée superficielle de type eczématoïde Dermatolog 91 334 1947

### Anthrax

- Lebowich R J McPhillip B G and Conboy J R Cutaneous anthrax Am J Clin Path 13 505 1943

### Tularemia

- Hitch J M and Smith D C Cutaneous manifestations of tularemia Arch Dermat & Syph 38 859 1938
- Lawless T K Tularemia Arch Dermat & Syph 44 147 1941 (Good review)
- Reich H Zur Kenntnis der Tularamie hautnaher (regionaler) Lymphknoten Arch f Dermat u Syph 192 175 1950

## Chancroid

- Heyman A Beeson P B and Sheldon W H Diagnosis of chancroid J A M A 129 935 1945  
Fund E R Greenblatt R B and Hue G B The role of biopsy in diagnosis of venereal disease Am J Syph Gon & Ven Dis 22 495 1938  
Sheldon W H and Heyman A Studies on chancroid Am J Path 22 415 1916

### Granuloma Inguinale

- Alexander L. J. and Schoch A. G. Granuloma inguinale *Am J Syph Gon & Ven Dis* 24 180 1940
- and Shields T. L. Squamous cell carcinoma of the vulva secondary to granuloma inguinale *Arch Dermat & Syph* 67 393 1955
- Allen A. C. Survey of pathologic studies of cutaneous diseases during World War II *Arch Dermat & Syph* 57 19 1948
- Beerman H. and Sonck C. E. The epithelial changes in granuloma inguinale *Am J Syph* 36 501 1952
- Margolis G. The diagnosis of granuloma venereum from frozen sections stained with polychrome methylene blue *Am J Path* 21 543 1945
- Pund E. R. and Greenblatt R. B. Specific histology of granuloma inguinale *Arch Path* 23 221 1957
- Irwin R. Greenblatt R. B. and Pund E. R. Granuloma inguinale (venereum) in the female *Am J Surg* 44 551 1959

### Rhinoviruses

- Kline P R and Brody E R Scleroma Arch Dermat & Syph 59 606 1919  
(Good review)  
Simpson L A and Ellis F A Rhinoscleroma Arch Dermat & Syph 39 903  
1939  
Weiss J A Scleroma (rhinoscleroma) histologic changes following teluradium  
therapy Arch Otolaryng 30 38 1939

# 12

## Spirochetal Diseases

### SYPHILIS

Acquired syphilis may be divided into three stages primary, secondary and tertiary. Primary and secondary syphilis comprise the early phase and tertiary syphilis the late phase of the infection. During the early phase, the causative organism, *Treponema pallidum* (*Spirochaeta pallida*), often can be demonstrated in the cutaneous lesions by dark-field examination. In the late phase, no spirochetes can be demonstrated in the cutaneous lesions.

Primary syphilis is characterized by the syphilitic or hard chancre which is usually a single lesion but may be multiple. The typical, or Hunterian, chancre is represented by a brownish red, indurated round papule or plaque with an eroded surface. Occasionally, the chancre shows ulceration. The regional lymph nodes are enlarged.

Secondary syphilis is characterized by a more or less generalized eruption, which is composed usually of macules or papules having a brownish red color. In the anogenital region, the papules may become large, verrucous and moist, they are then called condylomata lata. (They must be differentiated from condylomata acuminata, a variety of verruca, see page 251.) In some instances the cutaneous eruption, due to the presence of scaling, resembles psoriasis (psoriast form syphilid). Occasionally, pustules develop and are followed by heavy crusting (rupial syphilid). Ulceration of lesions in secondary syphilis is very rare and occurs only in severe cases. Alopecia syphilitica is characterized by numerous small patches of partial, "moth eaten" alopecia.

Tertiary syphilis often shows only a single lesion, but occasionally several lesions. A superficial nodular type and a deep, gummatous

center. Lesions of the gummatous type begin as a soft cutaneous or subcutaneous swelling, which breaks down to form one or several ulcers having a punched out appearance. In rare instances, joint articular nodes occur in tertiary syphilis. They are painless, slowly growing, subcutaneous, fibrous nodules, often symmetrically situated

in the vicinity of joints. The elbows and the knees are the sites of predilection.

*Congenital, or prenatal, syphilis* shows the same cutaneous manifestations as acquired secondary and tertiary syphilis with the following exceptions: first occasionally bullous lesions occur especially on the palms and the soles as a manifestation of secondary syphilis in infants and second juxta articular nodes do not occur.

**Histopathology.** The fundamental pathologic changes in syphilis are (1) a predominantly perivascular infiltrate composed of lymphocytes and many plasma cells and (2) endarteritis and endophlebitis. In tertiary syphilis one usually finds in addition a tuberculoid infiltrate and caseation necrosis.

**PRIMARY SYPHILIS.** In the typical primary lesion the epidermis shows at the margin of the lesion acanthosis. Toward the center the epidermis gradually becomes thinner until in the central portion it is entirely absent (Fig 99). An infiltrate composed of lymphocytes and many plasma cells is present in the dermis. It is compact in the center while at the margin it consists of individual perivascular islands. Both blood and lymph vessels show changes. The capillaries and the lymphatics are increased in number and show considerable proliferation of their endothelial cells (Fig 100). The larger blood vessels show proliferation of all their coats and invasion of their walls by the cellular infiltrate. Obliteration and thrombosis of the lumen of some of the vessels is common and results in small foci of necrosis. Occasionally the lumen of so many vessels are occluded that massive necrosis results. In such cases the primary lesion presents itself as an ulcer instead of an erosion.

On staining with Levadits stain spirochetes usually can be found often in large number. They are found throughout the tissue but especially in and around the walls of capillaries and lymphatics.

Histologic examination of the regional lymph nodes in primary syphilis may reveal only a nonspecific inflammation, not infrequently however small foci of tuberculoid reaction are encountered in addition (Michelson).

**SECONDARY SYPHILIS.** In secondary syphilis the number of spirochetes seen in sections stained with Levadits stain varies with the age of lesion. In the irregular lesions of early secondary syphilis spirochetes cannot be found as a rule. In papular lesions they are seen occasionally. In condylomata lata they are also

# 12

## Spirochetal Diseases

### SYPHILIS

Acquired syphilis may be divided into three stages primary, secondary and tertiary. Primary and secondary syphilis comprise the early phase and tertiary syphilis the late phase of the infection. During the early phase, the causative organism, *Treponema pallidum* (*Spirochaeta pallida*), often can be demonstrated in the cutaneous lesions by dark field examination. In the late phase, no spirochetes can be demonstrated in the cutaneous lesions.

*Primary syphilis* is characterized by the syphilitic or hard chancre which is usually a single lesion but may be multiple. The typical, or Hunterian, chancre is represented by a brownish red, indurated round papule or plaque with an eroded surface. Occasionally, the chancre shows ulceration. The regional lymph nodes are enlarged.

*Secondary syphilis* is characterized by a more or less generalized eruption, which is composed usually of macules or papules having a brownish red color. In the anogenital region, the papules may become large, verrucous and moist, they are then called condylomata lata. (They must be differentiated from condylomata acuminata a variety of verruca, see page 251.) In some instances the cutaneous eruption, due to the presence of scaling, resembles psoriasis (psoriatic form syphilid). Occasionally pustules develop and are followed by heavy crusting (rupial syphilid). Ulceration of lesions in secondary syphilis is very rare and occurs only in severe cases. Alopecia syphilitica is characterized by numerous small patches of partial, 'moth eaten' alopecia.

*Tertiary syphilis* often shows only a single lesion, but occasionally several lesions. A superficial nodular type and a deep gummatous

Lesions of the nodular type show an ac-  
posed of nodules and a smooth atrophic  
center. Lesions of the gummatous type begin as a soft cutaneous or subcutaneous swelling, which breaks down to form one or several ulcers having a punched out appearance. In rare instances, juxta articular nodes occur in tertiary syphilis. They are painless, slowly growing, subcutaneous, fibrous nodules, often symmetrically situated

in the vicinity of joints. The elbows and the knees are the sites of predilection.

*Congenital, or prenatal syphilis* shows the same cutaneous manifestations as acquired secondary and tertiary syphilis with the following exceptions: first, occasionally bullous lesions occur, especially on the palms and the soles as a manifestation of secondary syphilis in infants; and second, juxta-articular nodes do not occur.

**Histopathology.** The fundamental pathologic changes in syphilis are (1) a predominantly perivascular infiltrate composed of lymphocytes and many plasma cells, and (2) endarteritis and endophlebitis. In tertiary syphilis one usually finds in addition a tuberculoid infiltrate and caseation necrosis.

**PRIMARY SYPHILIS.** In the typical primary lesion the epidermis shows at the margin of the lesion acanthosis. Toward the center the epidermis gradually becomes thinner until in the central portion it is entirely absent. The dermis contains a dense cellular infiltrate in

lar islands. Both blood and lymph vessels show changes. The capillaries and the lymphatics are increased in number and show considerable proliferation of their endothelial cells (Fig. 100). The larger blood vessels show proliferation of all their coats and invasion of their walls by the cellular infiltrate. Obliteration and thrombosis of the lumina of some of the vessels is common and results in small foci of necrosis. Occasionally the lumina of so many vessels are occluded that massive necrosis results. In such cases the primary lesion presents itself as an ulcer instead of an erosion.

On staining with a Levaditi's stain spirochetes usually can be found often in large number. They are found throughout the tissue but especially in and around the walls of capillaries and lymphatics.

Histologic examination of the regional lymph nodes in primary syphilis may reveal only a nonspecific inflammation, not infrequently accompanied by a marked reaction are encountered.

In secondary syphilis the number of spirochetes seen in sections stained with Levaditi's stain varies with the type of lesion. In the inacular lesions of early secondary syphilis spirochetes cannot be found as a rule. In papular lesions they are seen occasionally. In condylomata but they are not

# 12

## Spirochetal Diseases

### SYPHILIS

Acquired syphilis may be divided into three stages primary, secondary and tertiary. Primary and secondary syphilis comprise the early phase and tertiary syphilis the late phase of the infection. During the early phase, the causative organism, *Treponema pallidum* (*Spirochaeta pallida*), often can be demonstrated in the cutaneous lesions by dark field examination. In the late phase, no spirochetes can be demonstrated in the cutaneous lesions.

*Primary syphilis* is characterized by the syphilitic or hard chancre which is usually a single lesion but may be multiple. The typical or Hunterian, chancre is represented by a brownish red, indurated round papule or plaque with an eroded surface. Occasionally the chancre shows ulceration. The regional lymph nodes are enlarged.

*Secondary syphilis* is characterized by a more or less generalized eruption, which is composed usually of macules or papules having a brownish red color. In the anogenital region the papules may become large, verrucous and moist, they are then called condylomata lata. (They must be differentiated from condylomata acuminata, a variety of verruca, see page 251.) In some instances the cutaneous eruption, due to the presence of scaling, resembles psoriasis (psoriasis form syphilid). Occasionally pustules develop and are followed by heavy crusting (crustal syphilid). *Ulceration of lesions in secondary syphilis* is very rare and occurs only in severe cases. Alopecia syphilitica is characterized by numerous small patches of partial, "moth eaten" alopecia.

*Tertiary syphilis* often shows only a single lesion, but occasionally several lesions. A superficial nodular type and a deep gummatous type of tertiary syphilis occur. Lesions of the nodular type show an active, serpiginous border composed of nodules and a smooth atrophic center. Lesions of the gummatous type begin as a soft cutaneous or subcutaneous swelling which breaks down to form one or several ulcers having a punched out appearance. In rare instances, juxta articular nodes occur in tertiary syphilis. They are painless slowly growing, subcutaneous, fibrous nodules, often symmetrically situated

in the vicinity of joints. The elbows and the knees are the sites of predilection.

*Congenital, or prenatal syphilis* shows the same cutaneous manifestations as acquired secondary and tertiary syphilis with the following exceptions: first, occasionally bullous lesions occur especially on the palms and the soles as a manifestation of secondary syphilis in infants; and second, juxta-articular nodes do not occur.

**Histopathology.** The fundamental pathologic changes in syphilis are (1) a predominantly perivascular infiltrate composed of lymphocytes and many plasma cells, and (2) endarteritis and endophlebitis. In tertiary syphilis one usually finds in addition a tuberculoid infiltrate and caseation necrosis.

**PRIMARY SYPHILIS.** In the typical primary lesion the epidermis shows at the margin of the lesion acanthosis. Toward the center the epidermis gradually becomes thinner until in the central por-

tion. Both blood and lymph vessels show changes. The capillaries and the lymphatics are increased in number and show considerable proliferation of their endothelial cells (Fig. 100). The larger blood vessels show proliferation of all their coats and invasion of their walls by the cellular infiltrate. Obliteration and thrombosis of the lumina of some of the vessels is common and results in small foci of necrosis. Occasionally the lumen of so many vessels are occluded that massive necrosis results. In such cases the primary lesion presents itself as an ulcer instead of an erosion.

of

**SECONDARY SYPHILIS.** In secondary syphilis the number of spirochetes seen in sections stained with Levaditi's stain varies with the type of lesion. In the articular lesions of early secondary syphilis spirochetes cannot be found as a rule. In papular lesions they are seen occasionally. In condylomata lata they are almost always present in sufficient numbers to be found without difficulty. They occur not only in the dermis but also between epidermal cells.

The macule is a  
a diagn






FIG 99 Primary syphilis Low magnification The margin of an erosion is shown The epidermis gradually becomes thinner as it approaches the erosion ( $\times 100$ )




FIG 100 Primary syphilis High magnification of Figure 99 The capillaries are greatly increased in number and show marked proliferation of their endothelial cells Many plasma cells are present in the dense infiltrate ( $\times 400$ )

ing of their endothelium and are surrounded by a slight infiltrate of lymphocytes and plasma cells. However the number of plasma cells is not sufficiently large to be of diagnostic value.

*Papular lesions*, as a rule, have a diagnostic appearance. Not only the superficial but also the deeper vessels of the dermis are involved. They show marked endothelial swelling and are surrounded by a pronounced infiltrate (Fig. 101). Because of its perivascular coat-sleeve like arrangement the infiltrate has a patchy distribution. The patchy pattern stands out very clearly in the lower dermis. In the upper dermis in addition to the patchy infiltrate there may be a significantly high number of plasma cells (Fig. 102). In the differential diagnosis the exudative discoid and lichenoid chronic dermatosis of Sulzberger and Garbe have to be excluded.

The latter in particular may suggest secondary syphilis because of the patchy distribution of its infiltrate and the presence of many plasma cells, but neither chronic dermatitis nor exudative discoid and lichenoid chronic dermatosis shows the marked endothelial swelling of the vessels and the extension of the patchy infiltrate into the lower dermis.

*Condylomata lata* show the same changes in the dermis as the papular lesions. In addition the epidermis shows considerable acanthosis with broadening and elongation of the rete ridges, intra-cellular and intercellular edema of the rete cells and migration of polymorphonuclear leukocytes through the epidermis.

*Psoriasisform secondary syphilis* may show epidermal changes similar to those of psoriasis, but the dermal infiltrate is that of secondary

*subcutaneous pustules*. The diagnosis of syphilis usually can be made because the lower dermis shows the characteristic infiltrate of secondary syphilis.

*Eczematous lesions* are rare in secondary syphilis. They occur when the larger vessels at the border of the dermis and the subcutaneous tissue become completely occluded by syphilitic endarteritis and endophlebitis (Wile, Wieder and Warthin).

*Essential alopecia* is found in many instances, however no infiltrate is found (essential alopecia). Wile and Belote believe that the latter type of alopecia is due to the local action of

on endocrine glands or on the autonomic nervous system. Meningeal syphilis is common with that type.

It is not always possible to assign, on the basis of histologic examination, cutaneous lesions of syphilis to either secondary or tertiary syphilis. Not infrequently, one finds in lesions which clinically are to be classified as late secondary syphilis an admixture of epithelioid and giant cells. On the other hand, lesions of early tertiary syphilis



FIG. 101. Secondary syphilis. Low magnification. The vessels throughout the dermis show endothelial proliferation. The cellular infiltrate is located around the vessels in coat sleeve arrangement ( $\times 100$ ).

may show the histologic infiltrate of secondary syphilis without any epithelioid and giant cell reaction.

**TERTIARY SYPHILIS.** In tertiary syphilis one observes a granulation tissue composed of lymphocytes, plasma cells, histiocytes, fibroblasts and varying numbers of epithelioid and giant cells. Usually, but not always, the number of plasma cells is prominent. The infiltrate is massive in the center but tends to have a perivascular arrangement at the periphery. Vascular changes are prominent, particularly in the larger vessels. They show proliferation of their walls leading to narrowing of their lumina and invasion of their walls by the inflammatory infiltrate. Caseation necrosis is frequent. It is probable that caseation necrosis is not due to the vascular changes because often the vascular changes are just as severe in areas without caseation as in areas with caseation. It seems more likely that caseation represents

in allergic phenomenon caused by alteration in the reactivity of the tissue (Boyd). In the healing stage, lesions of tertiary syphilis show numerous fibroblasts. The process ends in fibrosis. Tertiary syphilis occurs in the skin in two forms: nodular and gummatous.

In nodular tertiary syphilis, the granulomatous process is limited to the dermis. The number of epithelioid and giant cells is small,




Figure 101. The perivascular infiltrate contains many plasma cells. (X200)

is a rule. Occasionally, however, these cells are fairly numerous in the center of the lesion. They tend to lie without arrangement into real tubercles, and giant cells of the foreign body type are more common than those of the Langhans type (Fig. 103). Caseation necrosis is usually not extensive and may be absent. If extensive, ulceration results.

In gummatous tertiary syphilis the granulation

... tissue caseation necrosis occurs in the center of the ... giant cells are



FIG 103 Tertiary syphilis nodular type. In the center of the field a large island of epithelioid and giant cells is located. Most giant cells are of the foreign body type rather than of the Langhans type. The infiltrate at the margins contains many plasma cells. ( $\times 100$ )




FIG 104 Tertiary syphilis gumma. On the left side of the field the mass of cells. In the center of the field is seen part of the large mass of the gumma. ( $\times 200$ )

lesion (Fig. 103) The epithelioid and the giant cells are located mainly in the vicinity of the areas of caseation. Because of the deep extension of the process not only the vessels of the dermis as in nodular tertiary syphilis but also the large vessels of the subcutaneous layer are markedly involved.

Differentiation of tertiary syphilis from tuberculosis may be difficult. In the absence of caseation, the nodular type may resemble lupus vulgaris and the gummatous type may suggest scrofuloderma or erythema induratum. The latter in particular, may be difficult to exclude because it shows obliterative vascular changes similar to those of tertiary syphilis. But aside from erythema induratum, vascular changes provide the most important point of differentiation between syphilis and tuberculosis. In addition the prevalence of plasma cells and the presence of giant cells of the foreign body type rather than of the Langhans type favor a diagnosis of syphilis.

In juxta-articular nodes the histologic picture varies with the age of the lesion. Early lesions are fairly cellular (Tuta and Coombs) and present embedded in a dense fibrous tissue granulomatous areas composed of epithelioid cells, lymphocytes and plasma cells with an occasional Langhans type of giant cell. The blood vessels

which the histologic picture is the same as just described for early lesions, an intermediate zone showing dense fibrous tissue with only few cells, and an inner zone showing hyalinized fibrous bands ramifying to form partitions for cystic spaces. The cystic spaces contain

in early nodes the presence of spirochetes (Allen) Jessner as well as Hu, Liu, Chen and Frazier have succeeded in producing syphilis in rabbits by inoculating them with portions of juxta-articular nodes.

**CONGENITAL SYPHILIS** The histologic changes in the cutaneous lesions of early congenital syphilis are similar to those seen in acquired secondary syphilis. Except for the bullous lesions of the palms and the soles. The latter may show no plasma cells in the dermal infiltrate but numerous polymorphonuclear leukocytes and a few lymphocytes. Obliterative vascular changes are present, however (Fraser).

**SYPHILIS OF INTERNAL ORGANS** It is not appropriate to discuss here in detail the histologic appearance of the lesions of syphilis in the

internal organs. However, it may be pointed out that late syphilis may cause two types of reactions in the internal organs: gumma and diffuse interstitial inflammation. The latter reaction is more common.

The gummas in internal organs show the same histologic changes as those observed in the skin.

Diffuse interstitial syphilitic inflammation manifests itself as accumulations of lymphocytes and plasma cells around the small vessels. Crystallization is usually absent. As a result of the long continued inflammation there is gradual degeneration of the parenchymatous structures and their replacement by fibrous tissue (Boyd). This type of reaction produces, for instance, syphilitic hepatic cirrhosis. Syphilis of the aorta is also of the diffuse type. The process starts around the intima of the aorta. The inflammatory infiltrate then extends along the intima into the media. In the media one observes foci of inflammatory cells, nuclear necrosis and extensive destruction of elastic tissue.

General paresis shows perivascular accumulations of lymphocytes and plasma cells in the meninges and in the cortex of the brain. In addition, there is extensive degeneration of the cortex. In contrast to general paresis, tabes does not show inflammatory changes. The essential lesion is degeneration of the posterior columns of the spinal cord.

## BIBLIOGRAPHY

### Syphilis

- Allen, A. C. Survey of pathologic studies of cutaneous diseases during World War II. *Arch. Dermat. & Syph.* 57:19, 1948.
- Boyd, W. A. Textbook of Pathology, ed. 4, pp. 181, 182, 875, 880. Philadelphia: Lea & Febiger, 1943.
- Fordyce, J. A. The pathology of syphilis. *J. Cutan. Dis.* 33:80<sup>9</sup>, 1915.
- Fraser, J. F. The pathology of congenital syphilis. *Arch. Dermat. & Syph.* 1:491, 1920.
- Freeman, H. L. Juxta-articular nodules. *Arch. Dermat. & Syph.* 43:206, 1941.
- Hu, C. K., Liu, Y., Chen, K. C. and Frazier, C. N. Isolation of virulent *Treponema pallidum* from human aorta 32 hours after death from card ova culture syphilis. *Am. J. Med.* 1:301, 1946.
- Jessner, M. Ueber syphilitische juxta-artikuläre Knotenbildungen. *Arch. f. Dermat. u. Syph.* 152:132, 1906.
- Katz, F. and Newton, B. L. Syphilitic juxta-articular nodules. *Arch. Dermat. & Syph.* 48:626, 1943.
- Michelson, H. C. The occurrence of tuberculoid reactions in the inguinal glands of early syphilis. *Arch. Dermat. & Syph.* 19:66, 1929.
- Tuttle, J. A. and Coombs, R. A. Symmetric syphilitic granulomas of the elbow. *Arch. Dermat. & Syph.* 46:375, 1912.
- Wile, U. J. and Belote, G. H. Syphilitic alopecia. *Arch. Dermat. & Syph.* 13:495, 1926.
- , Wieder, L. and Warthin, A. S. Malignant syphilis. *Am. J. Syph.* 14:1, 1930.

# 13

## Fungus Diseases

### DERMATOPHYTOSIS (TINIA)

The following are the most important fungi causing superficial fungus infections *Microsporum canis* (lanosum) *Microsporum audouinii* *Trichophyton mentagrophytes* (gypseum) *Trichophyton rubrum* (purpureum) *Epidermophyton floccosum* *Trichophyton* (*Achorion*) *schoenleini* *Malassezia furfur* and *Nocardia* (*Actinomyces*) *minutissima*

Clinically nine types of superficial fungus infections can be recognized (1) tinea capitis (2) tinea barbae (3) tinea corporis (4) tinea cruris (5) dermatophytosis of the feet and the hands (6) onychomycosis (7) furus (8) tinea versicolor and (9) erythrasma

Tinea capitis which occurs almost exclusively in children usually is caused either by *Microsporum canis* or by *Microsporum audouinii*. The affected hairs tend to break off. *Microsporum canis* may evoke pronounced inflammation of the scalp so called kerion Celsi while *Microsporum audouinii* usually produces little inflammatory reaction.

Tinea barbae rare in the United States is usually caused by *Trichophyton mentagrophytes*. It is characterized by a boggy inflammatory infiltration in the bearded region of men.

Tinea corporis if caused by *Microsporum canis* manifests itself as multiple annular lesions with an active often vesicular border and central clearing. If caused by *Trichophyton mentagrophytes* one finds only one or at the most a few annular lesions showing little or no central clearing. If caused by *Trichophyton rubrum* there are fairly sharply demarcated sheetlike areas of erythema with slight scaling.

Tinea cruris usually caused by *Epidermophyton floccosum*, produces areas of erythema and scaling with a slightly elevated often vesicular border on the opposing surfaces of the skin between the scrotum and the thighs and in the perineal and the gluteal regions.

Dermatophytosis of the feet and the hands usually caused by *Trichophyton mentagrophytes* and occasionally by *Epidermophyton floccosum* is characterized by maceration between and underneath



internal organs. However, it may be pointed out that late syphilis may cause two types of reactions in the internal organs: gumma and diffuse interstitial inflammation. The latter reaction is more common.

The gummas in internal organs show the same histologic changes as those observed in the skin.

Diffuse interstitial syphilitic inflammation manifests itself as accumulations of lymphocytes and plasma cells around the small vessels. Caseation is usually absent. As a result of the long continued inflammation there is gradual degeneration of the parenchymatous structures and their replacement by fibrous tissue (Boyd). This type of reaction produces, for instance, syphilitic hepatic cirrhosis. Syphilis of the aorta is also of the diffuse type. The process starts around the vasa vasorum of the adventitia. The inflammatory infiltrate then extends along the vasa vasorum into the media. In the media one observes foci of inflammatory cells, mitotic necrosis and extensive destruction of elastic tissue.

General paresis shows perivascular accumulations of lymphocytes and plasma cells in the meninges and in the cortex of the brain. In addition there is extensive degeneration of the cortex. In contrast to general paresis, tabes does not show inflammatory changes. The essential lesion is degeneration of the posterior columns of the spinal cord.

## BIBLIOGRAPHY

### Syphilis

- Allen, A. C. Survey of pathologic studies of cutaneous diseases during World War II. *Arch. Dermat. & Syph.* 57:19, 1948.
- Boyd, W. A. *Textbook of Pathology*, ed. 4, pp. 181, 182, 870, 880. Philadelphia: Lea & Febiger, 1943.
- Fordyce, J. A. The pathology of syphilis. *J. Cutan. Dis.* 33:802, 1915.
- Fraser, J. F. The pathology of congenital syphilis. *Arch. Dermat. & Syph.* 1:491, 1920.
- Freeman, H. E. Juxta-articular nodules. *Arch. Dermat. & Syph.* 43:206, 1941.
- Hu, C. K., Liu, Y., Chen, K. C. and Frazier, C. N. Isolation of virulent *Treponema pallidum* from human aorta 32 hours after death from cardiovascular syphilis. *Am. J. Med.* 1:301, 1946.
- Jessner, M. Ueber syphilitische juxta-artikuläre Knotenbildungen. *Arch. f. Dermat. u. Syph.* 152:132, 1926.
- Katz, F. and Newton, B. L. Syphilitic juxta-articular nodules. *Arch. Dermat. & Syph.* 48:626, 1943.
- Michelson, H. E. The occurrence of tuberculous reactions in the inguinal glands of early syphilis. *Arch. Dermat. & Syph.* 19:66, 1929.
- Tuta, J. A. and Coombs, R. A. Symmetric syphilitic granuloma of the elbow. *Arch. Dermat. & Syph.* 46:375, 1942.
- Wile, U. J. and Belote, G. H. Syphilitic alopecia. *Arch. Dermat. & Syph.* 13:40, 1926.
- , Wieder, I. and Warthin, A. S. Malignant syphilis. *Am. J. Syph.* 14:1, 1930.

## Fungus Diseases

## DERMATOPHYTOSIS (TINEA)

The following are the most important fungi causing superficial fungus infections *Microsporum canis* (lanosum) *Microsporum audouinii* *Trichophyton mentagrophytes* (gypseum) *Trichophyton rubrum* (purpureum) *Epidermophyton floccosum*, *Trichophyton* (*Achorion*) *schoenleinii* *Malassezia furfur* and *Noctuidia* (*Actinomyces*) *minutissima*

Clinically nine types of superficial fungus infections can be recognized (1) tinea capitis (2) tinea barbae (3) tinea corporis (4) tinea cruris (5) dermatophytosis of the feet and the hands (6) onychomycosis (7) favus (8) tinea versicolor and (9) erythrasma

*Tinea capitis* which occurs almost exclusively in children usually is caused either by *Microsporum canis* or by *Microsporum audouinii*. The affected hairs tend to break off. *Microsporum canis* may evoke pronounced inflammation of the scalp so called kerion Celsi while *Microsporum audouinii* usually produces little inflammatory reaction.

*Tinea barbae* rare in the United States is usually caused by *Trichophyton mentagrophytes*. It is characterized by a boggy inflammatory infiltration in the bearded region of men.

*Tinea corporis* if caused by *Microsporum canis*, manifests itself as multiple annular lesions with an acute often vesicular border and central clearing. If caused by *Trichophyton mentagrophytes*, one finds only one or at the most a few annular lesions showing little or no central clearing. If caused by *Trichophyton rubrum*, there are fairly sharply demarcated sheetlike areas of erythema with slight scaling.

*Tinea cruris* usually caused by *Epidermophyton floccosum*, produces areas of erythema and scaling with a slightly elevated often vesicular border on the opposing surfaces of the skin between the scrotum and the thighs and in the perineal and the gluteal regions.

Dermatophytosis of the feet and the hands usually caused by *Trichophyton mentagrophytes* and occasionally by *Epidermophyton floccosum*, is characterized by maceration between and underneath

the toes and an erythematous scaling and vesicular eruption on the feet and the hands especially the soles and the palms

Onychomycosis caused usually by *Trichophyton rubrum* shows disintegration of the nail substance

Fungus rare in this country is caused by *Trichophyton (Achorion) schoenleini*. Most commonly it affects the scalp where it produces inflammation with formation of perifollicular crusts called scutula. Destruction of the hair ensues. Healing takes place with scarring.

Tinea versicolor caused by *Malasseia furfur* most commonly affects the upper trunk where one finds areas of brownish discoloration. The surface of the discolored areas shows fine branny scaling.

Erythrasma caused by *Nocardia minutissima* consists of circumscribed reddish brown slightly scaling patches in the axillae and the groins.

**Histopathology** For the demonstration of fungi in histologic sections the periodic acid Schiff stain of Hotchkiss McManus is of very great value since it stains fungi deeply red while almost all other structures stain a very pale pink (Kligman and Mescon). It should always be used when the presence of fungi is suspected. The reason for the strongly positive reaction of fungi to the periodic acid Schiff (PAS) stain is that the cell walls of fungi are composed of mixtures of cellulose and chitin which are rich in polysaccharides (Kligman Mescon and DeLamater).

In histologic sections fungi may present two structures hyphae (or mycelia) and spores. Hyphae which are nonreproductive appear as threadlike structures they may be septate or nonseptate. The spores which represent reproductive cells appear as round bodies.

**TINEA OF THE GLABROUS SKIN** In tinea of the glabrous skin fungi are present in small numbers with the exception of tinea versicolor in which the causative organism *Malasseia furfur* is always present in abundance. The fungi are found situated mainly in the upper two thirds of the stratum corneum. They may penetrate as far down as the upper layer of the stratum granulosum but they do not occur between living cells. In tinea of the glabrous skin except tinea versicolor the fungi occur as short and long segmented and occasionally granular hyphae extending parallel to the surface. Spores are usually not observed (Peck). In tinea versicolor one sees not only hyphae but also spores.

In the absence of fungi no diagnostic picture is presented by fungus infections of the glabrous skin. In most instances depending on the degree of reaction of the skin to the presence of fungi one sees the histologic picture of either an acute a subacute or a chronic dermatitis (see p 68). Dermatophytosis of the hands and the feet

this with intra epidermal

**TINEA CAPITIS AND TINEA BARBAE** In tinea capitis and tinea barbae fungi are present not only in the horny layer of the skin but also within and around the hair. Kligman, who recently has studied histologically the sequence of events in infections of human hair with *Microsporum canis* and *Microsporum audouinii* found that




FIG. 102. Favus. Periodic acid Schiff stain. The cross section of a hair follicle is shown. The fungus *Trichophyton schoenleinii* is present largely as mycelia within and around the hair ( $\times 400$ ).

after infection of the scalp surface hyphae grow downward into the follicle on the hair's surface a short distance and then penetrate into the hair. Inside the hair they form branches and grow downward but not farther than the exact limit of the zone of keratin synthesis. Thus as in the surface epidermis only fully keratinized structures are invaded. After the invasion of the hair ectothrix spores are formed by segmentation of external branches of intrapillary hyphae. Within the hair, however, the hyphae form no spores. As the infection subsides there is gradual reduction in the quantity of intrapillary hyphae and ectothrix spores.

Aside from the presence of fungi the histologic picture in tinea capitis and tinea barbae may be merely one of subacute or chronic dermatitis. Infection Celsi and tinea barbae however folliculitis

the toes and an erythematous, scaling and vesicular eruption on the feet and the hands, especially the soles and the palms

Onychomycosis, caused usually by *Trichophyton rubrum*, shows disintegration of the nail substance

Favus, rare in this country, is caused by *Trichophyton (Achorion) schoenleinii*. Most commonly it affects the scalp, where it produces inflammation with formation of perifollicular crusts, called scutula. Destruction of the hair ensues. Healing takes place with scarring.

Tinea versicolor, caused by *Malassezia furfur*, most commonly affects the upper trunk, where one finds areas of brownish discoloration. The surface of the discolored areas shows fine branny scaling.

Erythrasma, caused by *Nocardia minutissima*, consists of circumscribed, reddish brown, slightly scaling patches in the axillae and the groins.

**Histopathology.** For the demonstration of fungi in histologic sections, the periodic acid Schiff stain of Hotchkiss McManus is of very great value since it stains fungi deeply red while almost all other structures stain a very pale pink (Kligman and Mescon). It should always be used when the presence of fungi is suspected. The reason for the strongly positive reaction of fungi to the periodic acid-Schiff (PAS) stain is that the cell walls of fungi are composed of mixtures of cellulose and chitin which are rich in polysaccharides (Kligman, Mescon and DeLamater).

In histologic sections fungi may present two structures: hyphae (or mycelia) and spores. Hyphae, which are nonreproductive, appear as threadlike structures, they may be septate or nonseptate. The spores which represent reproductive cells appear as round bodies.

**TINEA OF THE GLABROUS SKIN.** In tinea of the glabrous skin, fungi are present in small numbers with the exception of tinea versicolor in which the causative organism, *Malassezia furfur*, is always present in abundance. The fungi are found situated mainly in the upper two thirds of the stratum corneum. They may penetrate as far down as the upper layer of the stratum granulosum, but they do not occur between living cells. In tinea of the glabrous skin, except tinea versicolor, the fungi occur as short and long segmented and occasionally granular hyphae extending parallel to the surface. Spores are usually not observed (Peck). In tinea versicolor one sees not only hyphae but also spores.

In the absence of fungi, no diagnostic picture is presented by fungus infections of the glabrous skin. In most instances, depending on the degree of reaction of the skin to the presence of fungi, one sees the histologic picture of either an acute, a subacute or a chronic dermatitis (see p. 68). Dermatomyiasis of the hands and the feet

TABLE 4—HISTOLOGIC APPEARANCE OF THE TISSUE AND OF THE FUNGI IN FUNGUS DISEASES

DISEASE	HISTOLOGIC APPEARANCE OF TISSUE	AVERAGE SIZE OF FUNGUS (MICRONS)	APPEARANCE OF FUNGUS IN TISSUE
Moniliasis	When invasive nonspecific granulation tissue with abscess formation	4	Hyphae and budding as well as nonbudding yeast cells (spores)
Blastomycosis	Epithelial hyperplasia tuberculoid granulation tissue with formation of small abscesses	10	Thick walled spores in giant cells and tissue Budding forms
Torulosis	Chronic inflammatory infiltrate with <u>extensive caseation</u>	7	Spores with wide gelatinous <u>capsule</u>
Chromoblastomycosis	Like blastomycosis	10	Thick walled, dark brown spores, often in clusters Some cells possess <u>cross walls</u>
Coccidioidomycosis	Granulomatous nodules like blastomycosis except that <u>cystic</u> may occur Subcutaneous abscesses central caseation surrounded by tuberculoid granulation tissue	40	Thick walled spores with granular cytoplasm The larger spores contain <u>endospores</u>
Actinomycosis	Nonspecific granulation tissue with abscess formation	150	Large irregularly lobulated granules with branching filaments and club formation at the periphery
Sporotrichosis	Primary lesion nonspecific granulation tissue Subcutaneous nodules three zones chronic suppurative tuberculoid and syphilitic zone	(5)	Usually, no fungi are seen Occasionally, asteroid forms of spores are present
Histoplasmosis	Chronic granuloma with foci of necrosis	3	Numerous spores surrounded by a clear halo lie in the cytoplasm of large histocytes

and foreign body giant cells. The infiltrate may extend into the subcutis. As a rule *Candida albicans* is present only in the stratum corneum and not in the dermis (Hauser and Rothman).

In the rare instances of abscess formation and ulceration due to *Candida albicans* hyphae as well as spores have been found in the dermis especially in the abscesses and their walls. The hyphae measure about 4 microns in diameter, have a distinct capsule and

and perifolliculitis (see page 168) frequently are seen. At first, an intrafollicular abscess forms. The wall of the follicle may then rupture, whereafter the perifollicular tissue becomes the seat of an abscess. After discharge of the pus, the abscess cavity begins to fill in with granulation tissue. At that stage, epithelioid cells and foreign body giant cells are frequently present, digesting remnants of the follicular epithelium (McCarthy). It may be pointed out that the histologic pictures of tinea barbae and folliculitis barbae (sycosis vulgaris) are essentially the same (see page 169).

**FAVUS.** The fungus is present in the horny layer of the skin, within and around the hairs (Fig. 105), and in the scutula. In some instances, the fungus invades the stratum malpighii and even the dermis (Ormsby and Montgomery). The scutula are composed at their periphery of well preserved hyphae and spores while in the center one finds granular debris and degenerated spores. The scutula rest upon an atrophic epidermis. The hair follicles underlying the scutula are destroyed. The dermis shows a mild to moderately severe inflammatory reaction.

### MONILIASIS

*Candida albicans*, the cause of moniliasis, not only is a cutaneous pathogen but also may affect mucosal surfaces, such as the oral mucosa and the vagina. In rare instances *Candida albicans* may infect the lungs or the meninges and cause death.

Clinically, cutaneous moniliasis usually is a superficial infection characterized by fairly sharply demarcated areas of erythema with or without pustulation. The intertriginous areas are predominantly affected. Paronychia is common. In rare instances granulomatous and hyperkeratotic lesions form on the face and the scalp of children so called monilial granuloma and, in adults abscess formation and ulceration may occur.

The association of cutaneous and mucosal moniliasis with hypoadrenalism (Talbot, Butler and MacLachlin) and with hypoparathyroidism (Sutphin, Albright and McCune) is known to occur. Since this association is often familial, it is likely that both the endocrine deficiency and the susceptibility to monilial infection in these cases are due to some defect in the germ plasm.

**Histopathology.** Superficial cutaneous moniliasis presents a histologic picture like that found in subacute or chronic dermatitis. The fungus elements occur only in the stratum corneum. They consist of hyphae and spores, some of the latter in a budding stage.

Monilial granuloma shows hyperkeratosis, acanthosis and, in the dermis, a dense infiltrate of lymphocytes, plasma cells, neutrophils

TABLE 4 — HISTOLOGIC APPEARANCE OF THE TISSUE AND OF THE FUNGI IN FUNGUS DISEASES

DISEASE	HISTOLOGIC APPEARANCE OF TISSUE	AVERAGE SIZE OF FUNGUS (MICRONS)	APPEARANCE OF FUNGI IN TISSUE
Moniliasis	When invasive nonspecific granulation tissue with abscess formation	4	Hyphae and budding as well as nonbudding yeast cells (spores)
Blastomycosis	Epithelial hyperplasia, tuberculoid granulation tissue with formation of small abscesses	10	Thick walled spores in giant cells and tissue Budding forms
Torulosis	Chronic inflammatory infiltrate with <u>extensive caseation</u>	7	Spores with wide gelatinous <u>capsule</u>
Chromoblastomycosis	Like blastomycosis	10	Thick walled, dark brown <u>spores</u> , often in clusters Some cells possess <u>cross walls</u>
Coccidioidomycosis	Granulomatous nodules like blastomycosis except that <u>caseation</u> may occur Subcutaneous abscesses central caseation surrounded by tuberculoid granulation tissue	40	Thick walled spores with granular cytoplasm The larger spores contain <u>endospores</u>
Actinomycosis	Nonspecific granulation tissue with abscess formation	150	Large, irregularly lobulated granules with branching filaments and club formation at the periphery
Sporotrichosis	Primary lesion nonspecific granulation tissue Subcutaneous nodules three zones chronic suppurative tuberculoid and syphilitoid zone	(5)	Usually, no fungi are seen Occasionally, asteroid forms of spores are present
Histoplasmosis	Chronic granuloma with foci of necrosis	3	Numerous spores surrounded by a clear halo like in the cytoplasm of large histiocytes

and foreign body giant cells The infiltrate may extend into the subcutis As a rule *Candida albicans* is present only in the stratum corneum and not in the dermis (Hauser and Rothman)

In the rare instances of abscess formation and ulceration due to *Candida albicans*, hyphae as well as spores have been found in the dermis especially in the abscesses and their vicinity The spores measure about 4 microns in diameter, have a distinct capsule and



are Gram positive. They often lie in clusters and budding forms are occasionally seen. The hyphae are slender rods and have a beaded or segmented appearance (Rockwood and Greenwood, Danbolt)

### NORTH AMERICAN BLASTOMYCOSIS (GILCHRIST'S DISEASE)

North American blastomycosis, caused by *Blastomyces dermatitidis*, is characterized by granulomatous and suppurative lesions which may occur in any organ of the body but are found most commonly in the skin, the lungs and the bones. Two clinical forms can be differentiated, a primary cutaneous and a systemic form. In the benign primary cutaneous form, one observes either one or a few rather large lesions. In the fatal systemic form, the cutaneous lesions are numerous and usually small in size.

The cutaneous lesions consist in both forms of verrucous plaques showing an active border beset with a large number of minute abscesses. In the primary cutaneous form, the older lesions show central healing with atrophy. In the systemic form the cutaneous lesions may undergo ulceration and, in addition, subcutaneous abscesses may occur.

**Histopathology.** Histologic examination shows xanthosis, papillomatosis and considerable downward proliferation of the epidermis. The downward proliferation may be marked enough to present the picture of pseudo-epitheliomatous hyperplasia. Intra-epidermal abscesses are often present. Occasionally, one finds Langhans giant cells completely enclosed by the proliferated epidermis (Fig. 106).

The dermis is infiltrated by a polymorphous granulation tissue. Polymorphonuclear leukocytes usually are present in large numbers and form small abscesses in the dermis. Langhans giant cells are scattered throughout the dermis. They usually lie alone and not within groups of epithelioid cells. Occasionally, one observes tubercloid formations but never true tubercles.

The cells, oospores, of *Blastomyces dermatitidis* are fairly <sup>↑</sup>numerous in histologic sections. They are often found lying free in the tissue, particularly in the abscesses. They occur, however, in their largest number within the giant cells. One or several spores may lie within a giant cell (Fig. 107). When in this location, the spores are easily spotted, even at low magnification, in sections stained with routine stains. Being unstained, they resemble small, round holes punched out of the cytoplasm of the giant cells. On high magnification, the spores are seen to have a thick wall, which gives them a double contoured appearance. They measure from 8 to 15 microns in diameter, on the average 10 microns. *Blastomyces* reproduce by



FIG 107 North American blastomycosis High magnification of Figure 106 Three blastomycetes cells (B.C.) are shown lying in the cytoplasm of giant cells (X400)

budding, and occasionally budding forms are seen in sections. Like in most fungus infections, many more spores are seen in sections stained with the Hotchkiss McManus stain than in routinely stained sections (see page 218)

The histologic appearance of the visceral lesions is analogous to that of the cutaneous lesions. The number of neutrophils is often great and numerous abscesses tend to be present (Littman, Wicker and Warren)

**Differential Diagnosis.** Tuberculosis verrucosa cutis, torulosis and chromoblastomycosis must be considered in the differential diagnosis. Tuberculosis verrucosa cutis shows no spores in the tissue. In addition, the number of neutrophils is much smaller and one usually finds true tubercles and areas of caseation necrosis. The distinctive features of torulosis and chromoblastomycosis are discussed under these respective headings (see below)

### TORULOSIS (CRYPTOCOCCOSIS, EUROPEAN BLASTOMYCOSIS OF BUSSE AND BUSCHKE)

Torulosis, though very rare, occurs throughout the world. The disease, caused by *Cryptococcus neoformans* (*Torula histolytica*), usually is chronic and systemic and ends in death. The brain and the meninges are nearly always affected, resulting in meningoencephalitis with presence of the organisms in the spinal fluid. The lungs also are commonly involved. Cutaneous lesions are present in about 10 per cent of the cases, and, in rare instances, lesions of the oral mucosa occur (Cawley). The cutaneous lesions may be the first clinical sign of the disease (Linell, Magnusson and Norden), and in some instances the disease may remain limited to the skin (Gandy). In that case, the course of the disease is benign and self limited.

The occurrence of torulosis has been reported either as a silent or as an active infection in patients with various types of lymphoma such as Hodgkin's disease (Geudel, Ende and Norman) and various types of leukemia (Cawley). A similar association with lymphoma occurs also in histoplasmosis (see page 231).

The cutaneous lesions are variable and may consist of papules, pustules, nodules, infiltrated plaques, ulcers or subcutaneous abscesses.

**Histopathology.** Histologic examination of the skin shows, in most instances, a chronic inflammatory infiltrate with or without giant cells. Occasionally, the inflammatory reaction is mild and extensive caseation is present in the dermis.

The causative fungus usually is present in abundance, either within

giant cells or lying free in the tissue. It consists of a spherical spore measuring from 5 to 10 microns in diameter which multiplies like *Blastomyces dermatitidis* by budding. It usually is surrounded by a wide gelatinous capsule which does not stain with hematoxylin and eosin but stains red with the periodic acid Schiff reaction and methen

(Linnell Mag  
capsule is absent

(Wile) The fungus then greatly resembles *B. dermatitidis* and easily can be confused with it especially since both may occur in giant cells. On cultures however in contrast with *Blastomyces* *Torula* forms no mycelia (Benham)

### CHROMOBLASTOMYCOSIS

This disease is limited to the skin and is benign. It is caused by three closely related fungi which appear alike in the tissue. Hormo




FIG. 108. Chromoblastomycosis. Periodic acid Schiff stain. Epithelioid cells and Langhans giant cells form the wall of an abscess. Two long chains of fungus cells are located in the right upper corner. In addition three fungus cells lie in the right center. (X400)

*dendrium pedrosi*, *Phialophora verrucosa* and *Hormodendrum compactum* (French and Russell)

The cutaneous lesions usually limited to a single area consist of densely aggregated nodules and plaques with a hard verrucous surface.

**Histopathology.** The histologic appearance of the skin is very much like that of blastomycosis. The epidermis shows considerable hyperplasia although rarely to the extent that is usual in blastomycosis. The dermis shows an intensive infiltration, with a polymorphous granulation tissue containing many Langhans giant cells and small abscesses composed of polymorphonuclear leukocytes. Tuberculoid formations may be present, but true tubercles and caseation are absent, just as in blastomycosis.

The causative organisms are found in Langhans giant cells as well as free in the tissue and are especially numerous in the abscesses. They appear as sclerotic, dark brown, thick-walled, usually spherical spores varying in size from 8 to 15 microns. They lie either singly or in chains or clusters (Fig. 108). Reproduction is by intracellular wall formation and splitting not by budding. In some of the organisms cross walls can be seen.

**Differential Diagnosis.** Chromoblastomycosis cannot be differentiated from blastomycosis except through the different appearance of the fungi. Just as blastomycosis, chromoblastomycosis differs from tuberculosis verrucosa cutis by the presence of a fungus in the tissue, and the absence of true tubercles and of caseation.

### COCCIDIOIDOMYCOSIS (SAN JOAQUIN VALLEY FEVER)

Coccidioidomycosis is caused by *Coccidioides immitis*. It is endemic in the southwestern United States, especially in the San Joaquin valley of California, and in northern Mexico. Three forms are recognized, primary, intermediate and progressive (Duemling).

Primary coccidioidomycosis represents an acute respiratory infection. Development of erythema nodosum is not uncommon.

Progressive coccidioidomycosis (coccidioidal granuloma) follows the primary form in a small percentage of cases after a varying length of time. It has a very high mortality. Many organs, especially the meninges, the lungs, the bones and the lymph nodes may be involved. Cutaneous lesions are common. They consist either of verrucous and granulomatous nodules or of subcutaneous cold abscesses which may break through the skin.

**Histopathology.** The nodose lesions of primary coccidioidomycosis have the same histologic appearance as idiopathic erythema nodosum (Winer).

The verrucous and the granulomatous lesions of coccidioidal granuloma resemble blastomycosis in their histologic aspects. How

ever there is less tendency to abscess formation, and caseation necrosis may occur (Moore). The causative organisms are found free in the tissue as well as in Langhans giant cells. As a rule they are present in large number.




FIG 109 Coccidioides  
larger than those of size. Their cytoplasm

The subcutaneous abscesses of coccidioidal granuloma resemble scrofuloderma in their appearance.




FIG 110 Coccidioidomycosis. A large Coccidioides spore lies within a giant cell. The Coccidioides spore contains numerous endospores ( $\times 400$ ).

thelioid cells and some giant cells. Numerous spores are present extracellularly as well as intracellularly in giant cells.

The spores of *C. immitis* are 80 microns

10 to  
Thus

*Coccidioides* is much larger than *Blastomyces*, *Torula* or *Phialophora*. The spores are spherical and thick walled and have a granular cytoplasm. Multiplication is not by budding but by formation of endospores which may be seen lying inside the larger spores (Fig. 110). The endospores are released into the tissue by rupture of the wall of the spore. Endospores may measure up to 5 microns in diameter.

**Differential Diagnosis.** A diagnosis of coccidioidomycosis can be made only in the presence of the fungus.

## ACTINOMYCOSIS

Actinomycosis is caused either by *Actinomyces* bovis, which is anaerobic or microaerophilic, or by several species of *Nocardia* (e.g. *N. asteroides*, *N. madurae*), which are aerobic.

Actinomycosis frequently affects the skin. Involvement may be primary, as in Madura foot (mycetoma). More often, however, the infection reaches the skin from an internal focus. The most common form is cervicofacial actinomycosis, in which case the organism reaches the skin from the mouth, next in frequency are actinomycosis of the thoracic skin secondary to involvement of a lung and actinomycosis of the abdominal skin secondary to involvement of the cecum or the appendix.

The involved skin is dark red in color, possesses a "wooden" type of hardness and shows numerous sinuses discharging a serosanguineous or purulent fluid containing sulfur yellow granules which consist of masses of fungi.

**Histopathology.** Histologic examination shows extensive granulation tissue containing large abscesses. The fungus granules are found within the abscesses. The granulation tissue is nonspecific in its appearance. In the early phase of the disease it is composed of neutrophils, eosinophils, lymphocytes, plasma cells, histiocytes and fibroblasts. In the healing phase fibroblasts predominate. Thus, the diagnosis can be established only by finding the *Actinomyces* granules in the abscesses. When selecting an area for biopsy, an area containing purulent material should be chosen.


The fungus granules are large and may measure several hundred microns in diameter, large enough to be visible macroscopically, as the so called sulfur granules. In histologic sections, they appear briso

(Fig. 111). They are homogeneous branching filaments at the periphery.

The ends of the filaments frequently are surrounded by a gelatinous sheath, giving the ends a club shaped appearance. The filaments are much better seen in sections stained by Gram's method, which stains them Gram positive, than in sections stained with routine stains. The

cells immediately around the granules are usually polymorphonuclear leukocytes, but foreign body giant cells occasionally are seen in contact with the granules.

In general, *Nocardia* produces more necrosis and less granulation tissue and less fibrosis than *Actinomyces*. The granules of *Nocardia* are smaller and much less numerous than those of *Actinomyces* so that multiple sections are necessary to find them (Weed and Baggen



ACTINOMYCETOSIS A large sulfur granule is shown in the center of the field. The granule appears homogeneous in the center. Clubs are present at the periphery ( $\times 100$ )

stoss). Nevertheless, these differences are not sufficiently distinct and cultural studies are necessary for a differentiation of *Nocardia* from *Actinomyces*.

## SPOROTRICHOSIS

SPOROTRICHOSIS

*Sporotrichum*, affects only the benign disease. Most

type of infection. In that type there is a primary lesion (sporotrichotic chancre) which is commonly an ulcer but may be papillomatous. Secondly, multiple subcutaneous nodules appear along the lymphatic drainage area. The nodules are

Much seminate

are widely scattered lesions consisting the localized type, form



*Coccidioides* is much larger than *Blastomyces*, *Torula* or *Phialophora*. The spores are spherical and thick walled and have a granular cytoplasm. Multiplication is not by budding but by formation of endospores which may be seen lying inside the larger spores (Fig. 110). The endospores are released into the tissue by rupture of the wall of the spore. Endospores may measure up to 5 microns in diameter.

**Differential Diagnosis.** A diagnosis of coccidioidomycosis can be made only in the presence of the fungus.

### ACTINOMYCOSIS

Actinomycosis is caused either by *Actinomyces bovis* which is anaerobic or microaerophilic or by several species of *Nocardia* (e.g. *N. asteroides*, *N. madurae*) which are aerobic.

Actinomycosis frequently affects the skin. Involvement may be primary as in Madura foot (mycetoma). More often however the infection reaches the skin from an internal focus. The most common form is cervicofacial actinomycosis in which case the organism reaches the skin from the mouth. Next in frequency are actinomycosis of the thoracic skin secondary to involvement of a lung and actinomycosis of the abdominal skin secondary to involvement of the cecum or the appendix.

The involved skin is dark red in color, possesses a wooden type of hardness and shows numerous sinuses discharging a serosanguineous or purulent fluid containing sulfur yellow granules which consist of masses of fungi.

**Histopathology.** Histologic examination shows extensive granulation tissue containing large abscesses. The fungus granules are found within the abscesses. The granulation tissue is nonspecific in its appearance. In the early phase of the disease it is composed of neutrophils, eosinophils, lymphocytes, plasma cells, histiocytes and fibroblasts. In the healing phase fibroblasts predominate. Thus the diagnosis can be established only by finding the *Actinomyces* granules in the abscesses. When selecting an area for biopsy an area containing purulent material should be chosen.

The fungus granules are large and may measure several hundred microns in diameter, large enough to be visible macroscopically as the so called sulfur granules. In histologic sections they appear brown.

111) They are homogeneous in filaments at the periphery.

The ends of the filaments frequently are surrounded by a gelatinous sheath giving the ends a club shaped appearance. The filaments are much better seen in sections stained by Gram's method which stains them Gram positive, than in sections stained with routine stains. The

thick walled round spores are found free in the tissue visible even with hematoxylin eosin staining. Some of them have radiating asteroid elongations (Fig 112) (Moore and Ackerman Pinkus and Grekim)

**Differential Diagnosis** Without the demonstration of the fungus a diagnosis of sporotrichosis cannot be made but can only be suspected. The subcutaneous nodules of tularemia may have the same histologic appearance. Erythema induratum differs from sporotrichosis by its vascular changes and the presence of caseation which is absent in sporotrichosis.

## HISTOPLASMOVIS

Histoplasmosis caused by the fungus *Histoplasma capsulatum* in most instances is a fatal systemic disease. Occasionally the disease is limited to the skin or the mucous membranes and benign in its course (Curtis and Grekin).

In the systemic form of histoplasmosis the clinical picture is variable. The lymph nodes are often enlarged markedly. Pulmonary and adrenal involvement may be a prominent feature. Cutaneous or mucomembranous lesions occur in about half the patients with the systemic form of infection (Miller Keddie Johnstone and Bosick).

Several authors have reported the simultaneous occurrence of histoplasmosis and lymphoma (Ende and Curtis) and to the lymphoma especially in view of the fact that also torulosis has been observed in association with lymphoma (see page 224).

The skin lesions of histoplasmosis in both the systemic and the cutaneous form may be nodular granulomatous or ulcerative.

**Histopathology** Histologic examination reveals a chronic non specific granulomatous infiltrate with foci of necrosis. Throughout this infiltrate but especially in the vicinity of the necrotic foci one observes large histiocytes (macrophages) engorged with numerous organisms.

*Histoplasma capsulatum* appears in sections stained with hematoxylin and eosin as small round bodies surrounded by a thin capsule and a wide clear space. The organism measures 2 to 3 microns in diameter. On staining with the periodic acid Schiff reaction however *Histoplasma capsulatum* shows no capsule but instead the body appears larger and lined by a definite red stained cell wall. Therefore it appears that the so-called halo or

**Histopathology.** The primary lesion of sporotrichosis shows a non specific granulation tissue containing, among other cells, many plasma cells, epithelioid cells and some Langhans giant cells. Foci of suppuration are present. If the primary lesion is papillomatous, one finds, in addition, marked epithelial hyperplasia and intra epidermal abscesses (Moore and Ackerman).

The subcutaneous nodules show a more characteristic appearance than the primary lesion because of the arrangement of the infiltrate



FIG. 112 Sporotrichosis. A large spore with radiating clefts is shown ( $\times 1950$ ) (Hermann Pinkus M.D.)

in three zones—the chronic suppurative zone, the tuberculoid zone and the syphiloid zone. The zone in the center of the nodule, the chronic suppurative zone, is composed chiefly of neutrophils with a few histiocytes and lymphocytes. Small abscesses composed entirely of neutrophils are present within this zone. The middle or tuberculoid zone is characterized by numerous epithelioid cells and a large number of giant cells of the Langhans type. The giant cells may lie in groups. The peripheral or syphiloid zone consists of a rich cellular infiltrate of plasma cells, lymphocytes and fibroblasts. The zonal arrangement is not always distinct. In older nodules one merely sees a nonspecific granuloma.

The causative organisms are present in the tissue in small numbers as round to fusiform bodies which do not stain with hematoxylin and eosin but become visible when the periodic acid Schiff reagent is employed (Kligman and Baldrige). In rare instances a few large,

## Torulosis

- Benham R W The fungi of blastomycosis and coccidioidal granuloma Arch Dermat & Syph 30 385 1934
- Cawley E P Grekin R H and Curtis A C Torulosis J Invest. Dermat 14 327 1940 (Good review)
- Candy W M Primary cutaneous cryptococcosis Arch Dermat & Syph 62 97, 1950
- Ceudel B R Ende M and Norman S L Cryptococcosis a review with special reference to apparent association with Hodgkin's disease Am J Med 9 343 1950
- Linell F Magnusson B and Norden A Cryptococcosis Acta dermat venerol 33 103 1953
- Wile U J Cutaneous torulosis Arch Dermat & Syph 31 58 1935

## Chromoblastomycosis

- French A J and Russell S R Chromoblastomycosis Arch Dermat & Syph 67 129 1953
- Lane C G A cutaneous disease caused by a new fungus (*Phialophora terricola*) J Cutan Dis 33 840 1915
- Moore M Cooper Z K and Weiss R S Chromomycosis (chromoblastomycosis) J A M A 122 1237 1943
- Weidman F D and Rosenthal L H Chromoblastomycosis a new and important blastomycosis in North America Arch Dermat & Syph 43 62 1941

## Coccidioidomycosis

- Duerming W W Progressive disseminated coccidioidomycosis Arch Dermat & Syph 60 781 1949
- Jacobsen H P Coccidioidal granuloma Arch Dermat & Syph 21 790 1930
- Moore M Mycotic granulomata and cutaneous tuberculosis a comparative study

Went

## Actinomycosis

- Moore M Radiate formation on pathogenic fungi in human tissue Arch Path 42 113 1946
- Salaman H A and Hewler J Cutaneous actinomycosis Arch Dermat & Syph 36 131 1937
- Wied L A and Baggenstoss A H Actinomycosis Am J Clin Path 19 201 1949

## Sporotrichosis

- Kligman A M and Baldridge G D Morphology of *Sporotrichum schenckii*
- A comparison of  
13 (Good review)  
formation in tissue
- and Wile R L Generalized subcutaneous gummatous ulcerating sporotrichosis Arch Dermat & Syph 31 672 1935
- Pinkus H and Grekin J N Sporotrichosis with asteroid tissue forms Arch Dermat & Syph 61 813 1950

capsule is an artefact produced by plasmolysis which causes the cytoplasm to shrink away from the cell wall (Kligman and Baldridge). *Histoplasma capsulatum* is a fungus according to its cultural characteristics and because it multiplies by budding.

**Differential Diagnosis** The general appearance of the granulomatous infiltrate with its parasitized histiocytes is much like that of rhinoscleroma, granuloma inguinale and cutaneous leishmaniasis (for their differential diagnosis see Table 5 page 237)

## BIBLIOGRAPHY

### Dermatophytosis

- Kligman A M The pathogenesis of tinea capitis due to *Microsporum audouinii* and *Microsporum canis* II Histopathologic findings Arch Dermat & Syph to be published
- and Mescon H Periodic acid Schiff stain for demonstration of fungi in animal tissue J Bact 60 115 1950
- — — and DeLamater I D The Hotchkiss McManus stain for the histopathologic diagnosis of fungus diseases Am J Clin Path 21 86 1951
- McCarthy I Histopathology of Skin Diseases p 470 St Louis Mosby 1951
- Ormsby O S and Montgomery H Diseases of the Skin ed 7 p 1117 Philadelphia Lea & Febiger 1948
- Peck S Epidermophytosis of the feet and epidermophytids of the hands Arch Dermat & Syph 22 40 1930

### Moniliasis

- Dunbolt N Deep cutaneous moniliasis a fatal case of peculiar type Acta dermat venereol 21 98 1940
- Hauser I V and Rothman S Monilial granuloma Arch Dermat & Syph 61 297 1950
- Rockwood E M and Greenwood A M Monilial infection of the skin Arch Dermat & Syph 29 574 1934
- Sutphin A Albright I and McCune D J Five cases (three in siblings) of idiopathic hypoparathyroidism associated with moniliasis J Clin Endocrin 12 625 1943
- Talbot N B Butler A M and MacLachlan E A The effect of testosterone and allied compounds on the mineral and nitrogen and carbohydrate metabolism of a girl with Addison's disease J Clin Invest 22 383 1943

### North American Blastomycosis

- Baker R D Tissue reaction in human blastomycosis (North American blastomycosis) Am J Path 18 479 1942
- Bush J D Severe generalized blastomycotic dermatitis Arch Dermat & Syph 43 485 1941
- Littman M L Wicker E H and Warren A S Systemic North American blastomycosis Am J Path 24 339 1948
- Martin D S and Smith D T Blastomycosis (American blastomycosis Gilchrist's disease) Am Rev Tuberc 39 275 488 1939
- Moore M Mycotic granulomata and cutaneous tuberculosis a comparison of the histopathologic response J Invest Dermat 6 119 1945 (Good review)

## Torulosis

- Benham R W The fungi of blastomycosis and coccidioidal granuloma Arch Dermat & Syph 30 385 1934
- Cawley E P Grekin R H and Curtis A C Torulosis J Invest Dermat 14 377 1950 (Good review)
- Candy W M Primary cutaneous cryptococcosis Arch Dermat & Syph 62 97 1950
- Crudel B R Eide M and Norman S I Cryptococcosis a review with special reference to apparent association with Hodgkin's disease Am J Med 9 313 1950
- Linell F Magnusson B and Norden A Cryptococcosis Acta dermat venereol 33 103 1953
- Wile U J Cutaneous torulosis Arch Dermat & Syph 31 58 1935

## Chromoblastomycosis

- French A J and Russell S R Chromoblastomycosis Arch Dermat & Syph 67 129 1953
- Lane C G A cutaneous disease caused by a new fungus (*Phialophora verrucosa*) J Cutan Dis 33 840 1915
- Moore M Cooper Z A and Weiss R S Chromomycosis (chromoblastomycosis) J A M A 122 1237 1943
- Weidman F D and Rosenthal L H Chromoblastomycosis a new and important blastomycosis in North America Arch Dermat & Syph 43 62 1941

## Coccidioidomycosis

- Duerming W W Progressive disseminated coccidioidomycosis Arch Dermat & Syph 60 781 1919
- Jacobsen H P Coccidioidal granuloma Arch Dermat & Syph 21 790 1930
- Moore M Mycotic granulomata and cutaneous tuberculosis a comparison of the histopathologic response J Invest Dermat 6 149 1945
- Winer L H Histopathology of the nodose lesion of acute coccidioidomycosis Arch Dermat & Syph 61 1010 1950

## Actinomycosis

- Moore M Radiate formation on pathogenic fungi in human tissue Arch Path 42 113 1946
- Salzman H A and Kessler I Cutaneous actinomycosis Arch Dermat & Syph 36 131 1937
- Wied L A and Baggenstoss A H Actinomycosis Am J Clin Path 19 201 1949

## Sporotrichosis

- Kligman A M and Baldridge G D Morphology of *Sporotrichum schenckii* and *Histoplasma capsulatum* Arch Path 51 367 1921
- Moore M Mycotic granulomata and cutaneous tuberculosis A comparison of the histopathologic response J Invest Dermat 6 149 1945 (Good review)
- and Ackerman L A Sporotrichosis with radiate formation in tissue Arch Dermat & Syph 53 253 1916
- and Kile R L Generalized subcutaneous gummatous ulcerating sporotrichosis Arch Dermat & Syph 31 672 1935
- Pirkus H and Grekin J A Sporotrichosis with asteroid tissue forms Arch Dermat & Syph 61 813 1950

## Histoplasmosis

- Cawley E P and Curtis A C Histoplasmosis and lymphoblastoma are these diseases related? J Invest Dermat 11 443 1948
- Curtis A C and Grekin J N Histoplasmosis JAMA 134 1217 1947
- Ende N Pizzolito P and Ziskind J Hodgkin's disease associated with histoplasmosis Cancer 5 763 1952
- Kligman A M and Baldrige G D Morphology of *Sporotrichum schenckii* and *Histoplasma capsulatum* in tissue Arch Path 51 567 1951
- Miller H E Keddie F M Johnstone H G and Bostick W L Histoplasmosis Arch Dermat & Syph 56 715 1947 (Review of cutaneous lesions)
- Parsons R J and Zarafonitis C J D Histoplasmosis in man Arch Int Med 75 1 1945 (General review)

epidermis grafted on the chorioallantois of chick embryos with the fluid obtained from herpes zoster vesicles

It is probable that the viruses of varicella and herpes zoster are identical. There exists strong clinical evidence (Brausgaard) and experimental evidence (Brain) for this contention. Furthermore the two viruses have an identical appearance under the electron microscope (Baldrige Blank and Rake). Wise and Sulzberger believe that herpes zoster represents an immune or allergic form of varicella which occurs in patients who had previous clinical or subclinical varicella.

**Clinical Appearance** Varicella shows a generalized eruption which at first consists of papules. After two or three days the papules are transformed into vesicles which are characteristically umbilicated. After three more days the vesicles change into pustules. In severe cases in addition there may be purpuric lesions. It is characteristic of varicella that all lesions are at the same stage of development.

The eruption of Varicella is also generalized. The lesions begin as small papules which soon develop into vesicles. The vesicles crust over as a rule without changing into pustules. Lesions occur in successive crops so that one observes lesions in different stages of development.

b

c

Varicella eruption there is a generalized vesicular eruption indistinguishable from that of varicella. Herpes zoster (especially the generalized form) is apt to occur in lymphoma (see page 491).

Herpes simplex shows one or several groups of vesicles on a mildly inflammatory base. The eruption may occur anywhere on the skin but is found most commonly about the face and the genitalia.

**Histopathology** The characteristic histologic lesion in the four diseases is an intra epidermal vesicle produced by profound degeneration of epidermal cells. Because of the presence of these degenerative changes the vesicle differs histologically from those seen in other

1 of Bullae page 66) and

er it is important that as

ion be selected for biopsy

invasion of inflammatory

may obscure the diagnostic features. The degeneration of the epidermal cells occurs in two forms: ballooning degeneration and reticular degeneration.

Ballooning -  
cells. Such



eosinophilic cytoplasm (Figs. 114, 115). They may have no nucleus, or may have one or many. Because of the fact that the balloon cells lose their intercellular bridges, acantholysis occurs and the cells become separated from one another. Unilocular vesicles result. The process of ballooning degeneration occurs mainly at the base of virus vesicles leading to a dissolution of the lower epidermis, so that, ult



FIG 114 Herpes zoster. Low magnification. There is marked ballooning degeneration of the cells at the floor of the vesicle. The cells of a hair follicle, shown at the left, likewise show ballooning degeneration. Reticular degeneration observed at the top of the vesicle, is only slight. Since this is an early lesion, no inflammatory reaction is present ( $\times 100$ )

mately, the originally intra epidermal vesicles become subepidermal in many places. Ballooning degeneration affects also the epithelial cells of hair follicles and sebaceous glands.

Reticular cells become

the cell wall bursts. By coalescence of neighboring, similarly affected cells, a multilocular vesicle results, the septa of which are formed by the resistant cellular walls (Fig. 116). Reticular degeneration occurs mainly at the top and the periphery of virus vesicles. In older vesicles, the resistant cellular walls disappear and the multilocular vesicle then becomes unilocular. It may be pointed out that reticular degeneration is not specific for virus vesicles since it also occurs in the vesicles of dermatitis (see page 68).

numerous polymorphonuclear leukocytes as well as some macrophages (Sheldon and Heyman). These central abscesses tend to have a triangular or quadrangular shape with elongated corners, giving them a stellate appearance (Fig. 124). The epithelioid cells surrounding the abscesses are arranged in palisade formation. As the abscesses gradually enlarge, they coalesce and lose their stellate appearance.




FIG. 124. *Lymphogranuloma venereum*. In the lymph node, a stellate abscess is present. It is surrounded by epithelioid cells in palisade formation ( $\times 100$ ).

### MILKERS' NODULES

Milkers' nodules are acquired from cows infected with paravaccinia (natural cowpox). There are, usually on the fingers, one to three and occasionally more nodules, 1 to 2 cm. in diameter, which are bluish red, semiglobular and usually painless. Spontaneous healing occurs in 1 to 2 months.

**Histopathology.** The epidermis shows acanthosis and parakeratosis. A dense nonspecific chronic inflammatory infiltrate is present in the dermis. The capillaries are increased in number, are dilated and show swelling of their endothelial cells (Norland and McKee).

While most authors have found no inclusion bodies, Katzenellenbogen observed them in some but not in all of his cases. They were

located in the cytoplasm of vacuolated epidermal cells and often were Feulgen positive. Their appearance was similar to that of the inclusion bodies observed in variola and vaccinia.

Study of smears of tissue obtained from milkers' nodules under the electron microscope has shown the presence of typical elementary bodies of the paravaccinia type (Nasimunn and Deubner).

## BIBLIOGRAPHY

## Virus Diseases

- Blind J O W and Robinow C F The inclusion bodies of vaccinia and their relationship to the elementary bodies studied in cultures of the rabbit's cornea. *J Path & Bact* 48:381 1939
- Ebert M H and Otsuka M Virus diseases of the skin with special reference to elementary and inclusion bodies. *Arch Dermat & Syph* 48:633 1943 (Good review)
- Goodpasture E W Woodruff A M and Buddingh G J Vaccinal infection of the chorioallantoic membrane of the chick embryo. *Am J Pathol* 8:91 1932

## Variola Varicella Herpes Zoster, Herpes Simplex

- Baldrige G D Blank H and Rake G Dermatologic aspects of electron microscopy. *J Invest Dermat* 16:281 1951
- Bland J O W and Robinow C F The inclusion bodies of vaccinia and their relationship to the elementary bodies studied in cultures of the rabbit's cornea. *J Path & Bact* 48:381 1939
- Blank H Coriell L L and Scott T F M Human skin grafted onto the chorioallantois of the chick embryo for virus cultivation. *Proc Soc Exper Biol & Med* 69:341 1948
- Brain R T A relation between the viruses of zoster and varicella. *Brit J Exp Path* 14:67 1933
- Bruusgaard E The mutual relation between zoster and varicella. *Brit J Dermat* 44:1 1932
- Crouse H V Coriell L L Blank H and Scott T F M Cytochemical studies on the intranuclear inclusion of herpes simplex. *J Immunol* 65:119 1950
- Denny Brown D and Adams R D Pathologic features of herpes zoster. *Arch Neur & Psych* 51:216 1944
- Downie A W and Dumbell K R The isolation and cultivation of variola virus on the chorioallantois of the chick embryo. *J Path & Bact* 59:189 1947
- Ebert M H Histologic changes in sensory nerves of the skin in herpes zoster. *Arch Dermat & Syph* 60:641 1949
- Frank L Varicella pneumoniae. *Arch Path* 50:450 1950
- Goodpasture E W and Anderson K Infection of the human skin grafted on the chorioallantois of chick embryos with virus of herpes zoster. *Am J Path* 20:447 1944
- Head H and Campbell A W Pathology of herpes zoster and its bearing on sensory location. *Brain* 23:353 1900
- Johnson H N Visceral lesions associated with varicella. *Arch Path* 30:292 1940

- Michelson H E and Ikeda K. Microscopic changes in variola Arch Dermat & Syph 15 138 1907
- Smith M G Lennette E H and Reames H R. Isolation of the virus and the demonstration of intranuclear inclusions in a case of acute encephalitis Am J Path 17 55 1941
- Wise F and Sulzberger M B. The 1937 Yearbook of Dermatology and Syphilology p 212 Chicago Yearbook Publishers 1938
- Zarafonitis C J D Smadel J E Adams J W and Haymaker W. Fatal herpes simplex encephalitis in man Am J Path 20 429 1941

#### Eczema Vaccinatum and Kaposi's Varicelliform Eruption

- Barton R L and Brunsting L A. Kaposi's varicelliform eruption Arch Dermat & Syph 50 99 1944
- Lynch F W. Kaposi's varicelliform eruption Arch Dermat & Syph 51 129 1945
- Riley K A and Callaway J I. Eczema vaccinatum J Invest Dermat 9 321 1947

#### Molluscum Contagiosum

- Rake G and Blank H. The relationship of host and virus in molluscum contagiosum J Invest Dermat 15 81 1950
- van Rooijen C F. The micromanipulation and microdissection of the molluscum contagiosum J Path & Bact 46 425 1938

#### Verruca

- Becker S W. The pigmented form of verruca plana Arch Dermat & Syph 34 965 1936
- Blank H Buerk M and Weidman F. The nature of the inclusion body of verruca vulgaris a histochemical study of the nucleotids J Invest Dermat 16 19 1951
- Bunting H Strauss M J and Banfield W G. The cytology of skin papillomas that yield virus like particles Am J Path 28 985 1952
- Lund H Z and Leuchsenberger C. The peculiar changes of nuclear chromatin in verruca vulgaris Am J Path 27 750 1951
- Lutz W. A propos de l'épidermodysplasie verruciforme Dermatolog 92 30 1946
- Strauss M J Bunting H and Melnick J L. Virus like particles and inclusion bodies in skin papillomas J Invest Dermat 15 433 1950
- — and — —. Eosinophilic inclusion bodies and cytoplasmic masses in verrucae J Invest Dermat 17 209 1951
- Waisman M and Montgomery H. Verruca plana and epithelial nevus Arch Dermat & Syph 45:29 1942 (Good review of verruca vulgaris and verruca plana)
- Wile U J and Jangery L B. The etiology of common warts J A M A 73 970 1919 76 410 1921

#### Lymphogranuloma Venereum

- Dunoy R and von Haam E. Venereal lymphogranuloma Arch Path 27 1032 1939
- Kornblith B A. Observations on lymphogranuloma venereum Surg Gynec. & Obst 63 99 1936
- Reichste H S and Connor W H. Lymphogranuloma inguinale Arch Dermat & Syph. 32 196 1935

located in the cytoplasm of vacuolated epidermal cells and often were Feulgen positive. Their appearance was similar to that of the inclusion bodies observed in variola and vaccinia.

Study of smears of tissue obtained from milkers' nodules under the electron microscope has shown the presence of typical elementary bodies of the paravaccinia type (Nasemann and Deubner).

## BIBLIOGRAPHY

## Virus Diseases

- Bland J O W and Robinow C F The inclusion bodies of vaccinia and the relationship to the elementary bodies studied in cultures of the rabbit's cornea *J Path & Bact* 48:381 1939
- Ebert M H and Otsuka M Virus diseases of the skin with special reference to elementary and inclusion bodies *Arch Dermat & Syph* 48:635 1943 (Good review)
- Goodpasture L W Woodruff A M and Buddingh G J Vaccinal infection of the chorioallantoic membrane of the chick embryo *Am J Pathol* 39:1 1932

## Variola Varicella Herpes Zoster Herpes Simplex

- Baldridge G D Blank H and Rake G Dermatologic aspects of electron microscopy *J Invest Dermat* 16:281 1951
- Bland J O W and Robinow C F The inclusion bodies of vaccinia and the relationship to the elementary bodies studied in cultures of the rabbit's cornea *J Path & Bact* 48:381 1939
- Blank H Coriell L L and Scott T F M Human skin grafted onto the chorioallantois of the chick embryo for virus cultivation *Proc Soc Exper Biol & Med* 69:341 1948
- Brun R T A relation between the viruses of zoster and varicella *Brit J Exp Path* 14:67 1933
- Brunsgaard E The mutual relation between zoster and varicella *Brit J Dermat* 44:1 1932
- Crouse H V Coriell L L Blank H and Scott T F M Cytochemical studies on the intranuclear inclusion of herpes simplex *J Immunol* 65:119 1950
- Denny Brown D and Adams R D Pathologic features of herpes zoster *Arch Neur & Psych* 51:216 1944
- Downie A W and Dumbell K R The isolation and cultivation of variola virus on the chorioallantois of the chick embryo *J Path & Bact* 59:189 1947
- Ebert M H Histologic changes in sensory nerves of the skin in herpes zoster *Arch Dermat & Syph* 60:641 1949
- Frank L Varicella pneumoniae *Arch Path* 50:450 1950
- Goodpasture E W and Anderson K Infection of the human skin grafted on the chorioallantois of chick embryos with virus of herpes zoster *Am J Path* 20:447 1944
- Head H and Campbell A W Pathology of herpes zoster and its bearing on sensory location *Brain* 23:353 1900
- Johnson H N Visceral lesions associated with varicella *Arch Path* 30:292 1940

## Metabolic Diseases

## LIPOIDOSES

The term lipoidoses has been applied to a group of diseases in which the lesions, due to a local or generalized disturbance of the lipid metabolism, contain lipid substances.

No generally accepted classification of the lipoidoses exists. The following classification is based largely on the classifications published by Thannhauser and by Montgomery and Osterberg.

- I Lipoidoses with Increased Blood Lipids
  - 1 Primary hypercholesteremic xanthomatosis
  - 2 Biliary xanthomatosis
  - 3 Idiopathic hyperlipemia
  - 4 Secondary hyperlipemia
- II Lipoidoses with Normal Blood Lipids
  - 5 Lipid reticulo-endotheliosis (Hand Schuller Christian disease)
    - a Fulminating type Letterer-Siwe disease
    - b Remittent type Letterer-Siwe disease
  - 6 Lipoid proteinosis
  - 9 Extracellular cholesterosis
- III Localized lipoidoses
  - 10 Xanthelasma palpebrarum
  - 11 Necrobiosis lipoidica

## LIPOIDOSES WITH INCREASED BLOOD LIPIDS

## 1 Primary Hypercholesteremic Xanthomatosis

Primary

clear. *...xanthomatata* therefore the serum is

cutaneous lesions consist of flat or slightly raised xanthelasmata on the eyelids and tuberous xanthomata, especially on the elbows and the knees. In addition, there are tendon xanthomata, especially on the Achilles tendons, the patellar tendon, and the extensor ten

Sheldon W H and Heyman A Lymphogranuloma venereum Am J Path.  
23 653 1947

Milkers Nodules

Katzenellenbogen I Studies on milkers nodules Dermatolog 103 69 1959

Nasemann T and Deubner B Beitrag zur Virusatologie des Melkerknotens  
Hautarzt 4 210 1953

Nomland R and McKee A P Milkers nodules Arch Dermat & Syph 63 663  
1952

in xanthoma cells to be anisotropic (doubly refractile) in contrast with the fat in the sebaceous cells and the subcutaneous fat cells, which is isotropic (not doubly refractile) (See page 31)

Xanthoma cells form from perithelial cells, which are histiocytes. They have usually one nucleus but may have two and even many



FIG 126 Xanthoma tuberosum early lesion. Scarlet red stain for fat. The xanthoma cells are filled with lipid material (X100)

nuclei. In multinucleated xanthoma cells the nuclei either are irregularly distributed as in foreign body giant cells, or lie near the center of the cell grouped around a small island of nonfoamy cytoplasm and surrounded by foamy cytoplasm. This latter

is an assortment of inflammatory cells, particularly polymorphonuclear leukocytes, lymphocytes and histiocytes. In well-developed lesions the infiltrate is composed almost entirely of xanthoma cells (Fig 125). Fat stains in this stage show



dons of the hands Atherosclerotic cardiovascular disease is common and may lead to death due to coronary occlusion in early life (Bloom Kaufman and Stevens)

**Histopathology.** The histologic appearance of the cutaneous and the tendinous lesions is characterized by the presence of xanthoma or foam cells Xanthoma cells are phagocytic cells filled with lipid drop



FIG 125 Xanthoma tuberosum, early lesion Numerous xanthoma cells (foam cells) are present There is only little fibrosis ( $\times 200$ )

lets In routine sections, the lipid droplets have been dissolved and extracted in the process of fixation and embedding so that the cells have a reticulated or foamy appearance (Fig 125) However, the lipid droplets can be seen when formalin fixed frozen sections are stained with fat stains, such as scarlet red (Fig 126) With scarlet red, the lipid substance in the xanthoma cell stains a brownish red, in contrast to the lipid substance in sebaceous glands and the subcutaneous fat, which stains a bright orange red This is due to the fact that the lipid substance in xanthoma cells is predominantly cholesterol and phospholipids, whereas the lipid substance in the sebaceous cells and the subcutaneous fat cells is predominantly neutral fat

✓Polariscopic examination of frozen sections reveals the lipid droplets

xanthomatous lesions often by many years. The blood serum is clear but due to its high content of bilirubin it is intensely green.

The biliary cirrhosis which causes the xanthomatous lesions is produced as a rule either by cholangiolitis or by extrahepatic obstruction (MacMahon) but occasionally in children by congenital hypoplasia of bile ducts (MacMahon and Thannhauser).

**Histopathology** The cutaneous lesions show the same histologic picture as is found in primary xanthomatosis.

### 3 Idiopathic Hyperlipemia

In idiopathic hyperlipemia which usually is not familial the blood serum shows an elevation not only of the cholesterol and phospholipids as in primary hypercholesteremic xanthomatosis but also of the neutral fat. Therefore the blood serum is milky.

The cutaneous lesions are of two types: tuberos xanthomata which are found especially on the elbows and the knees and papular xanthomata which may be diffusely distributed but are most prevalent on the buttocks. The papular xanthomata have a tendency to come and go and therefore are often referred to as eruptive xanthomata. Xanthelasmata of the eyelids are absent but tendon xanthomata may occur (Lever, Smith and Hurley).

Visceral manifestations occasionally are present. They include hepatosplenomegaly, attacks of abdominal pain due to secondary pancreatitis and coronary heart disease which however occurs less frequently than in primary hypercholesteremic xanthomatosis. The association of idiopathic hyperlipemia with hepatosplenomegaly occasionally is referred to as Burger-Gruetz disease.


**Histopathology** The histologic appearance of the xanthomata in idiopathic hyperlipemia is the same as that of the xanthomata in primary hypercholesteremic xanthomatosis. The papular xanthomata which usually are of recent origin often show like most young xanthomata an admixture of inflammatory cells (see page 239).

Only few reports on the histology of the visceral lesions are available. In one patient with hepatosplenomegaly, conspicuous fat infiltration of the liver cells was observed on biopsy (Movitt, Gerstl, Sherwood and Epstein). The presence of pancreatitis in patients with attacks of abdominal pain has been confirmed repeatedly by exploratory operation (Klatskin and Gordon).

### 4 Secondary Hyperlipemia

Secondary hyperlipemia may occur secondary to severe diabetes (xanthoma diabeticorum), to nephrosis and to glycogen storage disease (von Gierke's disease). The amounts of cholesterol phospho

that all fat is intracellular. In involuting lesions, fibroblasts appear (Fig 127). Ultimately, fibrosis replaces the foam cells. Weidman and Schaffer express the belief that the foam cells themselves can become transformed into fibroblasts. In old fibrosing lesions cholesterol may be found not only in foam cells but also extracellularly.



center of the cell grouped around a small island of nonfoamy cytoplasm and surrounded by foamy cytoplasm ( $\times 400$ )

**Differential Diagnosis.** Differentiation between a fibrotic lesion of xanthoma tuberosum and histiocytoma may be very difficult and even impossible. It may be necessary to rely upon clinical and blood chemical data for the correct diagnosis (Montgomery and Osterberg).

## 2 Biliary Xanthomatosis

Biliary cirrhosis may cause high values for cholesterol and phospholipids in the serum and lead to xanthelasmata on the eyelids and tuberous xanthomata on the skin that are indistinguishable from those of primary hypercholesteremic xanthomatosis. However, deep jaundice is present and the jaundice precedes the appearance of

xanthomatous lesions often by many years. The blood serum is clear but due to its high content of bilirubin it is intensely green.

The biliary cirrhosis which causes the xanthomatous lesions is produced as a rule either by cholangiolitis or by extrahepatic obstruction (MacMahon) but occasionally in children by congenital hypoplasia of bile ducts (MacMahon and Thannhauser).

**Histopathology** The cutaneous lesions show the same histologic picture as is found in primary xanthomatosis.

### 3 Idiopathic Hyperlipemia

In idiopathic hyperlipemia which usually is not familial the blood serum shows an elevation not only of the cholesterol and phospholipids as in primary hypercholesteremic xanthomatosis but also of the neutral fat. Therefore the blood serum is milky.

The cutaneous lesions are of two types: tuberous xanthomata which are found especially on the elbows and the knees and papular xanthomata which may be diffusely distributed but are most prevalent on the buttocks. The papular xanthomata have a tendency to

visceral manifestations occasionally are present. They include hepatosplenomegaly, attacks of abdominal pain due to secondary pancreatitis and coronary heart disease which however occurs less frequently. The association occasionally is

**Histopathology** The histologic appearance of the xanthomata in at of the xanthomata in The papular xanthomata show like most young xanthomata an admixture of inflammatory cells (see page 259).

Only few reports on the histology of the visceral lesions are available. In one patient with hepatosplenomegaly conspicuous fat infiltration of the liver cells was observed on biopsy (Movitt, Gerstl, Sherwood and Epstein). The presence of pancreatitis in patients with attacks of abdominal pain has been confirmed repeatedly by exploratory operation (Klaumin and Gordon).

### 4 Secondary Hyperlipemia

Secondary hyperlipemia may occur secondary to severe diabetes (xanthoma diabetorum) to nephrosis and to glycogen storage disease (von Gierke's disease). The amounts of cholesterol phospho-

lipids and neutral fat in the serum are elevated and the serum is milky.

Papular eruptive xanthomata occur which usually are diffusely distributed but have a predilection for the buttocks. Occasionally, recently erupted papules have an inflammatory base.

**Histopathology.** The papular xanthomata of secondary hyperlipemia show the same histologic picture as other xanthomata. Since they usually are of recent origin, a rather marked inflammatory infiltrate may be present with large numbers of neutrophils. In the evolving stage, much of the cholesterol may be seen in phagocytes and extracellularly (Montgomery and Osterberg).

### LIPIDOSIS WITH NORMAL BLOOD LIPIDS

#### 5 Lipid Reticulo endotheliosis (Hand Schuller Christian Disease)

Letterer Siwe disease, Hand Schuller Christian disease and eosinophilic granuloma represent variations in degree, stages of development and localization of the same basic disease process (Farber). Formerly, Hand Schuller Christian disease was regarded as a primary disturbance of the lipid metabolism, as a normocholesteremic xanthomatosis (Thannhauser and Mgendantz), but this concept has been abandoned (Mallory) and the presence of cholesterol in the lesions is now regarded as a secondary infiltration. The three diseases are at present believed to be histiocytoses or reticulo endothelioses. If the reticulo endotheliosis occurs in infancy, it is generalized and rapidly fatal (Letterer Siwe disease), death occurs before sufficient time has elapsed for the development of the lesion into xanthogranuloma. In early childhood, the disease is chronic (Hand Schuller Christian disease), and lipidization is as a rule pronounced. In later childhood or in the adult the usual picture is that of eosinophilic granuloma which represents an abortive form of the disease. Transitional cases between these three forms of reticulo endotheliosis are common. The values for blood plasma lipids are normal in all three forms.

**LETTERER SIWE DISEASE (NONLIPID HISTIOCYTOSIS OF FOOT AND OLCOTT)** This disorder usually occurs in infants and is almost inevitably fatal within a few months. It is characterized by, fever, anemia, enlargement of the liver and the spleen, lymphadenopathy and multiple defects of the bones. In most cases cutaneous lesions are present. They may consist of petechiae, papules or pustules. In some cases one observes numerous closely set, brownish papules covered with scales or crusts. This type of eruption usually is extensive with a predilection to involve the scalp, the face and the trunk.

The resemblance of the eruption to seborrheic dermatitis and Darier's disease is often striking (Laymon and Sevenants)

**Histopathology** The cutaneous lesions show usually close to the epidermis and often invading into the epidermis (Fig. 128) accumulations of histiocytes (reticulum cells) intermingled with a few lymphocytes and varying numbers of eosinophils. Extravasated blood cells frequently lie in and about the masses of histiocytes. The histiocytes appear as large cells with irregularly shaped vesicular nuclei and abundant slightly eosinophilic cytoplasm (Fig. 129). In some areas these cells are distinctly outlined and even separated by edema but in other areas their cytoplasm is confluent. Occasionally some of the cells have a foamy cytoplasm and stain positive for fat with fat stains. The epidermis may become destroyed by pressure of the underlying cells (Foot and Olcott, Abt and Denenholz, Lane and Smith, case 1).

The visceral lesions consist of proliferations of large pale reticulo-endothelial cells which invade and replace the normal structure of spleen, liver, bone marrow, lymph nodes and other organs (Sweitzer and Laymon).

**HAND-SCHÜLLER-CHRISTIAN DISEASE** Diabetes insipidus, exophthalmos and multiple defects of the bones, especially of the cranium represent the triad of typical Hand-Schüller-Christian disease. However, any one or even all three of the cardinal symptoms may be absent and involvement may occur in entirely different organs. For example, enlargement of the liver, the spleen and the lymph nodes is common and dwarfism is observed occasionally. Hand-Schüller-Christian disease takes a chronic course usually extending over years and has a mortality of about 70 per cent.

Cutaneous lesions are quite uncommon. If present they are similar to those seen in Letterer-Siwe disease, consisting of an extensive eruption of coalescing scaly or crusted papules with a distinct clinical resemblance to Darier's disease. Occasionally the term xanthoma disseminatum is used for this eruption.

**Histopathology** Early cutaneous lesions of Hand-Schüller-Christian disease show the same histologic picture as the cutaneous lesions of Letterer-Siwe disease (Lane and Smith, Laymon and Sevenants). In mature lesions a large number of the histiocytes show a foamy cytoplasm and typical foam cells may be present (Thannhauser and Vigdantz). In old lesions the number of foam cells again may be small and foreign body giant cells may be found. As a rule the tendency to lipidization is less pronounced in the cutaneous lesions than in the lesions of other organs.



FIG 128 Hand Schuller Christian disease (eosinophilic granuloma) Low magnification The upper portion of the dermis contains an infiltrate composed almost entirely of loosely aggregated histiocytes The infiltrate has invaded the epidermis in many areas ( $\times 100$ )




FIG 129 Hand Schuller Christian disease (eosinophilic granuloma) High magnification The upper portion of the dermis contains an infiltrate composed almost entirely of loosely aggregated histiocytes The infiltrate has invaded the epidermis in many areas ( $\times 400$ )

loma) High  
shaped nu  
invaded the

Eosinophilic  
ease of this gr

They occur involved. The cutaneous lesions consist either of an extensive eruption of crusted papules as in Letterer Siwe and Hand Schuller Christian disease (Lever and Leeper) or of one or several erythematous granulomatous plaques which may undergo ulceration (Curtis and Cawley McCreary). The two types of lesions may be present simultaneously. (Granuloma faciale formerly called eosinophilic granuloma of the face is an entirely different disease from eosinophilic granuloma and is not related to it. For its description see page 111.)

Histopathology. The papules have the same histologic appearance as in Letterer Siwe disease, namely an infiltrate of loosely aggregated large cells which is and invading the epidermis. The number of eosino-

The granulomatous plaques also show numerous loosely arranged large histiocytes within an edematous stroma. Eosinophils are present in varying numbers. They usually lie in patches rather than diffusely distributed through the lesion. In addition a few lymphocytes and plasma cells are present (Curtis and Cawley McCreary, Lever and Leeper).

XLipid is absent in the cutaneous lesions but may be present in the associated osseous lesions.

#### 6 Niemann Pick Disease

Niemann Pick disease is characterized by abnormal deposits of sphingomyelin and diamorphosphatide in the reticulo endothelial cells of many organs but not in the skin. The level of lipids in the blood serum is normal.

The disease occurs as a rule in Jewish infants and is fatal. There are enlargement of the liver and the spleen, cachexia and brownish discoloration of the skin.

Histopathology. On histologic examination the brownish discoloration of the skin is found to be due to the presence of increased amounts of melanin.

#### 7 Gaucher's Disease

In Gaucher's disease ceroid, a cerebroside is deposited in the reticulo endothelial cells of many organs. The skin however is spared. The blood lipids are normal.

The disease tends to be familial, occurs predominantly in Jews, may start at any age and takes a chronic course. There is hepatosple-



nomegaly and rarefaction with cortical thickening of the long bones. The skin shows brownish discoloration.

**Histopathology** The skin shows increased amounts of melanin.

### ♀ Lipoid Proteinosis (Urbach Wiethe)

Lipoid proteinosis is characterized by lipid infiltrations of the skin of the oral mucosa and of the larynx. Although the blood lipids may




FIG. 130 Lipoid proteinosis. The dermis is occupied by thick wavy hyaline-like bundles which run perpendicular to the epidermis. In addition the hyaline-like material surrounds all blood vessels as a thick mantle. (X200)

be normal (Wise and Rein, Price, LaRosa and Settle) there may be an increase in the total lipids (Hansen, Ramos e Silva) or a relative increase in phospholipids (Montgomery and Havens, Wile and Snow). The disease is often familial.

Clinically, one observes nodular and verrucous lesions on the skin and on the mucous membranes of the mouth and the larynx. The nodules of the skin on regressing leave pitted scars giving the skin a

There is hoarse-

is striking and  
irregular acan



... Lipoid proteinosis. Scarlet red stain for fat. A large amount of lipid material is present. It occurs in the form of small droplets throughout the hyaline material, particularly around the blood vessels ( $\times 100$ )

thosis. The dermis is considerably thickened. The upper half of the dermis is occupied by thick, wavy bundles which are

... and of the vascular endothelium are well preserved. In the lower dermis, the collagen appears normal but the

homogeneous material is present around some of the vessels and sweat glands

On staining with scarlet red, a large amount of lipid material is visualized. It stains a rusty brown and occurs in the form of small droplets throughout the hyalin like material, but particularly around the blood vessels (Fig 131). It is located extracellularly. On polariscopic examination, the lipid material proves to be isotropic, not doubly refractile. Urbach and Wiethe interpreted the histologic findings as indicating a merging of lipid and protein and, therefore, suggested the name lipoid proteinosis.

No definite conclusion has so far been reached about the nature of the lipid substance in the dermis. In several cases, the chemical analysis of involved skin revealed a decided increase in the amount of lecithin, which is a monoaminophosphatide (Montgomery and Havens, Ramos e Silva). In other cases, however, no such increase was found (Wile and Snow, Hansen, Price, LaRosa and Settle).

### 9 Extracellular Cholesterosis (Urbach)

This disorder is characterized by extracellular cholesterol infiltrations in the skin.

Only three cases of this condition have been reported in the literature (Urbach, Epstein and Lorenz, Laymon, Sobel and Pollock). All three cases showed reddish brown verrucous plaques and nodules involving especially the dorsa of the hands and the feet and the extensor surfaces of the legs. Ulceration of some of the lesions occurred in Sobel's case. The amount of cholesterol in the blood/serum was normal in Laymon's case and subnormal in Urbach's and Sobel's cases.

**Histopathology.** The involved skin shows a dense, nonspecific cellular infiltrate composed mainly of histiocytes and lymphocytes. The blood vessels are dilated and their endothelium is swollen. No foam cells are present. In healing lesions there is considerable fibrosis.

✓ Fat stains reveal numerous droplets of fat in extracellular position throughout the lesions but especially about the blood vessels. On polariscopic examination the fat droplets are doubly refractile.

Chemical examination of tissue in Urbach's and Sobel's cases revealed the content of cholesterol three to five times greater in the lesions than in the normal skin.

The cause of the disease is unknown. Sobel and Pollock, impressed with the subnormal values for cholesterol in their as well as in Urbach's case, suggest that, for some unknown reason, the blood is unable to hold normal amounts of cholesterol which, therefore, is

deposited in the tissue. Being unable to metabolize the cholesterol the tissue responds not with foam cell formation but instead with a severe inflammatory reaction as if dealing with a foreign body of irritating or toxic nature.

## LOCALIZED LIPOIDOSES

### 10 Xanthelasma Palpebrarum

This disorder is characterized by the presence of soft yellowish plaques on the eyelids caused by the deposition of cholesterol. Although xanthelasma palpebrarum is common in primary hypercholesteremic xanthomatosis (see page 257) it frequently occurs in individuals with little or no elevation of the serum cholesterol (Epstein, Rosenman and Gofman). Since in such instances the deposition of cholesterol probably is caused by local degenerative changes in the skin of the eyelids, xanthelasma palpebrarum is best regarded as a localized lipoidosis.

**Histopathology** The histologic changes are similar to those of primary xanthomatosis. As a rule, however, fewer Touton cells are seen and they may be absent. Fibrotic changes may occur (Montgomery and Osterberg).

### 11 Necrobiosis Lipoidica

This disease represents a localized lipoidosis inasmuch as the deposition of lipids occurs in areas in which degeneration or necrobiosis of collagen has taken place. The necrobiosis of collagen is due to vascular changes (Roederer, Wotinger and Burgun). In those cases in which diabetes exists, it can be assumed that the diabetes has caused the vascular changes. However, diabetes originally thought to be a prerequisite for this disease is present in only about one third of the cases (Kaalund-Jørgensen). In the other two thirds the cause of the vascular changes is unknown.

**Clinically** one observes on the legs and rarely elsewhere on the skin one or several sharply demarcated, irregularly outlined, glazed patches which are yellow in the center and violaceous at the periphery. The center gradually becomes depressed and atrophic and may break down to form an ulcer.

**Histopathology** On histologic examination the epidermis —

In these areas the collagen appears homogeneous swollen and partly basophilic (Fig. 132). The collagen bundles often are broken up and,

homogeneous material is present around some of the vessels and sweat glands

On staining with scarlet red a large amount of lipid material is visualized. It stains a rusty brown and occurs in the form of small droplets throughout the hyalin like material but particularly around the blood vessels (Fig. 131). It is located extracellularly. On polariscopic examination the lipid material proves to be isotropic, not doubly refractile. Urbach and Wiethe interpreted the histologic findings as indicating a merging of lipid and protein and therefore suggested the name lipoid proteinosis.

No definite conclusion has so far been reached about the nature of the lipid substance in the dermis. In several cases the chemical analysis of involved skin revealed a decided increase in the amount of lecithin which is a monaminophosphatide (Montgomery and Havens, Ramos e Silva). In other cases however no such increase was found (Wile and Snow, Hansen, Price, LaRosa and Settle).

### 9 Intracellular Cholesterosis (Urbach)

This disorder is characterized by extracellular cholesterol infiltrations in the skin.

Only three cases of this condition have been reported in the literature (Urbach, Epstein and Lorenz, Lyndon, Sobel and Pollock). All three cases showed reddish brown verrucous plaques and nodules involving especially the dorsa of the hands and the feet and the extensor surfaces of the legs. Ulceration of some of the lesions occurred in Sobel's case. The amount of cholesterol in the blood/serum is as normal in Lyndon's case and subnormal in Urbach's and Sobel's cases.

**Histopathology.** The involved skin shows a dense nonspecific cellular infiltrate composed mainly of histiocytes and lymphocytes. The blood vessels are dilated and their endothelium is swollen. No norm cells are present. In healing lesions there is considerable fibrosis.

At stains reveal numerous droplets of fat in extracellular position throughout the lesions but especially about the blood vessels. On polariscopic examination the fat droplets are doubly refractile.

Chemical examination of tissue in Urbach's and Sobel's cases revealed the content of cholesterol three to five times greater in the lesions than in the normal skin.

The cause of the disease is unknown. Sobel and Pollock, impressed with the subnormal values for cholesterol in their as well as in Urbach's case, suggest that for some unknown reason the blood is unable to hold normal amounts of cholesterol which therefore is

degeneration. In contrast with the neutral fat in the subcutaneous layer and the sebaceous glands, which stains a brilliant orange red, the granules stain a rusty brown. According to Hildebrand,  $\frac{105}{101}$   $\frac{105}{101}$   $\frac{105}{101}$




FIG. 133 *Necrobiosis lipoidica*. Several foreign body giant cells are located within an area of collagen degeneration. Two fibrous vessels are present. ( $\times 200$ )

(Laymon and Fisher) The fact that occasionally, lipids are absent in the lesions (Sachs) indicates that their presence is purely a secondary phenomenon.

*Granulomatosis disciformis chronica progressiva* recently described by several European authors (Miescher and Leder, Arzt) probably represents *necrobiosis lipoidica* without lipid deposits.

**Differential Diagnosis.** In the differential diagnosis, *granuloma annulare* must be considered, because both *necrobiosis lipoidica* and *granuloma annulare* show areas of collagen degeneration and the same type of reactive inflammation and fibrosis. However in *granuloma annulare* there are no vascular changes, no deposits of lipids, few or no giant cells and no atrophy or ulceration of the epidermis.

instead of lying parallel to the surface of the skin extend in various directions. There often is evidence of formation of young collagen between the degenerated bundles.

Within and near the areas of necrobiosis often extending into the subcutaneous fat one finds a predominantly perivascular inflammatory infiltrate composed of lymphocytes, histiocytes, fibroblasts and




FIG. 132 Necrobiosis lipoidica. Much of the collagen appears degenerated. An inflammatory infiltrate is scattered through the areas of degeneration. A vessel in the center shows endothelial proliferation and fibrosis of its wall ( $\times 100$ ).

occasional groups of epithelioid cells. Foreign body giant cells commonly are present and thus are of considerable diagnostic value (Michelson and Layman; Belote and Welton) (Fig. 133). Occasionally a few foam cells are noted (Kluber, Nichols).

The blood vessels, particularly in the middle and the lower dermis, exhibit fibrosis of their walls with proliferation of their endothelial lining. This process may lead to partial and occasionally even to complete occlusion of the lumen. Thrombosis of small vessels occurs sometimes. These vascular changes account for the degeneration of the collagen.

Staining for fat with scarlet red frequently, but not always, reveals numerous granules of lipid extracellularly in the areas of collagen



FIG 131 Primary systemic amyloidosis Round amorphous fissured masses of amyloid are present in the uppermost dermis. They resemble those of colloid milium but in contrast to colloid milium are present throughout the dermis ( $\times 900$ )




FIG 132 Primary systemic amyloidosis Subcutaneous fat. Amyloid is deposited in the walls of two blood vessels and also around fat cells forming so-called amyloid rings ( $\times 400$ )



on the other hand, deposits of mucin usually are present. In spite of these differences, Ellis and Kirby Smith regard the two diseases as closely related.

### AMYLOIDOSIS

Three forms of amyloidosis exist: (1) primary systemic amyloidosis, (2) primary localized amyloidosis of the skin (lichenoid amyloidosis) and (3) secondary systemic amyloidosis. In primary systemic amyloidosis, the skin is frequently involved, in secondary systemic amyloidosis, very rarely.

The Congo red test is of great value in the diagnosis of cutaneous amyloidosis. If 1.0 cc. of a 1 per cent solution of Congo red is injected subcutaneously, or 0.1 cc. intradermally, into the affected region, the areas in which amyloid is deposited will be stained strongly with red dye after from 24 to 48 hours, whereas the interposed skin will appear only very slightly stained (Nomland, Dostrovsky and Sigler). In histologic sections stained with hematoxylin and eosin, amyloid appears as light pink, probably because

of the process of fixation. With the Van Gieson stain, amyloid stains yellow and collagen red. Amyloid is a protein polysaccharide complex and therefore it stains at least in lichenoid amyloidosis and in secondary amyloidosis red with Congo red, purple red with methyl violet (indicating metachromasia) and deeply red with the periodic acid Schiff reaction. In primary systemic amyloidosis, however, these staining reactions are not always present and vary from patient to patient and even from organ to organ, probably because of differences in the amount of polysaccharides or in the nature of the union between the protein and the carbohydrate molecules (Goltz).

### PRIMARY SYSTEMIC AMYLOIDOSIS

Mesenchymal tissue is affected while the parenchyma of parenchymatous organs is spared. Amyloid deposits are found mainly in the smooth and striated musculature in the small blood vessels and in the gastrointestinal tract. Macroglossia has been present in almost one half, and cutaneous lesions in about one fourth, of the 57 cases reported up to 1949 (Dahlin).

Of interest is the frequency with which primary systemic amyloidosis is associated with multiple myeloma and Bence Jones proteinuria (Brunsting and MacDonald). Examination of the urine for Bence Jones protein, roentgenograms of the bones and sternal biopsy in search of atypical plasma cells in the bone marrow, therefore, should be performed in all cases of primary systemic amyloidosis.

about capillaries (Nomland) In some cases a mild chronic inflammatory infiltrate will be found

**Differential Diagnosis** Lichenoid amyloidosis must be differentiated from colloid milium For differential diagnosis see page 165

### SECONDARY SYSTEMIC AMYLOIDOSIS

This condition occurs in chronic suppurative diseases, such as tuberculosis and in chronic cachectic diseases associated with marked loss of protein from the body Amyloid deposits are found in the parenchymatous organs especially in the liver, the kidney, the spleen (sago spleen) and the adrenals The skin is involved very rarely Michelson and Lynch have reported a case of secondary systemic amyloidosis due to tuberculosis in which there were diffuse nodular lesions of the lips due to deposits of amyloid

### CALCINOSIS CUTIS

Two forms of calcinosis cutis exist metastatic and metabolic calcification

#### METASTATIC CALCIFICATION

Metastatic calcification develops as a result of hypercalcemia The hypercalcemia may be due to parathyroid neoplasm hypervitaminosis D chronic renal disease or destruction of bone in such conditions as osteomyelitis and malignant growths (Mulligan) Calcium is apt to be deposited in organs in which the cells excrete acid and therefore have a low carbon dioxide tension As a result of the low carbon dioxide tension

cells of these organs excrete acid substances The skin and the subcutaneous tissue are affected only rarely However instances of metastatic calcification of either skin or subcutaneous tissue have been reported as caused by parathyroid neoplasm (Penecke Laubmann) by hypervitaminosis D (Bevens and Taylor) by chronic renal disease (Platt and Owen) and by osteomyelitis (Weidman and Shaffer)

**Histopathology** Calcium deposits are recognized easily in histologic sections since they stain deeply blue with hematoxylin and eosin and black with von Kossa's stain for calcium In most instances of metastatic calcification the calcium occurs as individual granules as well as massive deposits in the dermis and in the subcutaneous fat (Fig 136) Larger deposits often evoke a foreign body reaction so that giant cells an inflammatory infiltrate and fibrosis may be present around them In Weidman and Shaffer's case the calcium deposits consisted of small granules which were found not only in the

Clinically, the skin shows discrete and coalescing papules and nodules which are semitranslucent and of a waxy, amber color. In addition, yellowish plaques resembling those of xanthomatosis are present. Petechiae and ecchymoses frequently occur at the site of eruption. Itching is absent. The face is predominantly affected especially the periorbital regions.

**Histopathology.** Histologic examination of the skin reveals large faintly eosinophilic, amorphous masses of amyloid which may be deposited anywhere in the dermis (Fig. 134) as well as in the subcutaneous tissue. In addition, small deposits of amyloid may occur in the membrana propria of the sweat glands around blood vessels and in the walls of blood vessels. In most cases, no inflammatory reaction is present (Michelson and Lynch), but, in some cases, one finds foci of lymphocytes, plasma cells and foreign body giant cells (Pearson, Rice and Dickens).

In the subcutaneous tissue, one may find besides large masses of amyloid and deposits of amyloid in the walls of the blood vessels, so called amyloid wings which are formed by the deposition of amyloid around individual fat cells (Fig. 135) (Pearson, Rice and Dickens). The fat cells thus appear as if cemented together by the eosinophilic amyloid substance (Iverson and Morrison).

Not only in the skin and the subcutaneous tissue, but also throughout the body, the small arteries and the veins show amyloid deposits often entirely replacing their media and adventitia. The vessels of the tongue, the skeletal muscle, the respiratory tract, the heart and the gastro intestinal and the genito urinary tracts usually are affected most severely. In the skeletal muscle and in the tongue, numerous muscle fibers show amyloid deposits in the form of amorphous nodular swellings (Iverson and Morrison).

#### PRIMARY LOCALIZED AMYLOIDOSIS OF THE SKIN (LICHENOID AMYLOIDOSIS)

In this form of amyloidosis only the skin is involved. The lesions are seen most commonly on the legs but they may occur elsewhere. They consist of closely set discrete conical or flat, brownish red papules which resemble the papules of lichen planus. Occasionally the papules have a translucent appearance. In some instances the papules, by coalescence form plaques which may develop a verrucous surface and then resemble lichen simplex chronicus. The lesions usually itch severely in contrast with those of primary systemic amyloidosis.

**Histopathology.** The amyloid deposits are much smaller in size than those found in primary systemic amyloidosis and are limited to the subepidermal region of the dermis. The earliest deposits occur

In sclerodermic calcinosis, the histologic appearance of the skin and the subcutaneous tissue is that of scleroderma. The calcium deposits are usually located within areas of sclerotic collagen. The deposition of calcium probably is due to the decreased metabolic activity in the sclerotic tissue. Decreased metabolic activity causes the carbon dioxide tension in the tissue to be lower than normal and thus reduces the solubility of calcium (Brody and Bellin).

In patients with idiopathic calcinosis cutis no perceptible histologic changes may precede the deposition of the calcium (Epstein, Bauer, Marble and Bennett). Bauer, Marble and Bennett found, in their case, that the initial lesion consisted of deposition of finely divided particles of calcium salts around apparently normal fat cells in the subcutaneous tissue. The granules seemed to coalesce slowly to form large masses. Other authors found that, even in idiopathic calcinosis, mild degenerative changes preceded the deposition of calcium (Rothstein and Welt, Atkinson and Weber).

## GOUT

Gout is a disturbance of purine metabolism characterized by uricemia. Deposits of urates are found in the tissues of the joints particularly their cartilages in the cartilage of the ears and in the



FIG. 137 Gout Deposits of sodium urate are surrounded by a foreign body giant cell reaction. On the left the sodium urate is present as needle shaped crystals. ( $\times 100$ )

dermis but also in the epidermis, in the sweat glands, in the sweat ducts and in the nerve trunks

### METABOLIC CALCIFICATION

Metabolic calcification is due to local metabolic disturbances and is not associated with hypercalcemia. Deposits of calcium occur, as a rule, only in the skin and the subcutaneous tissue, but occasionally




FIG. 136 Metastatic calcification as a result of hypercalcemia (produced by prolonged and excessive administration of vitamin D). Von Kossa stain for calcium. Irregular masses of calcium surrounded by a foreign body giant cell reaction are present in the subcutaneous fat ( $\times 100$ ).

also in the muscles and the tendons. The internal organs are spared. There may be just a few deposits (calcinosis circumscripta) or innumerable deposits everywhere in the skin (calcinosis universalis). Between 30 and 40 per cent of the cases of calcinosis circumscripta and universalis described in the literature occurred in patients with scleroderma or dermatomyositis (Atkinson and Weber) (See "Scleroderma," page 310, and "Dermatomyositis," page 303). In the rest, no reason for the calcinosis was evident.

**Histopathology.** As in metastatic calcification, the calcium may be present as individual granules or as massive deposits in the dermis and in the subcutaneous fat. In the subcutaneous fat, extensive areas of calcification may be observed. A foreign body giant cell reaction is often found around the larger deposits of calcium.

In the delayed cutaneous form the bullae arise as pressure bullae (see page 66) subepidermally (Zeligman and Baum Robert) However due to regeneration of the epidermis older bullae may be located partially or entirely within the epidermis The bullae, thus, do not differ histologically from those observed in epidermolysis bullosa The milia like those of epidermolysis bullosa dystrophica consist of small intradermally located epidermal cysts (Robert)

## MYXEDEMA

Three types of myxedema occur generalized myxedema circumscribed myxedema and papular myxedema Generalized myxedema is a manifestation of hypothyroidism Circumscribed myxedema is associated with or preceded by hyperthyroidism and occurs almost invariably together with exophthalmos commonly it follows thyroidectomy or therapy with thiouracil Papular myxedema is not associated with any disturbance of thyroid function

The mucin present in the tissue in these three diseases appears light blue on staining with hematoxylin and eosin It stains red with the periodic acid Schiff reaction and is strongly metachromatic with methylene blue thionine cresyl violet and toluidene blue indicating that it is a protein polysaccharide complex (Brewer) The mucin is digested in sections by hyaluronidase which points to the fact that it contains a large amount of hyaluronic acid (Palitz and Brunner) It also stains red with muckarmine (The latter staining method requires fixation with absolute alcohol—see page 29)

## GENERALIZED MYXEDEMA

Clinically the entire skin appears swollen dry pale and waxy It feels firm to the touch In spite of its edematous appearance the skin does not pit on pressure The facies is characteristic the nose is broad and thick and the lips are swollen

**Histopathology** The dermis is increased in thickness The collagen bundles as well as the individual fibers of the collagen bundles are separated by edema The collagenous fibers show signs of degener-

ation. The mucin is usually small (Reuter)

## CIRCUMSCRIBED MYXEDEMA

The lesions usually are limited to the anterior aspects of the legs They consist of hard raised nodular yellow waxy plaques with prominent hair follicles

subcutaneous tissue. The deposits in the ears and in the skin manifest themselves as nodules of varying size. These nodules are called tophi.

**Histopathology.** The tophi show needle shaped crystals of sodium biurate lying closely packed in the form of bundles or shewes. The crystals often have a brownish color. They stain well with von Kossa's stain. The accumulations of urates are surrounded by granulation tissue containing many foreign body giant cells (Fig. 137).

## PORPHYRIA

Porphyria represents an inborn metabolic error in which large amounts of uroporphyrin and coproporphyrin are excreted in the urine. Three forms of porphyria exist: the congenital form, the acute intermittent form and the delayed cutaneous form. In the congenital form, the porphyrins are formed in the bone marrow (porphyria erythropoietica) while in the acute intermittent form and in the delayed cutaneous form they are formed in the liver (porphyria hepatica) (Watson).

In the congenital form, cutaneous lesions identical with those seen in hydroa vacciniforme appear from earliest childhood on the exposed portions of the skin following exposure to the sun. The lesions result in scarring and mutilation. The teeth may appear red.

The acute intermittent form starts in adult life and is characterized by attacks of abdominal pain, peripheral neuropathy and mental disturbances. Cutaneous lesions do not occur, as a rule. However, in rare instances, cutaneous lesions like those in the delayed cutaneous form have been observed (Nesbitt and Watkins).

The delayed cutaneous form (porphyria cutanea tarda) like the acute intermittent form, remains latent until adult life. Frequently chronic alcoholism by causing hepatic dysfunction precipitates the onset of clinical manifestations. Blisters form on exposure to light as well as on minor mechanical or thermal trauma. They often heal with scarring and formation of milia. Abdominal pain and nervous system manifestations as in the acute intermittent type may be present, although to a lesser degree (Brunsting and Mason). Because of the occurrence of blisters following trauma and the presence of milia, some authors have referred to the lesions as acquired epidermolysis bullosa. However, the excretion of uroporphyrin and coproporphyrin in the urine and the presence of sensitivity to light make it evident that the disease is not related to epidermolysis bullosa (Brunsting and Mason).

**Histopathology.** In the congenital form the vesicles have the same appearance as in hydroa vacciniforme (see page 46).

In the delayed cutaneous form the bullae arise as pressure bullae (see page 66) subepidermally (Zeligman and Baum Robert). However, due to regeneration of the epidermis older bullae may be located partially or entirely within the epidermis. The bullae thus do not differ histologically from those observed in epidermolysis bullosa. The milia like those of epidermolysis bullosa dystrophica consist of small intradermally located epidermal cysts (Robert).

## MIXEDEMA

Three types of myxedema occur: generalized myxedema, circumscribed myxedema, and papular myxedema. Generalized myxedema is a manifestation of hypothyroidism. Circumscribed myxedema is associated with or preceded by hyperthyroidism and occurs almost invariably together with exophthalmos; commonly it follows thyroidectomy or therapy with thiouracil. Papular myxedema is not associated with any disturbance of thyroid function.

The mucin present in the tissue in these three diseases appears light blue on staining with hematoxylin and eosin. It stains red with the periodic acid-Schiff reaction.

methylenel

that it is a

digested in sections by hyaluronidase which points to the fact that it contains a large amount of hyaluronic acid (Palitz and Brunner). It also stains red with mucicarmine. (The latter staining method requires fixation with absolute alcohol—see page 29.)

## GENERALIZED MIXEDEMA

Clinically, the entire skin appears swollen, dry, pale, and waxy. It feels firm to the touch. In spite of its edematous appearance, the skin does not pit on pressure. The facies is characteristic: the nose is broad and thick, and the lips are swollen.

Histopathology: The dermis is increased in thickness. The collagen bundles as well as the individual fibers of the collagen bundles are separated by edema. The collagenous fibers show some of the

## CIRCUMSCRIBED MIXEDEMA

The lesions usually are limited to the anterior aspects of the legs. They consist of hard, raised, nodular, yellow, waxy plaques with prominent hair follicles.



subcutaneous tissue. The deposits in the ears and in the skin manifest themselves as nodules of varying size. These nodules are called tophi.

**Histopathology.** The tophi show needle shaped crystals of sodium biurate lying closely packed in the form of bundles or sheaves. The crystals often have a brownish color. They stain well with von Kossa's stain. The accumulations of urates are surrounded by granulation tissue containing many foreign body giant cells (Fig. 137).

## PORPHYRIA

Porphyria represents an inborn metabolic error in which large amounts of uroporphyrin and coproporphyrin are excreted in the urine. Three forms of porphyria exist: the congenital form, the acute intermittent form and the delayed cutaneous form. In the congenital form, the porphyrins are formed in the bone marrow (porphyria erythropoietica) while in the acute intermittent form and in the delayed cutaneous form they are formed in the liver (porphyria hepatica) (Watson).

In the congenital form cutaneous lesions identical with those seen in hydrot vacciniforme appear from earliest childhood on the exposed portions of the skin following exposure to the sun. The lesions result in scarring and mutilation. The teeth may appear red.

The acute intermittent form starts in adult life and is characterized by attacks of abdominal pain, peripheral neuropathy and mental disturbances. Cutaneous lesions do not occur, as a rule. However, in rare instances, cutaneous lesions like those in the delayed cutaneous form have been observed (Nesbitt and Watkins).

The delayed cutaneous form (porphyria cutanea tarda) like the acute intermittent form, remains latent until adult life. Frequently chronic alcoholism by causing hepatic dysfunction precipitates the onset of clinical manifestations. Blisters form on exposure to light as well as on minor mechanical or thermal trauma. They often heal with scarring and formation of milia. Abdominal pain and nervous system manifestations as in the acute intermittent type may be present, although to a lesser degree (Brunsting and Mason). Because of the occurrence of blisters following trauma and the presence of milia some authors have referred to the lesions as acquired epidermolysis bullosa. However, the excretion of uroporphyrin and coproporphyrin in the urine and the presence of sensitivity to light make it evident that the disease is not related to epidermolysis bullosa (Brunsting and Mason).

**Histopathology.** In the congenital form the vesicles have the same appearance as in hydrot vacciniforme (see page 46).

### PAPULAR MYXEDEMA (PAPULAR MUCINOSIS)

There is a widespread eruption of asymptomatic soft, yellowish papules which may coalesce into irregular patches.

**Histopathology** The appearance of the mucinous infiltrate is like that in circumscribed myxedema inasmuch as there are rather large amounts of mucin as well as stellate cells (Dalton and Seidell). However the extent of the mucinous infiltrate is more limited than in circumscribed myxedema because it is present only in the upper dermis and in the case of individual papules within relatively small areas.

### SCLEREDEMA ADULTORUM (BUSCHKE)

Scleredema adultorum is characterized by diffuse edema and induration of the skin and the subcutaneous tissue. Its cause is unknown but it is noteworthy that it frequently follows an infectious disease such as grippe or tonsillitis. It usually begins on the face and spreads rapidly to involve the neck and the upper trunk. Complete resolution takes place in a few months. The occurrence of pleural and pericardial effusions and of hydrarthrosis has been reported (Vallee).

**Histopathology** Throughout the dermis one observes swelling and splitting up of the collagen bundles by edema. The edema may be severe enough to produce in the dermis clear unstained spaces of various sizes (fenestration). Freund noted that the edema substance which did not stain with hematoxylin and eosin stained metachromatically with cresyl violet like mucin but not with mucicarmine. Vallee found this staining reaction in one case but not in another. Braun-Falco observed purplish metachromasia with toluidine blue which was no longer present when the sections prior to the staining were incubated with hyaluronidase. He concluded that the edema substance consisted of hyaluronic acid.

### ADDISON'S DISEASE

Addison's disease which is caused by hypofunction of the adrenal glands is characterized by weakness, loss of weight, low blood pressure and diffuse hyperpigmentation of the skin and the mucous membranes.

The hyperpigmentation in Addison's disease

It is known that in Addison's disease the damaged adrenal gland responds but weakly to pituitary stimulation and

**Histopathology.** Large amounts of mucin are present in the dermis particularly in the lower dermis. There it occurs not only as individual threads and granules, but also as massive deposits causing wide separation of the collagenous fibers (Fig. 138). The number of




FIG. 138. Circumscribed myxedema. Considerable amounts of mucin are present especially in the lower dermis separating the collagen bundles as well as individual collagen fibers. (The empty spaces are due to shrinkage and filling out of mucin) ( $\times 200$ )

fibroblasts is increased and newly formed collagen is present. Some fibroblasts have a stellate shape and are surrounded by mucin. Because of the much larger amount of mucin and the new formation of collagen, the dermis is greatly thickened, much more than in generalized myxedema.

The epidermis shows hyperkeratosis. The rete ridges are often flattened. The vessels in the upper dermis are dilated and surrounded by a mild inflammatory infiltrate. Elastic tissue stains show the elastic tissue to be frayed and greatly decreased (O'Leary).

**Differential Diagnosis** Differentiation of acanthosis nigricans from nevus verrucosus may be impossible (Curth). As a rule however nevus verrucosus shows more marked acanthosis than acanthosis nigricans and overdevelopment rather than atrophy of the rete ridges



FIG. 139 Acanthosis nigricans. There are hyperkeratosis and papillomatosis. Several papillae project up and as finger like projections. As is usually the case acanthosis and hyperpigmentation are slight ( $\times 100$ )

### RIEHL'S MELANOSIS MELANODERMATITIS TOXICA POIKILODERMA RETICULARE (CIVATTE)

Riehl's melanosis and melanodermatitis toxica have the same appearance clinically as well as histologically (Storck). In both conditions one observes ill defined symmetrical bluish brown hyperpigmentation of the face and occasionally also of the neck and the chest. There may be slight atrophy of the skin and follicular hyperkeratosis. The cause of Riehl's dermatosis is not fully known but it has been suggested that it is caused by a lack of vitamin B and provoked by exposure to the sun and that thus it is related to pellagra (Pechlmann). Melanodermatitis toxica is due to contact of the affected skin with tars oils or greases.

Poikiloderma reticulare differs from these two diseases clinically by its predominant localization on the neck, the reticular arrangement of the hyperpigmentation and the presence of telangiectases (Pierini and Bosq). Histologically the telangiectases usually are not sufficiently evident to differentiate poikiloderma reticulare from the other two diseases.

consequently, as a compensatory phenomenon, the pituitary gland is overactive.

**Histopathology.** Histologic examination often shows no changes other than hyperpigmentation. Occasionally, one observes slight flattening of the rete ridges and slight thinning of the epidermis. The amount of melanin is increased in both the epidermis and the dermis. In the epidermis, the melanin is present chiefly in the basal layer, but may be found also in the lower layers of the stratum malpighii. Because the pigment formation proceeds slowly over a long period of time, the number of clear cells is increased only slightly. In the upper dermis, a moderate number of melanin laden chromatophores (melanophores) are present. There is no inflammatory reaction in the dermis.

**Differential Diagnosis.** A diagnosis of Addison's disease cannot be made from histologic sections, because the same histologic picture is observed in nonspecific hyperpigmentation of the skin and in the normal skin of the Negro.

### ACANTHOSIS NIGRICANS

Three types of acanthosis nigricans exist: malignant acanthosis nigricans, benign acanthosis nigricans and pseudo acanthosis nigricans. Clinically and histologically, the three forms look alike (Curtis).

The malignant form occurs in adults and is associated with internal cancer, usually of the glandular type. The benign or juvenile form may start at any time before puberty. It represents a genodermatosis related to nevus verrucosus. Pseudo acanthosis nigricans occurs in the body creases of obese, brunette persons. It disappears when the patient loses weight.

Clinically, all three forms of acanthosis nigricans present verrucous, hyperpigmented patches, predominantly in the axillae, on the neck and in the submammary and the genital regions.

**Histopathology.** Histologic examination reveals marked hyperkeratosis and papillomatosis. In addition, acanthosis and hyperpigmentation are present but are slight (Fig. 139). Thus, the name acanthosis nigricans has little histologic justification.

In a typical lesion, the papillae project far upward as finger like projections and are covered with a not unduly thickened stratum malpighii. The valleys between the finger like projections are filled in largely by keratin. In areas where there is no papillomatosis the epidermis shows areas of moderate acanthosis with adjacent areas of atrophy of the stratum malpighii. The rete ridges, as a rule, are developed only poorly. There may be a slight increase in the amount of melanin in the basal layer.

## VITILIGO

Vitiligo is characterized by variously sized and shaped sharply demarcated patches of depigmentation surrounded by hyperpigmented skin

**Histopathology** In the depigmented areas the epidermis is devoid of melanocytes. The amount of melanin in the basal layer is increased and numerous melanophores are present in the upper dermis

The dopa reaction is negative in the basal layer of the depigmented areas and positive in the basal layer of the adjoining hyperpigmented areas

## HEMOCHROMATOSIS (BRONZE DIABETES)

This disturbance of the metabolism is characterized by deposition of hemosiderin in various organs of the body. The presence of hemosiderin causes bronzing of the skin, cirrhosis of the liver and sclerosis of the spleen and the pancreas. Sclerosis of the pancreas is responsible for the diabetes usually associated with the disease.

Clinically the pigmentation of the skin is diffuse and is indistinguishable from that seen in Addison's disease.

**Histopathology** Granules of hemosiderin are found within chromatophores in the upper dermis and in the membrana propria of the sweat glands. Occasionally they may also occur in the basal cells of the epidermis. Hemosiderin is best demonstrated by staining sections with potassium ferrocyanide. With this method hemosiderin granules on account of their content of iron stain blue. Not infrequently melanin is present in excess in the basal layer and within chromatophores.

In selecting a site for biopsy it is not necessary to choose a pigmented area because hemosiderin is present throughout the skin in hemosiderosis. However it is important not to take a specimen from the legs where deposits of hemosiderin frequently occur in association with stasis dermatitis and other vascular disturbances (Montgomery and O'Leary).

**Differential Diagnosis** It is impossible to differentiate the granules of hemosiderin from those of melanin in routine stains. They can however be easily differentiated by staining with potassium ferrocyanide which stains hemosiderin blue but does not stain melanin. Furthermore melanin does not occur in the membrana propria of the sweat glands.

**Histopathology.** The clinically visible hyperpigmentation in these three diseases is brought about by pigmentary incontinence of the basal layer, resulting in an accumulation of melanin in the upper dermis. The epidermis shows mild hyperkeratosis, thinning of the stratum malpighii and varying degrees of degeneration of the cells in the basal layer. The amount of melanin in the basal layer is decreased. The papillary and the subpapillary layers of the dermis how-




FIG. 110 Melanoderma. The basal layer shows numerous clear cells. The upper dermis shows large amounts of melanin, mainly within but also outside of chromatophores. A mild perivascular inflammatory infiltrate is present ( $\times 200$ ).

ever, show large amounts of melanin, mainly within, but also outside of, chromatophores (Fig. 110). In addition, there are various degrees of inflammatory reaction in the upper dermis. In some cases, the inflammatory infiltrate is limited to the perivascular areas, in others, it is extensive, bandlike and close to the epidermis and thus resembles that of lichen planus (Storck).

At a later stage, the degeneration of the basal layer is slight and many clear cells may be present. The degree of inflammation lessens with the age of the lesion.

**Differential Diagnosis.** Incontinentia pigmenti differs from the three diseases just discussed by showing no inflammatory infiltrate in the upper dermis. Addison's disease shows no inflammatory infiltrate either and only a slight amount of melanin in the dermis.

there may be considerable regeneration of the cutaneous appendages (Steffens Bair and Sheard)

**Differential Diagnosis** Differentiation of phrynoderma from ichthyosis may be difficult. As a rule, however, ichthyosis shows atrophy or absence of the granular layer, thinning of the epidermis and elongation and branching of the rete ridges. Pityriasis rubra pilaris differs from phrynoderma by showing, in addition to hyperkeratosis and follicular plugging, spotted parakeratosis, irregular acanthosis and a more pronounced inflammatory infiltrate, not limited to the hair follicles.

### PELLAGRA

Pellagra is caused by a deficiency of the vitamin B complex, but particularly of nicotinic acid. Besides cutaneous lesions, pellagra usually presents also a stomatitis, which is characterized by edema and redness of the tongue and atrophy of the lingual papillae. In addition, gastro-intestinal symptoms and nervous and mental changes may be present.

Cutaneous lesions occur predominantly on exposed areas, such as the dorsa of the hands, the wrists, the face, the V area of the upper chest and the dorsa of the feet. In the early stage, there is erythema, which in severe cases may be accompanied by bullae. Later, the erythema assumes a livid shade and the skin becomes thickened and scaling. Ultimately, the affected areas become atrophic and deeply pigmented.

**Histopathology** Early lesions present a chronic inflammatory infiltrate with moderate edema in the upper dermis. Vesicles and bullae may be present. They may be located intra-epidermally as well as subepidermally.

Older lesions show hyperkeratosis with areas of parakeratosis and a moderate degree of acanthosis. Follicular plugging occasionally is observed. The amount of melanin in the epidermis is increased. The

deeper portions of the dermis (Moore, Spies and Cooper).

In the end stage, hyperkeratosis and hyperpigmentation are still present, but the stratum malpighii now shows considerable atrophy with flattening of the rete ridges. The dermis shows moderate fibrosis.

**Differential Diagnosis** The histologic picture of pellagra is not diagnostic. As a rule, it is merely one of chronic dermatitis. In the end stage, the presence of hyperpigmentation and of atrophy of the stratum malpighii serves to distinguish pellagra from chronic dermatitis.



The granules of silver, present in argyria, are also frequently located in the membrana propria of the sweat glands, but they differ from those of hemosiderin by being much smaller and more uniform in size. Furthermore, they do not stain with potassium ferrocyanide and are refractile with dark field illumination (see page 153).

### OCHRONOSIS

In ochronosis, due to an inborn metabolic error, the catabolism of tyrosine cannot proceed beyond homogentisic acid. The disease is characterized by blackening of the cartilages and by osteoarthritis and may show bluish discoloration of the sclerae and brown or bluish mottled pigmentation of the skin. The urine darkens on exposure to the air. The discoloration of the cartilages, the sclerae and the skin as well as the darkening of the urine, are due to the presence of homogentisic acid which by oxidation is converted into a dark-colored insoluble product.

**Histopathology.** Varying amounts of a light brown pigment are present in the dermis. The pigment is either diffusely distributed or present as clumps of varying size and shape. The clumps may be large, measuring more than 100 microns in diameter. The pigment does not stain with silver nitrate as melanin does but becomes black when stained with polychrome methylene blue (Layman).

### VITAMIN A DEFICIENCY (PHYRYNODERMA)

Vitamin A plays an important part in the metabolism of the epithelial structures of the skin and the mucous membranes. In addition to cutaneous changes, deficiency of vitamin A may cause night blindness, xerophthalmia and keratomalacia.

The cutaneous changes, to which the name phrynoderma has been given, consist of dryness and roughness of the skin and the presence of follicular hyperkeratosis.

Lichen spinulosus of Crocker is probably identical with phrynoderma (Lehman and Rapaport).

**Histopathology.** The skin shows moderate hyperkeratosis with marked distention of the upper part of the hair follicles by large horny plugs. In contrast with ichthyosis the granular layer is present and may even be increased in thickness. The lower part of the hair follicles is atrophic and usually surrounded by a slight, chronic inflammatory infiltrate. There are only few remnants of sebaceous glands. In addition, one may find evidence of atrophy of the sweat glands, such as flattening of the secretory cells (Frazier and Hu). In severe cases the sweat glands and the sebaceous glands may undergo keratinizing metaplasia (Bessey and Wolbach). When adequate amounts of vitamin A are supplied to a patient with phrynoderma,

- Kaallund Jørgensen O. Necrobiosis lipoidica (diabeticorum) Acta dermat venereol 28 214 1948
- Klaber R. Necrobiosis lipoidica diabeticorum Report of a case Brit J Dermat 46 926 1954
- 1939
- Laymon C W. Extracellular cholesterosis Arch Dermat & Syph 35 269 1937
- and Fisher I. Necrobiosis lipoidica (diabeticorum ?) Arch Dermat & Syph 59 150 1949
- and Sevenants J J. Systemic reticuloendothelial granuloma Arch Dermat & Syph 57 873 1948
- Lever W F and Leeper R W. Eosinophilic granuloma of the skin Arch Dermat & Syph 62 85 1950
- Smith P A J and Hurley N A. Idiopathic hyperlipemia and primary hypercholesteremic xanthomatosis I Clinical data and analysis of the plasma lipids J Invest Dermat 22 35 1954
- McCreary J H. Eosinophilic granuloma Arch Dermat & Syph 58 572 1948
- MacMahon H E. Biliary xanthomatosis (xanthomatous biliary cirrhosis) Am J Path 21 527 1948
- and Thannhauser S J. Biliary xanthomatosis (congenital acholangic biliary cirrhosis) due to hypoplasia or partial agenesis of bile ducts Am J Path 27 750 1951
- Mallory T B. Pathology diseases of bone New England J Med 227 955 1942
- Michelson H E and Laymon C W. Necrobiosis lipoidica diabeticorum Arch Dermat & Syph 35 1130 1937
- Miescher G and Leder M. Granulomatosis disciformis chronica et progressiva Dermatolog 97 25 1948
- Montgomery H and Havens F Z. Xanthomatosis IV I no 1 1950
- 
- Munt E R, Gerstl B, Sherwood F and Epstein C C. Essential hyperlipemia Arch Int Med 87 79 1951
- Nicholas L. Necrobiosis lipoidica diabeticorum with xanthoma cells Arch Dermat & Syph 48 606 1943
- Price H, LaRosa W V and Seale E B. Lipoidosis cutis et mucosae Arch Dermat & Syph 55 42 1947
- Ram s e Silva J. Lipid protemosis (Urbach Wiethe) Arch Dermat & Syph 47 301 1945
- Roederer J, Woringer F and Burgun R. Considérations sur un cas de nécrobiose lipoidique Dermatolog 99 131 1949
- Sachs P. in discussion of Rein C R and Kanof N B. Necrobiosis lipoidica, Arch Dermat & Syph 61 130 1950
- Schubert
- Sue
- The
- Th. — — — J. Lipidoses ed. 2 New York Oxford Univ Press, 1950

## VITAMIN C DEFICIENCY (SCURVY)

Scurvy, caused by a deficiency of ascorbic acid, is characterized by bleeding and spongy gums and petechial hemorrhages which often are perifollicular.

**Histopathology.** Scurvy is characterized by inability of the supporting tissues to produce and maintain intercellular substances (Wolbach and Howe). Thus, the endothelial cells of capillaries fail to form adequate amounts of intercellular ground substance. Extravasation of red blood cells results. The extravasation occurs without inflammatory changes around the capillaries. Scurvy, consequently, belongs to the group of noninflammatory purpuras (see page 127) (Peck, Rosenthal and Erf).

Histologic examination of the skin shows hemorrhages around the capillaries. As a rule, these hemorrhages are most pronounced in the vicinity of the hair follicles. In many instances one finds hyperkeratosis and follicular plugging as in vitamin A deficiency. According to Scheer and Keil, these changes in the epidermis are purely secondary.

## BIBLIOGRAPHY

## Lipoidoses

- Abt A F and Denenholz E J Letterer-Siwe's disease. *Am J Dis Child* 51:499 1936
- Arzt L Zur Differentialdiagnose granulomatöser Prozesse (Granulomatosis disciformis chronica progressiva, Necrobiosis lipoidica diabetorum atypisches The Granulom). *Hautarzt* 3:488 1952
- Belote G H and Welton D G Necrobiosis without diabetes. *Arch Dermat & Syph* 40:887 1939
- Bloom D Kaufman S R and Stevens R A Hereditary xanthomatosis. *Arch Dermat & Syph* 45:1, 1942
- Bürger M and Grütz O Ueber hepatosplenomegalie Lipoidose mit xanthomatösen Veränderungen in Haut und Schleimhaut. *Arch f Dermat u Syph* 166:542, 1932
- Curtis A C and Cawley E P Eosinophilic granuloma of bone with cutaneous manifestations. *Arch Dermat & Syph* 55:810 1947
- Ellis F A and Kirby Smith H Necrobiosis lipoidica and granuloma annulare. *Arch Dermat & Syph* 45:40 1942
- Epstein N N Rosenman R H and Golman J W Serum lipoproteins and cholesterol metabolism in xanthelasma. *Arch Dermat & Syph* 65:70 1952
- Farber S The nature of solitary or eosinophilic granuloma of bone. *Am J Path* 17:625 1941
- Foot N C, and Olcott C T Report of a case of non lipoid histiocytosis (reticuloendotheliosis) with autopsy. *Am J Path* 10:81 1934
- Hand, A Defects of membranous bones, exophthalmos and polyuria in childhood: is it dyspituitarism? *Am J M Sc* 162:509 1921
- Hansen P Ein Fall von Lipoidproteinose. *Arch f Dermat u Syph* 175:618

- Rothstein J L and Welt S Calcinosis universalis and calcinosis circumscripta in infancy and in childhood Am J Dis Child 52:368 1936
- Weidman F D and Shaffer L W Calcification of the skin including the epiderm in connection with extensive bone resorption Arch Dermat & Syph 14:503 1926

## Goat

- Brogsitter A M Histopathologie der Gelenkgicht Deutsch Arch f klin Med 153:257 1926 154:1 1927

## Porphyria

- Brunsting L V and Mason H L Porphyria with cutaneous manifestations Arch Dermat & Syph 60:66 1949
- Nesbitt S and Watkins C H Acute porphyria Am J Med Sc. 203:74 1942
- Robert P Epidermolysis bullosa dystrophica atige Dermatose bei kongenitaler Iorphunurie Dermatolog 97:106 1948
- Watson C J Some recent studies of porphyrin metabolism and porphyria Lancet 1:539 1951
- Zeligman I and Baum M Porphyric bullous dermatosis Arch Dermat & Syph 58:337 1948

## Myxedema

- Brewer O B Myxoedema an autopsy report with histochemical observations on the nature of the mucoid infiltrations J Path & Bact 63:503 1951
- Dalton J E and Seidell M A Studies on lichen myxedematosus (papular mucinosis) Arch Dermat & Syph 67:194 1953
- O'Leary P A Localized solid edema of the extremities in association with exophthalmic goiter Arch Dermat & Syph 21:57 1950
- Pink L L and Brunner M J The mucinoses A classification with histochemical studies on the nature of mucin J Invest Dermat 11:159 1950
- Libbury D M and Stokes J H Circumscribed myxedema of the skin Arch Dermat & Syph 21:255 1931
- Reuter M J Histopathology of the skin in myxedema Arch Dermat & Syph 21:35 1931

## Scleredema Adultorum

- Braunfalko O Neues zur Histopathologie des Scleroedema adultorum (Buschke) Dermat Wchnschr 125:409 1952
- Freund H Leber Sklerodem (Buschke) Arch f Dermat u Syph 161:92 1930
- Sweitzer S E and Lajmon C W Scleredema adultorum (Buschke) Arch Dermat & Syph 37:420 1938
- Valley B L Scleredema a systemic disease New England J Med 235:207 1946

## Addison's Disease

- Hall T C, Mcracken B H and Thorn G W Skin pigmentation in relation to adrenal cortical function J Clin End & Met 13:243 1953
- Montgomery H and O'Leary P A Pigmentation of the skin in Addison's disease acanthosis nigricans and hemochromatosis Arch Dermat & Syph 21:570 1950

## Acanthosis Nigricans

- Curtis H O Benign type of acanthosis nigricans Arch Dermat & Syph. 34:353 1936
- Significance of acanthosis nigricans Arch Dermat & Syph 66:60 1952
- Herold W C, Kaufman W H and Smith D C Acanthosis nigricans, Arch Dermat & Syph 44:789 1941

- Thannhauser S J and Magendintz H The different clinical groups of xanthomatous diseases a clinical physiological study of 22 cases Am J Int Med 11 1662 1938 (Good review of the lipoidoses)
- Urbach E Epstein E and Lorenz K Extrazelluläre Cholesterinose Arch f Dermat u Syph 166 213 1932
- and Wiethe C Lipoidosis cutis et mucosae Virch Arch f path Anat 273 285 1929
- Weidman F D and Schaffer H W Xanthoma of the skin and larynx Arch Dermat & Syph 35 767 1937
- Wile U J and Snow D S Lipoid proteinosis Arch Dermat & Syph 43 131 1941
- Wise F and Rein C R Lipoidosis cutis et mucosae (lipoid proteinosis of Urbach) Arch Dermat & Syph 37 201 1938

### Amyloidosis

- Brunsting I A and MacDonald J D Primary systemic amyloidosis with microglossia a syndrome related to Bence Jones proteinuria and myeloma J Invest Dermat 8 145 1917
- Dahlin D C Primary amyloidosis with report of six cases Am J Path 23 105 1919
- Dostrovsky A and Sigher F Localized amyloidosis of the skin Arch Dermat & Syph 41 891 1911
- Goltz R W Systematized amyloidosis Medicine 31 81 1932
- Iverson L and Morrison A B Primary systemic amyloidosis Arch Path 45 1 1918
- Michelson H E and Lynch F W Systematized amyloidosis of the skin and muscles Arch Dermat & Syph 29 805 1931 32 363 1935
- Nomland R Localized (lichen) amyloidosis of the skin report of two cases with vital staining of the amyloid nodules by Congo red injected intracutaneously or subcutaneously Arch Dermat & Syph 33 85 1936
- Pearson B Rice M M and Dickens K L V Primary systemic amyloidosis Arch Path 32 1 1941

### Calcinosis Cutis

- Atkinson F R B and Weber F P Cutaneous and subcutaneous calcinosis Brit J Dermat 50 267 1938
- Bauer W Marble A and Bennett G A Further studies in a case of calcification of subcutaneous tissue (calcinosis universalis) in a child Am J M Sc 182 237 1931
- Bevans M and Taylor H K Lesions following use of Ertron in rheumatoid arthritis Am J Path 23 367 1917
- Brody J and Bellin D E Calcinosis with scleroderma Arch Dermat & Syph 36 85 1937
- Epstein E Idiopathic calcinosis cutis Arch Dermat & Syph 34 367 1936
- Laubmann W Hochgradige Kalkmetastasierung bei Epithelkörperchentumoren Verhandl d dtisch pathol Gesellsch 27 229 1934
- Mulligan R M Metastatic calcification Arch Path 43 177 1917 (Good review)
- Penecke Ueber zwei Fälle von Ostitis fibrosa Recklinghausen mit Epithelkörperchentumoren Centrbl f allg Pathol u path Anat 37 535 1926
- Platt R and Owen T K Renal dwarfism associated with calcification of arteries and skin Lancet 2 135 1934

# Systemic Diseases of Unknown Cause

## LUPUS ERYTHEMATOSUS

Three types of lupus erythematosus are generally recognized (1) chronic discoid lupus erythematosus (2) subacute disseminate lupus erythematosus and (3) acute systemic lupus erythematosus. In the second type visceral lesions may be absent. Inter

mediary forms between the three types occur clinically as well as histologically. They cannot be regarded as separate diseases.

**Clinical Appearance** Chronic discoid lupus erythematosus is limited usually to the face, where the flush areas of the cheeks are affected predominantly. The scalp and the ears also may be involved. The lesions consist of well defined erythematous slightly infiltrated patches showing adherent keratotic scaling and follicular plugging. Older lesions show in addition atrophic scarring. Systemic symptoms are absent.

Subacute disseminate lupus erythematosus usually is superimposed upon chronic discoid lupus erythematosus but may begin as such. In addition to the face other areas especially the thorax and the arms and the legs are involved. Systemic symptoms such as fever, malaise and leukopenia are often present. The lesions consist of erythematous often slightly raised patches which tend to coalesce and show only slight scaling. The prognosis as to life is fairly good since development into the final acute form of the disease is not common.

Acute systemic lupus erythematosus has severe systemic symptoms and almost invariably fatal. Even corticotropin and cortisone as a rule merely delay death because the renal lesions do not respond to these drugs. Systemic symptoms often precede the cutaneous eruption and occasionally cutaneous manifestations are absent throughout the course of the disease. The cutaneous eruption frequently begins on the face as a diffuse ill-defined erythema with some edema.

## 292 Metabolic Diseases

Knowles F C Sidlick D M and Ludy J B Acanthosis nigricans Arch Dermat & Syph 19 391 1929

### Riehl's Melanosis Melanodermatitis Toxica Poikiloderma Reticulare (Cavatte)

Loerster H R and Schwartz I Occupational melanosis from pitch Arch Dermat & Syph 39 955 1939

Pierini L E and Bosq P Maladie de Cavatte Ann de dermat et de syph 9 381 1938

Pochlmann A Ueber Melanosen der Gesichtshaut in der Kriegs und Nachkriegszeit Dermat Wchnschr 119 454 1947

Storck H Ueber Riehl'sche Melanose Dermatolog 92 246 1946

### Vitiligo

Becker S W Vitiligo Arch Dermat & Syph 28 497 1933

Oliver E A Schwartz I and Warren I H Occupational leukoderma Arch Dermat & Syph 42 993 1940

### Hemochromatosis

Montgomery H and O'Leary P A Pigmentation of the skin in Addison's disease acanthosis nigricans and hemochromatosis Arch Dermat & Syph 21 970 1930

### Ochronosis

Laymon C W Ochronosis Arch Dermat & Syph 67 553 1953

### Vitamin A Deficiency

Bessey O A and Wolbach S B Vitamin A physiology and pathology J A M A 110 2072 1938

Irazier C N and Hu C Nature and distribution according to age of cutaneous manifestations of vitamin A deficiency Arch Dermat & Syph 33 855 1936

Lehman E and Rapaport H G Cutaneous manifestations of vitamin A deficiency in children J A M A 114 386 1940

Steffens L F Bur H I and Sheard C Dark adaptation and dietary deficiency in vitamin A Am J Ophth 23 1325 1940

### Pellagra

Moore R A Spies T D and Cooper Z K Histopathology of the skin in pellagra Arch Dermat & Syph 46 100 1942

### Vitamin C Deficiency

Dilldorf G The pathology of vitamin C deficiency J A M A 111 1376 1938

Peck S M Rosenthal N and Erf L Purpura Arch Dermat & Syph 35 831 1937

Scheer M and Keil H Follicular lesions in vitamin A and C deficiencies Arch Dermat & Syph 30 177 1934

Wolbach S B and Howe P R Intercellular substances in experimental scorbutus Arch Path 11 1946

fore, the hyperkeratosis may not be present until after the lesion is several weeks old. Parakeratosis is usually completely absent. The keratotic plugs are found mainly in the follicular openings but occur also in the sweat ducts and independent of either. The follicles inside the dermis may contain concentric layers of keratin instead of hairs (Fig 141). The atrophy of the stratum malpighii is not always uni-




FIG. 141. There are degenerative basophilic areas of collagen and a patchy, predominantly lymphocytic infiltrate ( $\times 200$ ).

form but may alternate with areas of anarthrosis. Focal liquefaction degeneration of the basal layer represents the most significant histologic change in lupus erythematosus (Fig 142). In its absence, a histologic diagnosis of lupus erythematosus is rarely, if ever, justified. In older lesions the basal layer often is partially absent as a consequence of this degeneration.

The dermis shows considerable edema in its upper portion. In darkly skinned persons melanin is often present in the upper dermis since the degeneration of the cells in the basal layer causes them to lose their melanin (pigmentary incontinence). The capillaries and larger vessels are dilated and their walls may show edema, however,



("erythema perstans"). Soon other lesions appear, often widely spread over the body. They are poorly defined, purplish and often edematous, and may be purpuric or vesicular. The systemic symptoms include irregular fever, malaise, weakness, pains in muscles and joints and pleural pains. Laboratory findings include marked leukopenia, ↑ hypergammaglobulinemia, proteinuria and often, hematuria



FIG 141 Chronic discoid lupus erythematosus. Low magnification. There is keratotic plugging and the follicles inside the dermis contain, instead of hairs, concentric layers of keratin. The epidermis is atrophic and devoid of rete ridges. The inflammatory infiltrate is distinctly patchy and tends to be located in the vicinity of hair follicles (×50).

in most mbina in the skin. (1) hyperkeratosis with keratotic plugging, (2) atrophy of the stratum malpighii, (3) liquefaction degeneration of the basal cells, (4) a patchy, perivascular, cellular inflammatory infiltrate with tendency to arrangement about follicles degeneration of the collagen (Fig. 141). However, not all these changes are present in every case.

The epidermal changes are secondary to those in the dermis. There-

degeneration of the collagen (see under Acute Systemic Lupus Erythematosus)

1. Section of Acute Systemic Lupus Erythematosus The fibrinoid degeneration is observed also in the skin. In order to study fibrinoid degeneration of the col-



FIG. 143. Acute systemic lupus erythematosus. The epidermis is atrophic and shows marked liquefaction degeneration of the basal layer. The dermis shows fibrinoid degeneration. Only a mild perivascular inflammatory infiltrate is present. ( $\times 200$ )

lagen in the skin sections should be taken from covered areas of the body; in exposed areas basophilic degeneration of the collagen is

lygen fibers

usually not vis-

phibic refract

thickened st.

... more deeply eosinophilic than nor-  
mally. In more advanced lesions the degenerated collagen fibers be-  
come fragmented and fuse with the altered ground substance. The  
fibrinoid material shows metachromasia upon staining with toluidine  
blue and stains positive with the periodic acid-Schiff reaction.

proliferative or obliterative changes are absent. The inflammatory infiltrate is distinctly patchy. It is located mainly in the vicinity of the hair follicles and the sebaceous glands, presses upon these structures and causes their gradual atrophy and disappearance (Fig 141). The infiltrate is composed predominantly of lymphocytes but contains also a small number of plasma cells and histiocytes.


Basophilic degeneration of the collagen in the upper dermis is common in sections of discoid lupus erythematosus obtained from exposed areas (where discoid lupus erythematosus usually occurs); it is absent, however, in sections taken from covered areas (Montgomery). In spite of its common occurrence in lupus erythematosus of exposed areas, basophilic degeneration of the collagen cannot be regarded in any way as diagnostic of lupus erythematosus because it occurs also in simple senile atrophy of the exposed skin (see page 157) as well as in many other dermatoses when located in exposed areas. The elastic tissue shows, irrespective of whether the section is obtained from an exposed or a nonexposed area, at first fraying and later destruction throughout the dermis wherever the inflammatory infiltrate occurs.

In the differential diagnosis, two diseases have to be considered which share with chronic discoid lupus erythematosus the presence of a patchy infiltrate—namely, secondary syphilis and lymphocytic lymphoma. Secondary syphilis differs from chronic discoid lupus erythematosus by the arrangement of the patchy infiltrate predominantly around blood vessels by the presence of numerous plasma cells in the infiltrate and by the presence of vascular changes. In lymphocytic lymphoma the patchy infiltrate is composed entirely of lymphocytes and does not show a tendency to arrangement near the epidermal appendages. Neither secondary syphilis nor lymphocytic lymphoma show epidermal changes comparable with those of chronic discoid lupus erythematosus.

**Histopathology of Subacute Disseminate Lupus Erythematosus**  
The histologic changes in the lesions differ only in degree from those of chronic discoid lupus erythematosus. The atrophy of the epidermis, the liquefaction degeneration of the basal cells and the edema of the dermis are more prominent than in chronic discoid lupus erythematosus whereas the hyperkeratosis and the inflammatory infiltrate are less marked.

Occasionally, the edema in the upper dermis and the liquefaction degeneration of the basal cells is severe enough to result in the formation of clefts and even vesicles between the epidermis and the dermis (McCreight and Montgomery). There may be evidence of fibrinoid

fibrillary ground substance normally not visible may be present as homogeneous eosinophilic fibrinoid clumps (Fig 143) (Klemperer Pollack and Baehr 1941) There is a rather mild perivascular infiltrate. In addition one sees diffusely scattered histiocytes and fibroblasts some of which show pyknosis of their nuclei. Extravasations of red blood cells are frequently observed. Changes in the walls of the vessels other than edema are usually absent. Only oc-



generation ( $\times 100$ )

asionally does one see ~~degenerative~~ changes in the walls. They are best demonstrated by the use of the periodic acid Schiff reaction (Stoughton and Wells).

The subcutaneous fat is often involved. It may show focal mucoid degeneration with reactive lymphocytic infiltration. The collagen bundles separating the fat lobules may be increased in thickness and show edema and fibrinoid degeneration similar to those in the dermis (Fig 145).

**Histopathology of the Visceral Lesions of Acute Systemic Lupus Erythematosus.** Visceral lesions usually are widespread but often minute in size so that they may easily be overlooked on gross inspection and even on histologic examination unless they are especially looked for. The endocardium, the serous membranes, the heart and the skeletal muscle, the renal glomeruli, the spleen, the lymph nodes and the fat depots are affected most commonly.

Altshuler and Angevine believe that the fibrinoid material is formed by the precipitation of the acid mucopolysaccharides of the ground substance by an alkaline protein derived from either the necrosis of tissue or the interaction of the tissue with a damaging agent. The fibrinoid degeneration may affect the collagenous ground substance of capillaries, arterioles and venules. Thus vascular damage if pres-




FIG. 144. Acute systemic lupus erythematosus. There are marked liquefaction degeneration of the basal layer and edema of the upper dermis. Within the zone of subepidermal edema the collagen is present as homogeneous fibrinoid material which lies in irregular conglomerates and also surrounds a capillary ( $\times 400$ ).

ent does not represent an independent lesion but is part of the basic injury to the collagen (Klemperer, Pollack and Biehr, 1941).

The histologic appearance of the skin in acute systemic lupus erythematosus resembles that of subacute disseminate lupus erythematosus (Madden, Montgomery). One observes considerable liquefaction degeneration in the basal cell layer and pronounced edema in the uppermost dermis (Fig. 143). Within the zone of subepidermal edema homogeneous eosinophilic fibrinoid material may be seen lying in irregular conglomerates as well as around capillaries (Fig. 144). Deeper in the dermis some of the collagen bundles are swollen and stain intensely eosinophilic. The inter-

strated only very rarely in the blood of patients with diseases other than acute systemic lupus erythematosus a positive test for L. E. cells is almost pathognomonic for it (Dubois)

The formation of L. E. cells is due to the presence in the plasma of a factor which causes disintegration of the nuclei of neutrophils, as well as of other cells with subsequent phagocytosis of this material by neutrophils (Haserick, Gold) The L. E. cell is a neutrophil con-




FIG. 146 Acute systemic lupus erythematosus. Smear of the buffy coat of a patient's blood after its incubation with heparin. In the left upper corner lies an L. E. cell a neutrophil containing a large smoky body. In the right lower corner a rosette is seen consisting of amorphous material surrounded by phagocytosing neutrophils ( $\times 400$ ).

taining a round structureless smoky basophilic mass of such size that it presses the lobes of the nucleus against the cell membrane (Fig. 146) (Hargraves Richmond and Morton Smith). L. E. cells can be demonstrated most easily in smears made from the buffy coat of the patient's blood after its incubation with heparin (Dubois). Smears of the buffy coat of heparinized sternal marrow may also be used. The smears are stained with Wright's stain. In addition to L. E. cells

the rosettes are precursors of the L. E. cell (Haserick).

The hematoxylin staining bodies are found in various organs, especially in the kidney and the endocardium but also occasionally in the

The verrucous endocarditis of lupus erythematosus (the so-called Libman-Sacks syndrome) is caused by focal fibrinoid degeneration of the subendothelial connective tissue of the endocardium and the subsequent round cell and fibroblastic proliferation and fibrosis. Since fibrinoid degeneration occurs again and again in the newly formed fibrous tissue considerable amounts of granulomatous tissue are formed resulting in raised verrucous formations (Libman and Sacks, Belote and Ratner, Gross, Klemperer, Pollack and Biehr 1911).

The serous membranes such as the pleura, the pericardium and the peritoneal covering of the liver and the spleen may show fibrinoid degeneration of their collagen with mild, reactive inflammation composed of lymphocytes, plasma cells, histiocytes and fibroblasts (Klemperer, Pollack and Biehr 1911).

The myocardium and less frequently the skeletal muscle may show small foci of degeneration in the interfascicular connective tissue and in the muscle bundles usually associated with a mild reactive inflammation. These changes are identical with those of dermatomyositis though much milder (Klemperer, Pollack and Biehr 1911, Jager).

The renal glomeruli not infrequently present wire loop lesions. In this type of lesion individual glomerular loops appear thickened, rigid and deeply eosinophilic because of fibrinoid degeneration and thickening of the basement membrane of glomerular capillaries (Klemperer, Pollack and Biehr 1911, Jager). In addition small foci of necrosis may occur in some of the glomeruli (Jager).

Periarterial fibrosis of the central arteries in the spleen is one of the most common lesions in acute systemic lupus erythematosus. Thick concentrically layered rings of sclerotic collagen fibers surround these arteries (Klemperer, Pollack and Biehr 1911, Krüger).

The lymph nodes often show foci of necrosis.

The body fat may show the same changes as those described for the subcutaneous fat.

Vascular changes are usually not conspicuous. Occasionally, however, the fibrinoid degeneration affects the collagenous ground substance of blood vessels. In rare instances vascular changes are marked and resemble those of periarthritis nodosa (Jircho, Mallory). It is possible that in these cases lupus erythematosus and periarthritis nodosa are present simultaneously.

Two structures are encountered frequently in acute systemic lupus erythematosus. L. E. cells in the blood and hematoxylin staining bodies in the tissue. They are absent in chronic discoid and chronic disseminated lupus erythematosus. Since L. E. cells can be demon-

strated only very rarely in the blood of patients with diseases other than acute systemic lupus erythematosus a positive test for L. E. cells is almost pathognomonic for it (Dubois)

The formation of L. E. cells is due to the presence in the plasma of a factor which causes disintegration of the nuclei of neutrophils, as well as of other cells with subsequent phagocytosis of this material by neutrophils (Haserick Gold) The L. E. cell is a neutrophil con-



Fig 14b Acute systemic lupus erythematosus Smear of the buffy coat of a patient's blood after its incubation with heparin In the left upper corner lies an L. E. cell a neutrophil containing a large smoky body In the right lower corner a rosette is seen consisting of amorphous material surrounded by phagocytizing neutrophils ( $\times 100$ )

taining a round structureless smoky basophilic mass of such size that it presses the lobes of the nucleus against the cell membrane (Fig 14b) (Hargraves Richmond and Morton Smith) L. E. cells can be demonstrated most easily in smears made from the buffy coat of the patient's blood after its incubation with heparin (Dubois) Smears of the buffy coat of heparinized sternal marrow may also be used The smears are stained with Wright's stain In addition to L. E. cells one often

Leukematoxilin staining bodies are found in various organs especially in the kidney and the endocardium but also occasionally in the



The verrucous endocarditis of lupus erythematosus (the so-called Libman Sacks syndrome) is caused by focal fibrinoid degeneration of the subendothelial connective tissue of the endocardium and the subsequent round cell and fibroblastic proliferation and fibrosis. Since fibrinoid degeneration occurs again and again in the newly formed fibrous tissue, considerable amounts of granulomatous tissue are formed resulting in raised verrucous formations (Libman and Sacks 1906 and Ratner Gross Klemperer Pollack and Baehr 1911).

The serous membranes such as the pleura the pericardium and the peritoneal covering of the liver and the spleen may show fibrinoid degeneration of their collagen with mild, reactive inflammation composed of lymphocytes, plasma cells, histiocytes and fibroblasts (Klemperer Pollack and Baehr 1911).

The myocardium and less frequently the skeletal muscle may show small foci of degeneration in the interfascicular connective tissue and in the muscle bundles usually associated with a mild reactive inflammation. These changes are identical with those of dermatomyositis though much milder (Klemperer Pollack and Baehr 1911 Jäger).

The renal glomeruli not infrequently present wire loop lesions. In this type of lesion individual glomerular loops appear thickened rigid and deeply eosinophilic because of fibrinoid degeneration and thickening of the basement membrane of glomerular capillaries (Klemperer Pollack and Baehr 1911 Jäger). In addition small foci of necrosis may occur in some of the glomeruli (Jäger).

Periarterial fibrosis of the central arteries in the spleen is one of the most common lesions in acute systemic lupus erythematosus. Thick concentrically layered rings of sclerotic collagen fibers surround these arteries (Klemperer Pollack and Baehr 1911 Krüger).

The lymph nodes often show foci of necrosis.

The body fat may show the same changes as those described for the subcutaneous fat.

Vascular changes are usually not conspicuous. Occasionally however the fibrinoid degeneration affects the collagenous ground substance of blood vessels. In rare instances vascular changes are marked and resemble those of perarteritis nodosa (Jucho Mallory). It is possible that in these cases lupus erythematosus and perarteritis nodosa are present simultaneously.

Two structures are encountered frequently in acute systemic lupus erythematosus. L. E. cells in the blood and hematoxylin staining bodies in the tissue. They are absent in chronic discoid and chronic disseminated lupus erythematosus. Since L. E. cells can be demon-

## LUPUS ERYTHEMATOSUS PROFUNDUS (KAPOSI URGANS)

In this rare condition cutaneous lesions of chronic discoid lupus erythematosus are present and in addition one or more firm sharply outlined ~~inmovable subcutaneous nodes covered by normal appearing skin~~

s that because subcutaneous ~~sarcom~~ of Darier Roussy. He concludes that lupus erythematosus profundus constitutes a coexistence of two different diseases—discoid lupus erythematosus and subcutaneous sarcom of Darier Roussy

## DERMATOMYOSITIS

In dermatomyositis the skin and the skeletal muscles are predominantly affected

The cutaneous lesions consist of extensive rather sharply defined areas of erythema and edema involving predominantly the face the chest and the arms. The eruption often greatly resembles that of subacute or acute lupus erythematosus. The lesions in the muscles cause progressive weakness with vague muscular pain and later atrophy of the muscles. Involvement of the esophagus often results in dysphagia. Prior to the use of corticotropin or cortisone the disease was often fatal. Even with its use atrophy and fibrosis of the skin and the muscles may result producing a clinical picture resembling that of generalized scleroderma.

**Histopathology** In the skin particularly in early lesions the histologic changes may resemble those of subacute disseminate lupus erythematosus (Kinney and Maher O Leary and Waisman). Not infrequently however the histologic picture is that of a nonspecific chronic dermatitis. In old lesions the collagen bundles of the dermis may show thickening homogenization and sclerosis and the cutaneous vessels fibrotic thickening of their walls so that the changes are indistinguishable from those of scleroderma (Dowling Freudenthal).

The subcutaneous fat may show mucoid degeneration of the fat cells and focal lymphocytic infiltration in the early stage and areas of fibrosis and calcification in the late stage (Norregaard Wainger and Lever) (see Calcinosi Cutis page 276).

During the active phase of the disease the skeletal muscles show

skin (Klemperer, Gueft, Lee, Leuchtenberger and Pollister, Gueft) They appear in sections stained with hematoxylin and eosin as red purple, homogeneous bodies of the size and the shape of fibroblasts or histiocytes They may be present singly or in aggregates They develop from degenerating nuclei and are identical with the smoky body within L. E. cells Occasionally, one may see histiocytes or neutrophils which have phagocytized a hematoxylin staining body The similarity of these cells to L. F. cells is striking Histochemical investigations have shown that, in the smoky bodies of L. E. cells and in the hematoxylin staining bodies, the Feulgen reaction for desoxyribonucleic acid (DNA) is positive This is evidence that they are derived from nuclear chromatin (Gueft)

**Relationship of Acute Systemic Lupus Erythematosus to Dermatomyositis and Generalized Scleroderma.** These three diseases—lupus erythematosus, dermatomyositis and scleroderma—are related to one another (Banks) The fundamental pathologic lesion, namely fibrinoid degeneration of the collagen, is the same in all three diseases (Klemperer, Pollack and Baehr, 1941, Baehr and Pollack) Only the response to the fundamental lesion and the organs affected differ in these three diseases One may summarize the response to fibrinoid degeneration of the collagen in these three diseases as follows

In lupus erythematosus, the degeneration is associated with mild inflammation, occurring, as a rule, in small foci The lesions are widespread

In dermatomyositis, the degeneration is associated with usually pronounced inflammation Lesions are present mainly in the skin and the striated musculature, but other organs, particularly the heart may be involved

In scleroderma the degeneration is associated with only slight inflammation but with marked fibrosis As in dermatomyositis the principal lesions are in the skin and the striated musculature In many cases, other organs (such as the heart the esophagus, the lungs and the kidneys) are involved also

The cause of fibrinoid degeneration of the collagen in these three diseases is not known It is assumed by some that the degeneration represents an allergic or hyperergic reaction to infection, particularly streptococcal infection However, Klemperer, Pollack and Baehr (1942) point out that, although fibrinoid degeneration of the collagen may occur in hypersensitivity reactions, it does not necessarily follow that all instances of fibrinoid degeneration are on the basis of hypersensitivity.

In older lesions the changes in the muscles may resemble those of scleroderma. The muscle bundles show sclerosis and atrophy and fibrotic connective tissue replaces the muscle bundles in many areas. On the basis of a study of these late changes as just described in the skin and the skeletal muscle several authors have concluded that dermatomyositis and generalized scleroderma represent one and the same disease (Dowling, Freudenthal).

Changes in organs other than the skin and the skeletal muscles occur but not so regularly as in acute systemic lupus erythematosus or in generalized scleroderma. The heart may show changes identical with those in the skeletal muscle though less severe (Kinney and Maher, O'Leary and Waisman, Winger and Lever). The body fat may be affected similarly as the subcutaneous fat (Greenway and Lambie, Kinney and Maher, Winger and Lever). Inflammatory changes in the serous membranes may occur (Kinney and Maher). Ulcerative lesions in the gastro-intestinal tract due to vascular occlusions have been described (Karelitz and Welt, Horn, Winger and Lever).

Of interest is the relatively common occurrence of visceral carcinoma in patients with dermatomyositis (Dostrovsky and Siglier). So far 31 cases of carcinoma in dermatomyositis have been reported (Schuermann). In a statistical analysis Schuermann concludes that carcinoma occurs at least five times more frequently in patients with dermatomyositis than in the normal population. In contrast with this only one case of carcinoma in scleroderma has ever been reported.

#### POIKILODERMA ATROPHICANS VASCULARE (JACOBI)

Poikiloderma atrophicans vasculare has been described in a few instances as an independent disease (Marchionini and Rosser, Dow-

As a rule

It occurs

usually in dermatomyositis but also in lupus erythematosus and in mycosis fungoides (see page 484). When associated with dermatomyositis the term poikilodermatomyositis is often employed.

Clinically poikiloderma atrophicans vasculare presents large ill defined areas usually in symmetrical distribution which in the early stage show erythema and slight scaling associated with

is pronounced. The clinical picture then resembles chronic radiodermatitis.

*degenerative changes and inflammation.* The degree of these changes varies not only in different muscles but also within each affected muscle. Even in severely affected muscles, close to areas of severe damage, one often finds areas of mild degeneration and areas in which the muscle bundles appear normal (Fig. 147). In areas of mild degeneration, the muscle bundles exhibit effacement of transverse striation,




FIG 147 Dermatomyositis, muscle The muscle bundles show various degrees of degeneration. In addition, one sees edema and focal collections of inflammatory cells ( $\times 100$ )

coagulation or hyalinization of the sarcoplasm and proliferation of nuclei. In more severely degenerated areas, the muscle bundles show fragmentation of fibers, granular and vacuolar degeneration, basophilic staining and phagocytosis by large histiocytes. Inflammatory changes are secondary and not essential; they represent a reaction to the parenchymatous damage. One observes a cellular infiltrate, composed largely of lymphocytes, but containing also plasma cells, histiocytes and fibroblasts, between the muscle bundles, either in perivascular arrangement or distributed diffusely. In addition, edema usually separates the muscle bundles. The blood vessels are dilated, but, as a rule, their walls reveal no abnormalities (O'Leary and Waisman).

In older lesions the changes in the muscles may resemble those of scleroderma. The muscle bundles show sclerosis and atrophy and fibrotic connective tissue replaces the muscle bundles in many areas. On the basis of a study of these late changes as just described in the skin and the skeletal muscle several authors have concluded that dermatomyositis and generalized scleroderma represent one and the same disease (Dowling, Freudenthal).

Changes in organs other than the skin and the skeletal muscles occur but not so regularly as in acute systemic lupus erythematosus or in generalized scleroderma. The heart may show changes identical with those in the skeletal muscle though less severe (Kinney and Maher, O'Leary and Waisman, Winger and Lever). The body fat may be affected similarly as the subcutaneous fat (Greenway and Lambie, Kinney and Maher, Winger and Lever). Inflammatory changes in the serous membranes may occur (Kinney and Maher). Ulcerative lesions in the gastro-intestinal tract due to vascular occlusions have been described (Karelitz and Welt, Horn, Winger and Lever).

Of interest is the relatively common occurrence of visceral carcinoma in patients with dermatomyositis (Dostrovsky and Sagher). So far 31 cases of carcinoma in dermatomyositis have been reported (Schuermann). In a statistical analysis Schuermann concludes that carcinoma occurs at least five times more frequently in patients with dermatomyositis than in the normal population. In contrast with this only one case of carcinoma in scleroderma has ever been reported.

#### POIKILODERMA ATROPHICANS VASCULARE (JACOBI)

Poikiloderma atrophicans vasculare has been described

as a disease of the skin. It occurs most frequently in dermatomyositis but also in lupus erythematosus and in mycosis fungoides (see page 484). When associated with dermatomyositis the term poikilodermatomyositis is often employed.

Clinically poikiloderma atrophicans vasculare presents large ill defined areas usually in symmetrical distribution which in the early stage show erythema and slight scaling, a mottled pigmentation and numerous telangiectases. In the late stage the skin appears atrophic, the erythema has largely disappeared but the mottled pigmentation and the telangiectases are more pronounced. The clinical picture then resembles chronic radiodermatitis.

**Histopathology** In the idiopathic form of poikiloderma and in poikiloderma associated with dermatomyositis or lupus erythematosus the histologic changes are identical. In the early active stage the epidermis shows moderate atrophy of the stratum malpighii, effacement of the rete ridges and hydropic degeneration of the basal cells (Fig. 148). In the upper dermis one finds a fairly dense cellular



FIG. 148 Poikilodermatomyositis, early stage. The epidermis shows atrophy and hydropic degeneration of the basal cells. In the upper dermis one sees a fairly dense inflammatory infiltrate which in places invades the epidermis. The collagen of the upper dermis shows edema and hyalinization ( $\times 200$ ).

infiltrate which in places invades the epidermis and often has a band-like arrangement. The infiltrate consists of lymphocytes, histiocytes and fibroblasts. Many melanophores may be present. Some of the superficial capillaries are dilated. The collagen is edematous and shows hyaline degeneration. The elastic tissue is largely destroyed. Hair follicles and sebaceous glands are absent (Horn).

In the late stage the epidermis is atrophic. There is marked dilatation of superficial capillaries. The dermis shows homogenization and sclerosis of the collagen with little or no inflammatory infiltrate. Melanophores are present in varying numbers. In contrast with scleroderma the dermis is greatly thinned (Guy, Gruver and Jacob).

In poikiloderma associated with mycosis fungoides the histologic changes in the dermis are those of mycosis fungoides (see page 489).

**Differential Diagnosis** Subacute disseminate lupus erythematosus like poikiloderma atrophicum vasculare shows atrophy of the stratum malpighii and vacuolization of the basal layer. However the presence of the superficially located bandlike infiltrate in poikiloderma atrophicum vasculare usually makes a differentiation possible.

## SCLERODERMA

Two forms of scleroderma occur: circumscribed scleroderma (morphea) and systemic or generalized scleroderma.

**Clinical Appearance** In circumscribed scleroderma (morphea) one or several round, oval or irregularly shaped, smooth, indurated patches are present. They are at first dull red or violaceous in color but soon assume an ivory color. As long as there is peripheral extension the patches tend to have a purplish halo (lilac ring). The disease is benign. Internal lesions are absent.

In systemic or generalized scleroderma large areas of the skin are affected. At first the involved areas present diffuse induration. As the disorder progresses the skin and the subcutaneous tissue become firmly bound to the underlying structures so that motion is difficult. Gradually the skin and the subcutaneous tissue undergo atrophy but even at this stage the skin retains some of its induration. The face and the hands are often the most severely affected areas (acro-sclerosis). In addition to the skin the striated musculature variably is affected resulting in weakness and muscular atrophy. Involvement of the esophagus may lead to difficulties in deglutition of the heart to cardiac insufficiency of the lungs to dyspnea.

**Histopathology of Circumscribed Scleroderma (Morphea)** An early inflammatory and a late sclerotic stage exist. Most sections obtained routinely show a histologic picture intermediary between the two stages.

In the early lesion the collagenous bundles appear swollen and homogeneous. They are separated by edema. An inflammatory infiltrate predominantly lymphocytic is present between the collagenous bundles and around blood vessels (Fig. 149). The walls of the vessels are edematous. These changes are present throughout the dermis. In addition the inflammatory infiltrate extends between the fat cells of the subcutaneous layer causing degeneration of the fat cells. The elastic fibers are frayed and may be destroyed.

In a late lesion the dermis is markedly thickened. The collagen bundles are hypertrophic, sclerotic and closely packed (Fig. 150). Fibroblasts are fewer than in the normal dermis. The inflammatory infiltrate has disappeared almost completely except around the vessels where as a rule one still finds a few inflammatory cells. Most of



the vessels in the dermis show marked thickening and sclerosis of their walls with narrowing of their lumen. Sebaceous glands and hair structures are often completely absent. Sweat glands on the other hand, are still present. They are reduced in number and are atrophic. Instead of lying close to the cutaneous-subcutaneous border and being surrounded by fat cells they lie in the midst of sclerotic




FIG. 149 Scleroderma, early stage. The collagenous bundles appear swollen and homogeneous and are separated by edema. An inflammatory infiltrate, predominantly lymphocytic, is present between the collagenous bundles. The wall of the blood vessel seen on the right side is thickened by edema and beginning fibrosis. ( $\times 100$ )

collagenous bundles tightly bound down by them. The fact that the sweat glands lie inside the dermis rather than at its lower border is evidence that the thickening of the dermis is produced not alone by hypertrophy of the pre-existing collagen bundles but also by new formation of collagen at the lower border of the dermis. The border between the dermis and the subcutaneous layer is not so sharp as it usually is, because thick strands of sclerotic collagen extend from the dermis into the subcutaneous layer and replace much of the subcutaneous fat. The thickness of the subcutaneous layer may be greatly reduced. The vessels in the subcutaneous layer, including those of large caliber, often show marked thickening of all their coats with

narrowing of the lumen (O Leary and Nomland) It is worth noting that the rete ridges of the epidermis usually remain well preserved in spite of the fact that the thickened collagen bundles extend right up to it




FIG 150 Scleroderma late stage The collagen bundles

( $\times 100$ ) are bound down by collagen

**Histopathology of Generalized Scleroderma** The cutaneous changes are essentially the same as in circumscribed scleroderma so that a histologic differentiation of the two types is not possible in the early stage. In the circumscribed form the collagen bundles are initially small and

the vessels in the dermis show marked thickening and sclerosis of their walls with narrowing of their lumina. Sebaceous glands and hair structures are often completely absent. Sweat glands, on the other hand, are still present, they are reduced in number and are atrophic. Instead of lying close to the cutaneous subcutaneous border and being surrounded by fat cells, they lie in the midst of sclerotic




FIG. 149 Scleroderma, early stage. The collagenous bundles appear swollen and homogeneous and are separated by edema. An inflammatory infiltrate, predominantly lymphocytic, is present between the collagenous bundles. The wall of the blood vessel seen on the right side is thickened by edema and beginning fibrosis. ( $\times 100$ )

collagenous bundles, tightly "bound down" by them. The fact that the sweat glands lie inside the dermis rather than at its lower border is evidence that the thickening of the dermis is produced not alone by hypertrophy of the pre-existing collagen bundles but also by new formation of collagen at the lower border of the dermis. The border between the dermis and the subcutaneous layer is not so sharp as it usually is, because thick strands of sclerotic collagen extend from the dermis into the subcutaneous layer and replace much of the subcutaneous fat. The thickness of the subcutaneous layer may be greatly reduced. The vessels in the subcutaneous layer, including those of large caliber, often show marked thickening of all their coats with

up to it




FIG. 150. Scleroderma, late stage. The collagen bundles are hypertrophic sclerotic and closely packed together. Only very little inflammatory infiltrate is present. Fibroblasts are fewer than in the normal dermis. Groups of rather atrophic sebaceous glands (SG) are tightly bound down by collagen ( $\times 100$ ).

**Histopathology of Generalized Scleroderma** The cutaneous changes are essentially the same as in circumscribed scleroderma, so that a histologic differentiation of the two types is not possible. In the early stage degenerative changes are often more severe than in the circumscribed type. One may observe fibrinoid degeneration of the collagen (Pollack). The walls of the vessels may show marked intimal proliferation, fibrinoid degeneration and inflammatory infiltration (Masugi and Ya Shu). An occasional vessel may show throm-

bosis. In the late stage, focal or even extensive calcification may take place in the lower dermis and in the subcutaneous layer (Talbot Gall, Consolazio and Coombs, Brody and Bellin, Kamee) (See also "Calcinosis Cutis," page 276) Also, in the late stage of generalized scleroderma, the epidermis may show, in contrast to circumscribed scleroderma, atrophy with disappearance of the rete ridges

**Histopathology of the Visceral Lesions of Generalized Scleroderma.** There are often extensive systemic manifestations. The skeletal muscles are nearly always affected severely. In early lesions degeneration of muscle bundles with accompanying inflammation similar to *dermatomyositis* but less severe, may be observed occasionally. However, the characteristic changes are homogenization, sclerosis and atrophy of the muscle bundles with increase in the interstitial connective tissue. In contrast with *dermatomyositis*, the interseptal blood vessels often show marked obliterative changes. Foci of chronic inflammation may be present around the vessels. The musculature of the esophagus is often affected similarly (Lindsay, Templeton and Rothman). Extensive atrophy and fibrosis may occur in the heart muscle (Weiss, Stead, Warren and Bailey) and in the intestinal musculature (Bevans). Widespread sclerosis of the medium sized and the small sized vessels in the myocardium has been observed.

Pathologic changes may occur also in the endocardium and the epicardium (Pollack), in the serous membranes (Bevans) and in the esophageal and the intestinal mucosa (Bevans). The changes consist in early lesions, of fibrinoid degeneration of the collagen with reactive inflammation, as in acute systemic lupus erythematosus. In older lesions, homogenization and sclerosis of the collagen predominate.

Additional findings in occasional instances include glomerulitis of the wire loop type, as in acute systemic lupus erythematosus (Pollack, Bevans), extensive fibrosis with cystic changes in the lungs (Dostrovsky), and fibrosis of the thyroid (Bevans).

### PERIARTERITIS NODOSA

Periarteritis nodosa is a manifestation of hypersensitivity. Although, in most cases, the cause of the disease is not apparent, the administration of foreign serum or of sulfonamides may be the cause (Rich). The disease affects mainly small arteries and arterioles, but occasionally also veins. The principal vessels affected are those of the gastro intestinal tract, the kidneys and the heart, but those of the brain, the lungs and the skin as well as those of other organs may be involved.

Depending on the sites of involvement the clinical symptoms may vary. Severe abdominal pain and symptoms of nephritis and of myocardial disease are the most common manifestations. Irregular fever and marked prostration are present. The disease is fatal in the vast majority of cases.

Cutaneous manifestations are found in about one third of the cases. They are manifold and may consist of macules, papules, nodules, petechiae, ecchymoses and necrotic ulcers (Fig. 151). Subcutaneous nodules, usually movable and painless, are observed occasionally (Ketron and Bernstein). In some cases extensive areas of cutaneous hemorrhage with subsequent necrosis occur (Melzer and Venkei).

**Histopathology.** The name periarteritis is misleading for the lesions actually represent a panarteritis. On a histologic basis the changes which occur in the arteries and the arterioles may be divided into four stages (Arkin Weir).

In the first, the degenerative stage, sections of the intima and the media undergo necrosis. The necrosis often affects only a segment rather than the entire circumference of the vessel, and only portions of the vessels are involved. Milium aneurysms may form in areas of segmental necrosis.

In the second, the inflammatory stage, the necrotic area and the adjoining adventitia are densely infiltrated with polymorphonuclear leukocytes and eosinophils, and sometimes also with lymphocytes and plasma cells (Fig. 152). The infiltrate extends to the perivascular tissue. The lumen of most vessels thus affected is thrombosed.

In the third, the granulation stage, the necrotic part of the vascular wall is replaced by granulation tissue and the intima shows proliferation leading to partial or total occlusion of the lumen.

In the fourth, the fibrotic stage, the destroyed vascular wall is replaced by scar tissue. The lumen may show reduction in size, obliteration or recanalization.

In the skin, the arterioles at the cutaneous-subcutaneous border and in the subcutaneous tissue may show the typical changes of periarteritis nodosa (Fig. 151). However, the small blood

vessel walls composed largely of neutrophils and eosinophils, many of which show fragmentation of their nuclei. Thrombus formation with or without inflammatory reaction also may occur in the cutaneous vessels. The vasculitis and the throm-

bus formation cause extravasation of erythrocytes (Ketron and Bernstein).

It has been noted by several authors (Carol and Prakken, Miescher, Slinger and Starck) that periarteritis nodosa may occur in a benign,



FIG 151 Periarteritis nodosa. Low magnification An artery (A) at the cutaneous-subcutaneous border shows the granulation stage of periarteritis nodosa Above the artery is an ulcer which is probably caused by the occlusion of the artery ( $\times 50$ )

chronic form limited to the skin and the subcutaneous tissue These authors state that, histologically, the lesions of the benign form are indistinguishable from those of the malignant systemic form

#### ALLERGIC GRANULOMATOSIS

An attempt has been made recently to divide periarteritis nodosa into two types: "true periarteritis nodosa" and "allergic granulomatosis" (Churg and Strauss, Strauss, Churg and Zak; Zeek) In the

latter group allergic manifestations dominate the clinical picture to a much greater extent than in true periarteritis nodosa. Most patients have severe asthma as the first symptom and in addition to cutaneous lesions including intracutaneous and subpronounced. The course is




FIG. 139. Periarteritis nodosa. High magnification. A medium sized artery located in the lower dermis shows partial necrosis of its wall and invasion by inflammatory cells. (X200)

chronic and usually fatal. Strauss, Churg and Zak believe that periarteritis nodosa as a result of drug sensitivity usually manifests itself as allergic granulomatosis.

**Histopathology.** Histologically in addition to typical widespread lesions of periarteritis nodosa one finds extravascular granulomas in many organs. The cutaneous and the subcutaneous nodules show no necrotizing arteritis but only extravascular granulomas. They consist of areas of central necrosis surrounded by radially arranged histiocytes and foreign body giant cells which are embedded in a diffuse inflammatory infiltrate rich in eosinophils (Strauss, Churg and Zak).

The purpuric lesions show the same appearance as those of anaphylactoid purpura (see page 127).



bus formation cause extravasation of erythrocytes (Ketron and Bernstein).

It has been noted by several authors (Carol and Prakken, Miescher, Slinger and Starck) that periarteritis nodosa may occur in a benign



FIG 151 *Periarteritis nodosa* Low magnification An artery (A) at the cutaneous subcutaneous border shows the granulination stage of *periarteritis nodosa* Above the artery is an ulcer which is probably caused by the occlusion of the artery ( $\times 50$ )

chronic form limited to the skin and the subcutaneous tissue These authors state that, histologically the lesions of the benign form are indistinguishable from those of the malignant systemic form

#### ALLERGIC GRANULOMATOSIS

An attempt has been made recently to divide *periarteritis nodosa* into two types "true *periarteritis nodosa*" and "allergic granulomatosis" (Churg and Strauss, Strauss, Churg and Zak, Zeck) In the

- Pautrier L M A propos du pseudo-lupus érythémateux profond (Kaposi Irgang) Ann de dermat et de syph 80 233 1953
- Smith P A J The L. E. cell and its significance Brit J Dermat 64 10 1952 (Good review)
- Stoughton R and Wells G A histochemical study on polysaccharides in normal and diseased skin J Invest. Dermat 14 57 1950

### Dermatomyositis and Poikiloderma Atrophicum Vasculare (Jacobi)

- Dostrovsky A and Sagher F Dermatomyositis and malignant tumour Brit J Dermat 58 52 1946
- Dorling G B Generalized scleroderma Brit J Dermat 52 242 1940
- and Freudenthal W Dermatomyositis and poikiloderma atrophicum vasculare A clinical and histological comparison Brit J Dermat 50 519 1938
- Dorling J G Edelstein J M and Fitzpatrick T R Poikiloderma vasculare atrophicum Arch Dermat & Syph 56 740 1947
- Freudenthal W Generalized scleroderma and dermatomyositis a histological comparison Brit J Dermat 52 289 1940
- Greenway T M and Lambie C G Dermatomyositis Brit J Dermat 49 209 1937
- Guy W B Graver R C and Jacob F M Poikilodermatomyositis Arch Dermat & Syph 40 867 1939
- — — — — 44 1086 1941
- — — — — d 43 1134 1932
- — — — — ch. 16 561 1940
- — — — — hicans vasculare

— — — — — 1932

- Norregaard S A case of dermatomyositis with universal calcinosis Acta dermat venereol 27 479 1947
- O'Leary P A and Waisman M Dermatomyositis A study of forty cases Arch. Dermat & Syph 41 1001 1940 (Good review)
- Schuermann H Maligne Tumoren bei Dermatomyositis und progressiver Scleroderme Arch f Dermat u Syph 192 575 1951
- Wainger C K and Lever W F Dermatomyositis A report of three cases with postmortem observations Arch Dermat & Syph 59 196 1949 (Review of findings in internal organs)

### Scleroderma

- Bevan M Pathology of scleroderma with special reference to the changes in the gastrointestinal tract Am J Path 21 23 1945
- Brody J and Bellin D E Calcinosis with scleroderma Arch Dermat. & Syph 36 83 1937
- Dostrovsky A Progressive scleroderma of the skin w k r ch. 16 561 1940
- — — — — 44 1086 1941
- — — — — d 43 1134 1932
- — — — — ch. 16 561 1940
- — — — — hicans vasculare
- Lindsay J R Templeton F E and Rothman S 1941
- Matsu 1
- O'Leary P A and Normland R A clinical study of 103 cases of scleroderma Am J M Sc. 180 93 1930

## BIBLIOGRAPHY

## Lupus Erythematosus

- Altshuler C H and Angevine D M Histochemical studies on the pathogenesis of fibrinoid *Am J Path* 25 1061 1949
- Arnold H L Jr Lupus erythematosus profundus (Kaposi's Irgang) *Arch Dermat & Syph* 57 196 1948
- Baehr, G and Pollack A D Disseminated lupus erythematosus and diffuse scleroderma *JAMA* 134 1169 1947
- Banks B M Is there a common denominator in scleroderma dermatomyositis disseminated lupus erythematosus the Libman-Sacks syndrome and polyarteritis nodosa? *New England J Med* 225 433 1941
- Belote G H and Ratner H S V The so called Libman-Sacks syndrome *Arch Dermat & Syph* 33 642 1936
- Dubois E I Simplified method for the L E cell test *Arch Int Med* 92 168 1953
- Gold S C Some observations on the pathogenesis of systemic lupus erythematosus *J Invest Dermat* 19 333 1952
- Gross L The cardiac lesions in Libman-Sacks disease with a consideration of its relationship to acute diffuse lupus erythematosus *Am J Path* 16 375 1940
- Gueft B Depolymerization of nucleic acid in acute disseminated lupus erythematosus *Arch Dermat & Syph* 61 892 1950
- Hargraves M M Richmond H and Morton R J Presentation of two bone marrow elements the mitotic cell and the L E cell *Proc Staff Meet Mayo Clinic* 23 25 1948
- Haserick J R Blood factor in acute disseminated lupus erythematosus *Arch Dermat & Syph* 61 889 1950
- Irgang S Lupus erythematosus profundus *Arch Dermat & Syph* 42 97 1940
- Jager B V Disseminated lupus erythematosus *Arch Dermat & Syph* 46 362 1942
- Jarcho S Lupus erythematosus associated with visceral vascular lesions *Bull Johns Hopkins Hosp* 59 262 1936
- Kaiser I H The specificity of periarterial fibrosis of the spleen in disseminated lupus erythematosus *Bull Johns Hopkins Hosp* 71 31 1942
- Klemperer P Gueft B Lee S L Leuchtenberger C and Pollister A W  
*Arch Path* 49 503 1950  
 disseminated lupus erythematosus  
 pathology of visceral lesions)
- and —— Diffuse collagen disease *JAMA* 119 331 1942
- Libman E and Sacks B A hitherto undescribed form of valvular and mural endocarditis *Arch Int Med* 33 701 1924
- McCreight W G and Montgomery H Cutaneous changes in lupus erythematosus *Arch Dermat & Syph* 61 1 1950 (Good review)
- Madden J F Acute disseminated lupus erythematosus *Arch Dermat & Syph* 25 854 1932
- Mallory T B Cabot case 24 201 lupus erythematosus *New England J Med* 218 838 1938
- Montgomery H Pathology of lupus erythematosus *J Invest Dermat* 2 343 1939

# 18

## Epidermal Tumors

### HISTOGENESIS AND CLASSIFICATION OF EPIDERMAL TUMORS

The epidermal tumors may be divided into two classes namely tumors of the surface epidermis and tumors of the epidermal appendages. In each class nevroid tumors (or hamartomas) and carcinomas occur.

Nevroid tumors or hamartomas are defined as benign neoplasms which usually arise in accordance with the Cohnheim theory from immature ring adult life. Carcinomas posed according to Hanse mann's theory of anaplastic cells. They usually arise from mature cells due to their change into anaplastic cells but embryonal cells

able to exist more independently than embryonal cells and therefore survive when carried away through the lymphatics and multiply as metastases.

**Tumors of the Surface Epidermis** The tumors of the surface epidermis may be classified as follows:

- 1 Nevroid (benign) tumors
  - a. Nevus verrucosus (papilloma)
  - b. Epidermal cyst
- 2 Precancerous tumors
  - a. Keratosis senilis
  - b. Leukoplakia
- 3 Carcinomas
  - a. Squamous-cell carcinoma (epidermoid carcinoma)
  - b. Bowen's disease (intra epidermal squamous-cell carcinoma)
  - c. Paget's disease

**Tumors of the Epidermal Appendages** The nevroid tumors or hamartomas of this group can be divided according to the decreasing degree of organization and differentiation observed in them into

## 316 Systemic Diseases of Unknown Cause

- Pollack A D Visceral and vascular lesions in scleroderma Arch Path 29:859 1940
- Talbott J H Gail E A Consolizio W V and Coombs F S Dermatomyositis with scleroderma calcinosis and renal endarteritis associated with focal cortical necrosis Arch Int Med 63:476 1939
- Weiss S Stead E A Warren J V and Bailey O T Scleroderma heart disease Arch Int Med 71:749 1943

### Periarteritis Nodosa

- Arkin A A clinical and pathological study of periarteritis nodosa Am J Path 6:401 1930
- Carol W I L and Prakken J R Die kutane Form der Periarteritis nodosa Acta dermat venereol 18:102 1937
- Churg J and Strauss L Allergic granulomatosis allergic angitis and periarteritis nodosa Am J Path 27:277 1951
- Ketron L W and Bernstein J C Cutaneous manifestations of periarteritis nodosa Arch Dermat & Syph 40:929 1939 (Good review)
- Melzer N and Venken T Ueber die Hautformen der Periarteritis nodosa Dermatolog 91:214 1947
- Miescher G Ueber kutane Formen der Periarteritis nodosa Dermatolog 90:995 1946
- Rich A R The role of hypersensitivity in periarteritis nodosa Bull Johns Hopkins Hosp 71:123 375 1912
- Slinger W N and Starck V Cutaneous form of polyarteritis nodosa Arch Dermat & Syph 63:461 1951
- Strauss L Churg J and Zak F G Cutaneous lesions of allergic granulomatosis J Invest Dermat 17:349 1951
- Weir D R Polyarteritis nodosa Am J Path 15:79 1939
- Zeek P M Periarteritis nodosa a critical review Am J Clin Path 22:777 1952

## Epidermal Tumors

### HISTOGENESIS AND CLASSIFICATION OF EPIDERMAL TUMORS

The epidermal tumors may be divided into two classes, namely, tumors of the surface epidermis and tumors of the epidermal appendages. In each class nevoid tumors (or hamartomas) and carcinomas occur.

Nevoid tumors or hamartomas are defined as benign neoplasms which usually arise in accordance with the *Cohnheim theory*, from

Mann's theory of anaplastic cells. They usually arise from mature cells due to their change into anaplastic cells, but embryonal cells may also occasionally change into anaplastic cells and thus give rise to carcinoma. Anaplastic cells are dedifferentiated cells which behave differently from embryonal cells (*Foulds*). They are autonomous, able to exist more independently than embryonal cells and, therefore survive when carried away through the lymphatics and multiply as metastases.

**Tumors of the Surface Epidermis** The tumors of the surface epidermis may be classified as follows:

1. Nevoid (benign) tumors
  - a. *Nevus verrucosus* (papilloma)
  - b. Epidermal cyst
2. Precancerous tumors
  - a. *Keratosis senilis*
  - b. *Leukoplakia*
3. Carcinomas
  - a. Squamous-cell carcinoma (epidermoid carcinoma)
  - b. *Bowen's disease* (intra-epidermal squamous cell carcinoma)
  - c. *Paget's disease*

**Tumors of the Epidermal Appendages** The nevoid tumors, or hamartomas of this group can be divided, according to the decreasing degree of organization and differentiation observed in them, into

four groups organic nevi adenomas benign epitheliomas and basal cell epitheliomas (See Table 6) The carcinomas of the epidermal appendages can be divided into sebaceous gland carcinomas eccrine gland carcinomas and apocrine gland carcinomas

The author has advanced the thesis that the organic nevi the adenomas the benign epitheliomas and the basal cell epitheliomas develop from arrested embryonal primary epithelial germ cells and, as such are primary epithelial germ tumors Since the primary epithelial germ is an embryonal structure (see Chart 1 page 4) the tumors developing from it can accordingly be regarded as nevroid tumors or hamartomas It is the author's belief that the organic nevi the adenomas and the benign epitheliomas arise from primary epithelial germ cells which prior to the onset of neoplasia attained a certain degree of differentiation whereas the basal cell epitheliomas arise from primary epithelial germ cells which attained no or only little differentiation In accordance with the potentiality inherent in the primary epithelial germ to differentiate into sebaceous glands apocrine glands and hair differentiation in the tumors developing from the primary epithelial germ can be toward either sebaceous gland apocrine gland or hair structures (see Table 6) Pinkus recently has suggested that those tumors in this group which arise in later life—especially the basal cell epitheliomas—do not necessarily arise from congenitally preformed epithelial germ rests but from immature pluripotential cells forming in later life

The organic nevi are composed of mature organic structures The cutaneous adenomas show less differentiation than the organic nevi nonetheless well developed glandlike structures are present In the benign epitheliomas there is a further step down in respect to differentiation The basal cell epitheliomas are the least differentiated of the primary epithelial germ tumors

Basal cell epitheliomas are not considered carcinomas because they do not metastasize Their origin from basal cells is doubted because their cells in contrast with basal cells do not possess intercellular bridges They are believed to originate from the primary epithelial germ for the following reasons (1) they occur only in areas where primary epithelial germ structures (sebaceous glands apocrine glands or hair) are found (2) the cells of undifferentiated basal cell epitheliomas resemble the cells of primary epithelial germs as found in embryos from 4 to 6 months old (Fig. 1) (3) structures resembling sebaceous glands tubular glands and hair are present in many basal cell epitheliomas and (4) basal cell epitheliomas frequently are seen in the same lesion with primary epithelial germ tumors of higher differentiation

TABLE 6—CLASSIFICATION OF THE NEVOID TUMORS OF THE EPIDERMAL APPENDAGES (PRIMARY EPITHELIAL GERM TUMORS)

	WITH SEBACEOUS DIFFERENTIATION	WITH APOCRINE DIFFERENTIATION	WITH HAIR DIFFERENTIATION
I Organic hamartomas (organic nevi)	Sebaceous nevi 1 Nevus sebaceus (Jadassohn) 2 Adenoma sebaceum (Pringle) 3 Senile sebaceous nevus 4 Fordyce disease	Apocrine nevi	Hair nevi
II Organoid hamartomas (adenomas)	Sebaceous adenoma	Apocrine adenomas 1 Syringocystadenoma papilliferum 2 Hidradenoma papilliferum	
III Suborganoid hamartomas (benign epitheliomas)	Sebaceous epithelioma	Apocrine epitheliomas 1 Syringoma 2 Cylindroma 3 Myo-epithelioma	Hair epitheliomas 1 Tricho-epithelioma (epithelioma adenoides cysticum) 2 Calcifying epithelioma
IV Nonorganic hamartomas a Differentiated basal cell epitheliomas b Undifferentiated basal cell epitheliomas	Cystic basal-cell epithelioma  —	Adenoid basal cell epithelioma  —	Keratotic basal cell epithelioma  —



Some of the terms used to designate groups of tumors require definition

**NEVUS** This term is used in the literature in two different ways referring either to a tumor composed of nevus cells (nevocellular nevus, pigmented nevus) or to a lesion originating from embryonal cells and composed of mature or nearly mature structures (organic nevi, such as nevus vasculosus, nevus sebaceus, nevus pilosus and nevus verrucosus). In order to avoid confusion, it is advisable to use the term nevus when referring to an organic nevus always with a qualifying adjective, so that nevus without qualifying adjective designates a tumor composed of nevus cells. According to Jadassohn organic nevi are tumors and not hyperplasias because they develop on the basis of an abnormal germ anlage and not by an increase in the size and number of mature structures.

**NEVOID TUMOR (HAMARTOMA)** The term nevoid tumor is used widely as a designation for benign tumors of embryonal origin. However, this term lacks conciseness and therefore is unsatisfactory. The term hamartoma appears more satisfactory. This term, derived from the Greek word *hamartanem* (fail, miss, err) was coined by Albrecht as a designation for tumorlike malformations showing a faulty mixture of the normal components of the organ in which they occur. Van der Valk enlarged the concept of hamartoma to include all benign tumors of embryonal origin with an organoid structure. Since no satisfactory term exists for the entire group of benign tumors of embryonal origin, the author has suggested that the meaning of the term hamartoma be extended to include all such tumors even those with suborganoid and nonorganic structure. Thus, the tumors arising from the primary epithelial germ may be divided into organic hamartomas (organic nevi), organoid hamartomas (adenomas), suborganoid hamartomas (benign epitheliomas) and nonorganic hamartomas (basal cell epitheliomas).

**EPITHELIOMA** The term epithelioma is used by many authors as a synonym for carcinoma of the epidermis. Since, however, the true meaning of the word is tumor of the epithelium, the term may be used, as suggested by Jadassohn, as a designation of benign as well as malignant tumors of the epidermis provided that a qualifying adjective is added. It would perhaps be best if, as Becker has suggested, the term epithelioma be reserved for benign epidermal tumors and carcinoma for malignant epidermal tumors.

# I. TUMORS OF THE SURFACE EPIDERMIS

## NEVUS VERRUCOSUS

Nevus verrucosus is known also under various other clinical designations, such as hard nevus, epidermal nevus, nevus unius lateralis, linear nevus keratotic nevus and ichthyosis hystrix



FIG 153 Nevus verrucosus There are marked hyperkeratosis, acanthosis and papillomatosis The rete ridges are elongated (X50)

The lesions may be single or multiple and usually are present at birth They consist of verrucous growths of brownish color which often show linear configuration Large horny excrescences may be present

**Histopathology.** Nevus verrucosus shows hyperkeratosis, papillomatosis and acanthosis with elongation of the rete ridges (Fig 153) Thus, it has the histologic appearance of a papilloma

Some of the terms used to designate groups of tumors require definition

**NEVUS** This term is used in the literature in two different ways referring either to a tumor composed of nevus cells (nevocellular nevus pigmented nevus) or to a lesion originating from embryonal cells and composed of mature or nearly mature structures (organic nevus such as nevus vasculosus nevus sebaceus nevus pilosus and nevus verrucosus) In order to avoid confusion it is advisable to use the term nevus when referring to an organic nevus always with a qualifying adjective so that nevus without qualifying adjective designates a tumor composed of nevus cells According to Jadassohn organic nevi are tumors and not hyperplasias because they develop on the basis of an abnormal germ Anlage and not by an increase in the size and number of mature structures

**NEVOID TUMOR (HAMARTOMA)** The term nevoid tumor is used widely as a designation for benign tumors of embryonal origin However this term lacks conciseness and therefore is unsatisfactory The term hamartoma appears more satisfactory This term derived from the Greek word *hamartanein* (fail miss err) was coined by Albrecht as a designation for tumorlike malformations showing a faulty mixture of the normal components of the organ in which they occur Van der Valk enlarged the concept of hamartoma to include all benign tumors of embryonal origin with an organoid structure Since no satisfactory term exists for the entire group of benign tumors of embryonal origin the author has suggested that the meaning of the term hamartoma be extended to include all such tumors even those with suborganoid and nonorganic structure Thus the tumors arising from the primary epithelial germ may be divided into organic hamartomas (organic nevus) organoid hamartomas (adenomas) suborganoid hamartomas (benign epitheliomas) and nonorganic hamartomas (basal cell epitheliomas)

**EPITHELIOMA** The term epithelioma is used by many authors as a synonym for carcinoma of the epidermis Since however the true meaning of the word is tumor of the epithelium the term may be used as suggested by Jadassohn as a designation of benign as well as malignant tumors of the epidermis provided that a qualifying adjective is added It would perhaps be best if as Becker has suggested the term epithelioma be reserved for benign epidermal tumors and carcinoma for malignant epidermal tumors

cans These five diseases all show hyperkeratosis and papillomatosis. In typical instances, differentiation is easy, but occasionally one is unable to make any more specific diagnosis in these diseases than papilloma. Because the term papilloma is often used in such non




FIG 123 Comparison of epidermal cyst (left) with sebaceous cyst (right). The wall of the epidermal cyst is filled with

amorphous material which has formed by the gradual disintegration of the vacuolated cells ( $\times 200$ )

specific sense it is better to avoid this term as a designation for nevus verrucosus

In typical instances keratosis verrucosus differs from nevus verrucosus by its downward growth. In the verrucosus downward growth shows no

The degree of hyperkeratosis and papillomatosis differs considerably from lesion to lesion and depends on the size of the lesion. The thickness of the granular layer varies, areas of marked hyperplasia of the granular layer may alternate with areas in which it is atrophic



FIG. 151 Epidermal cyst. The wall is composed of true epidermis, i.e. squamous granular and horn cells. The cyst is filled with keratin. ( $\times 100$ )

There may be some proliferation of basal cells (as seen in basal-cell papilloma) and hyperpigmentation of the basal layer. In rare instances apocrine gland lumina are found deep in the dermis (see page 346). Nevus cells are absent in pure nevus verrucosus. However, the combination of nevus verrucosus and nevus pigmentosus is not infrequent, and in that case nevus cells are present.

**Differential Diagnosis.** Nevus verrucosus must be differentiated from other types of papillomas, namely senile keratosis (keratosis senilis), basal cell papilloma, verruca vulgaris and acanthosis nigra.

SEBACEOUS CYSTS, or steatomas, have a wall composed of epithelial cells that possess no intercellular bridges and do not undergo keratinization. Many of the cells are vacuolated because of their transformation into sebaceous cells (Fig. 155). The wall never possesses



FIG. 156. Calcified sebaceous cyst. The palisading of the basal layer makes it evident that this is a sebaceous cyst. It has ruptured and fibrous tissue has proliferated into the lumen ( $\times 100$ ).

rete ridges at its periphery. The peripheral cell layer shows a basal layer of cells.

Large amounts of cholesterol and cholesterol crystals may be present in the cysts. Calcification occurs frequently within sebaceous cysts (Fig. 156). As in epidermal cysts, a considerable foreign body reaction results when the wall of a sebaceous cyst ruptures, and the cyst may undergo partial disinte-

203) *Verruca Vulgaris* shows groups of large vacuolated cells in the upper stratum malpighii and the granular layer, and intermittent areas of parakeratosis (Figs 118, 119). *Acanthosis nigricans*, as a rule, shows less acanthosis than *nexus verrucosus* and atrophy of the rete ridges rather than elongation (Fig 139).

## EPIDERMAL CYST (WEN), SEBACEOUS CYST, MILIUM, DERMOID CYST

Epidermal cysts and sebaceous cysts are often indistinguishable clinically. Both are commonly called wens. On histologic examination, most wens prove to be epidermal rather than sebaceous cysts. In their material, Wary and Gates found 556 epidermal cysts as compared with only 3 sebaceous cysts.

Epidermal cysts are slow growing, elevated, round, firm, intracutaneous or subcutaneous tumors, varying from 0.2 to 5 cm. in diameter. They occur most commonly on the scalp. As a rule, they are freely movable with the skin. No orifice can be demonstrated in the overlying skin. The material within them is nearly solid and is odorless.

Sebaceous cysts often cannot be differentiated from epidermal cysts on a clinical basis. In general, however, sebaceous cysts are softer than epidermal cysts, and, occasionally, they show a small orifice on their surface. The material within them is semifluctuant and has a rancid odor.

Milia are multiple pinhead sized, whitish, globoid, hard lesions occurring most commonly on the face.

Dermoid cysts occur in rare instances in the subcutaneous tissue as soft, round or oval tumors of various sizes. The most frequently observed site is the periorbital region.

**Histopathology** EPIDERMAL CYSTS have a wall composed of true epidermis, i.e., squamous, granular and horn cells (Fig 154). Rete ridges may be present at the periphery of the wall. The cyst is filled with keratin which frequently is arranged in laminated layers. Foci of calcification are found only in rare instances. When the cyst ruptures and the contents of the cyst reach the dermis, a considerable foreign body giant cell reaction results. The foreign body reaction may cause breaking up and partial disintegration of the epidermal cyst.

Malignant degeneration is rare, occurring in approximately 1.5 per cent of cases (Love and Montgomery). If such degeneration occurs, it takes the form of squamous cell carcinoma. Usually, such carcinomas are of low grade malignancy and do not cause metastases (Peden).

SEBACEOUS CYSTS, or steatomas, have a wall composed of epithelial cells that possess no intercellular bridges and do not undergo keratinization. Many of the cells are vacuolated because of their transformation into sebaceous cells (Fig 155). The wall never possesses



FIG 156 Calcified sebaceous cyst. The palisading of the basal layer makes it evident that this is a sebaceous cyst. It has ruptured and fibrous tissue has proliferated into the lumen ( $\times 109$ ).

rete ridges at its periphery. The peripheral cell layer shows a distinct palisade arrangement such as is never seen in epidermal cysts. The cysts are filled with amorphous material which forms by the gradual disintegration of the sebaceous cells. Large amounts of cholesterol and cholesterol crystals may be present in the cysts. Calcification occurs frequently within sebaceous cysts (Fig 156). As in epidermal cysts a considerable foreign body reaction results when the wall of a sebaceous cyst ruptures, and the cyst may undergo partial disinte-



gration. In rare instances, basal-cell epitheliomas have been found arising in sebaceous cysts (Love and Montgomery).

MILIA present a histologic aspect similar to that of epidermal cysts. They are, however, much smaller. Milia are not tumors, like the epidermal and the sebaceous cysts, but are retention cysts caused by the occlusion of a pilosebaceous follicle (Love and Montgomery).

DERMOID CYSTS are lined by an epidermis endowed with rudimentary sebaceous glands, sweat glands and hair follicles. They contain sebaceous material as well as keratin. In addition, hairs are present in about 30 per cent of cutaneous dermoid cysts. Cartilage and bone are encountered occasionally (New and Erich).

Differential Diagnosis. For differentiation of epidermal and sebaceous cysts and, particularly, of sebaceous cyst with secondary basal cell epithelioma from calcifying epithelioma, see page 368.

### KERATOSIS SENILIS

Keratosis senilis occurs, frequently as multiple lesions, on the face and the dorsa of the hands in persons past middle life. The lesions



FIG. 157 Keratosis senilis. There are hyperkeratosis and papilloma

tion, and shows a rather pronounced chronic inflammatory infiltrate ( $\times 100$ )

usually measure less than 1 cm in diameter and show dry, hard scales firmly adherent to an erythematous base showing little or no infiltration. Occasionally lesions of senile keratosis show a verrucous surface. In from 20 to 25 per cent of the cases of senile keratosis squamous cell carcinoma develops in one or more of the lesions (Montgomery and Dorfman).




Fig. 158 Keratosis senilis. Clefts and two vesicles are present in the epidermis in the clefts the dermis as (X150)

*Cornu cutaneum*, a variant of keratosis senilis, shows a circumscribed horny excrescence sometimes suggesting the horn of an animal.

**Histopathology.** The shortest histologic definition that may be given to keratosis senilis is that it represents a squamous-cell carcinoma Grade  $1\frac{1}{2}$ . It is a precancerous lesion differing only in degree from squamous-cell carcinoma or Bowen's disease.

Hyperkeratosis is usually pronounced. Mild or moderate papillomatosis may be present. The stratum malpighi varies in thickness, may alternate. The stratum of the rete cells. In some of cells, individual cell

dyskeratosis, clumping of nuclei and prevalence of mitotic figures are observed, so that the histologic picture approaches that of Bowen's disease (Szodora). In other cases, the epidermis shows irregular downward proliferation but without frank invasion (Fig 157). The histologic picture then approaches that of squamous cell carcinoma Grade I.

Not infrequently, immediately above the basal layer, one observes clefts similar to those seen in Darier's disease (see page 48). These clefts, first described by Freudenthal, may contain a few acantholytic cells. In rare instances, even small suprabasal vesicles may be present. The basal layer underneath these clefts and vesicles usually consists of cells with large, hyperchromatic nuclei which appear closely set together. In addition, this type of atypical basal layer may proliferate into the dermis as short ductlike structures and, furthermore, it may surround as cell mantles the upper portion of pilosebaceous follicles and sweat ducts the epithelium of which otherwise appears normal (Fig 158) (Halter).

The upper dermis usually shows a fairly dense, chronic, inflammatory infiltrate in which plasma cells are prominent.

In instances in which a diagnosis of senile keratosis is made, it often is advisable to section deeper into the block of tissue, because actual progression into squamous cell carcinoma may have taken place in another area.

CORNÜ CUTANEUM is a type of keratosis senilis with particularly pronounced hyperkeratosis.

Differential diagnosis

from basaloid tumor of cells of the basal cell type (see page 382) (Freudenthal, Montgomery and Dorf, Rutter). For differentiation from squamous cell carcinoma see page 334. From arsenical keratosis see page 155.

## LEUKOPLAKIA

Leukoplakia represents senile keratosis of the mucous membranes and occurs on the lips, the oral mucosa and the vulva. On the vulva leukoplakia is apt to develop secondary to senile atrophy of the vulva (see page 162).

Clinically, the lesions consist of white patches which are sharply demarcated, irregular in outline and slightly elevated above the level of the mucous surface. The lesions are often multiple and may be discrete or confluent. In from 20 to 30 per cent of the cases of leukoplakia squamous cell carcinoma supervenes (MacKee and Cipollaro).

**Histopathology.** In leukoplakia, hyperkeratosis usually is less pronounced than in keratosis senilis, but the inflammatory infiltrate is

often more intense and may hug the epidermis just as in lichen planus. Formation of suprabasal clefts as in senile keratosis does not occur.

**Differential Diagnosis** Differentiation of leukoplakia from lichen planus may be very difficult and occasionally impossible. As a rule however the infiltrate in leukoplakia is less severe than in lichen planus and contains a rather large number of plasma cells whereas in lichen planus their number is small. In both diseases the epidermal-dermal border is apt to have a hazy outline due to the invasion of the lower epidermis by the inflammatory infiltrate but only leukoplakia shows atypicality of the cells in the stratum malpighii. On the buccal mucosa pressure of teeth or dentures may produce pressure calluses which show hyperkeratosis, acanthosis and a non specific inflammatory infiltrate. They differ from leukoplakia by the absence of atypicality in the epidermis.

### SQUAMOUS CELL CARCINOMA (EPIDERMOID CARCINOMA)

Squamous cell carcinoma may occur anywhere on the skin as well as on the mucous membranes. It may begin as such or develop from a senile keratosis or leukoplakia. Most commonly the lesion consists of a shallow ulcer surrounded by a wide elevated and indurated border. The ulcer often is covered by a crust which conceals a red granular base. Occasionally raised fungoid verrucous lesions with an ulceration occur. The latter are usually of a relatively low grade of malignancy whereas ulcerated lesions may grow rapidly and cause metastases within a short time.

**Histopathology** Squamous-cell carcinoma is a true invasive carcinoma. The tumor cells are found in nests and cords which invade the dermis. The invading tumor masses are composed of squamous and squamoid cells. The cells show a great variation in the size and the shape of the cells, hyperplasia and hyperchromasia of the nuclei, absence of prickles, keratinization of individual cells, prevalence of mitotic figures and presence of atypical mitotic figures.

Differentiation in squamous-cell carcinomas is in the direction of keratinization. Keratinization often takes place in the form of horn pearls. The horn pearls are very characteristic and consist of concentric layers of squamous

keratinization toward the center. The center may or may not show complete keratinization.

Broders introduced a system of grading squamous cell carcinoma. He established four grades according to the proportion of differentiated cells to atypical cells. In Grade I, more than 75 per cent, in Grade II, more than 50 per cent, in Grade III, more than 25 per cent, and in Grade IV, less than 25 per cent of the cells are differentiated. Since differentiation is in the direction of keratinization, the degree of keratinization is a good guide in grading. Broders' system of grading is most useful and has been accepted widely in spite of certain objections raised against it. In the first place, there is, of course, a large personal factor in the interpretation of cytologic changes, and, secondly, different degrees of malignancy may be present in different fields. In regard to the first objection, Broders has suggested to err rather on the side of the higher grade because malignant processes tend to be progressive rather than regressive. In regard to the second objection, one should examine several sections of every tumor and grade according to the least differentiated portion (Edmundson).

In squamous cell carcinoma, Grade I (Fig. 159), the tumor masses have not penetrated beyond the level of the sweat glands. They still show in some areas an intact basal layer at their periphery. In other areas the basal layer has become disorganized and has disappeared. In such areas the cell masses appear poorly demarcated from the surrounding stroma. The cells of the invading cell masses are predominantly mature squamous cells with well developed prickles. Nevertheless some of the squamous cells are atypical. Horn pearls are present in fairly large number. Some are well developed and have fully keratinized centers; others, however, show only partial keratinization of their centers and the concentric arrangement of the cells is not distinct. Besides horn pearls, sheets of partially keratinized cells may be present. The dermis often shows a rather marked inflammatory reaction. It is noteworthy that, in simple keratosis and squamous cell carcinoma, Grade I, the inflammatory reaction in the dermis is usually much more pronounced than in the more malignant forms of squamous cell carcinoma. This phenomenon is due to the fact that tissue when invaded by carcinomatous cells is able to defend itself to some extent provided the cells are only moderately malignant, but is overwhelmed without any fight if the cells are highly malignant. (The same observation incidentally can be made also in malignant melanoma and mycosis fungoides—see pages 460 and 485.) Metastases do not as a rule occur as long as a squamous-cell carcinoma remains Grade I.

In squamous-cell carcinoma, Grade II (Fig 160), the invading cell masses are, as a rule, poorly demarcated from the surrounding stroma. They may invade deeply. Keratinization is much less in evidence than in Grade I. There are only few horn pearls, and those present show incompletely keratinized centers. A fairly large number of the squamous cells are atypical.



FIG. 159 Squamous-cell carcinoma, Grade I. There is invasion of the dermis by epidermal masses. The cells of the invading epidermal masses are predominantly mature squamous cells showing relatively slight atypicality. Several horn pearls are present. The dermis shows a marked inflammatory reaction ( $\times 100$ ).

In squamous-cell carcinoma, Grade III (Fig 161), keratinization is minimal. Horn pearls are not found. Instead, keratinization occurs in small cell groups and in individual cells (individual cell keratinization, "malignant dyskeratosis" see page 499). The majority of cells are atypical. Mitotic figures are conspicuous and often atypical.

In squamous-cell carcinoma, Grade IV (Fig 162), keratinization is almost completely absent. Nearly all squamous cells are atypical and devoid of prickles. Thus, it is often difficult to arrive at the correct

keratinization toward the center. The center may or may not show complete keratinization.

Broders introduced a system of grading squamous cell carcinoma. He established four grades according to the proportion of differentiated cells to atypical cells. In Grade I, more than 75 per cent, in Grade II, more than 50 per cent, in Grade III, more than 25 per cent, and in Grade IV, less than 25 per cent of the cells are differentiated. Since differentiation is in the direction of keratinization, the degree of keratinization is a good guide in grading. Broders' system of grading is most useful and has been accepted widely in spite of certain objections raised against it. In the first place, there is, of course, a large personal factor in the interpretation of cytologic changes, and, secondly, different degrees of malignancy may be present in different fields. In regard to the first objection, Broders has suggested to err rather on the side of the higher grade because malignant processes tend to be progressive rather than regressive. In regard to the second objection, one should examine several sections of every tumor and grade according to the least differentiated portion (Edmundson).

In squamous cell carcinoma, Grade I (Fig. 159), the tumor masses have not penetrated beyond the level of the sweat glands. They still show in some areas an intact basal layer at their periphery. In other areas the basal layer has become disorganized and has disappeared. In such areas, the cell masses appear poorly demarcated from the surrounding stroma. The cells of the invading cell masses are predominantly mature squamous cells with well developed prickles. Nevertheless, some of the squamous cells are atypical. Horn pearls are present in fairly large number. Some are well developed and have fully keratinized centers, others, however, show only partial keratinization of their centers and the concentric arrangement of the cells is not distinct. Besides horn pearls, sheets of partially keratinized cells may be present. The dermis often shows a rather marked inflammatory reaction. It is noteworthy that, in senile keratosis and squamous cell carcinoma, Grade I, the inflammatory reaction in the dermis is usually much more pronounced than in the more malignant forms of squamous cell carcinoma. This phenomenon is due to the fact that tissue, when invaded by carcinomatous cells, is able to defend itself to some extent provided the cells are only moderately malignant, but is overwhelmed without any fight if the cells are highly malignant. (The same observation incidentally, can be made also in malignant melanoma and mycosis fungoides—see pages 460 and 485.) Metastases do not, as a rule, occur as long as a squamous cell carcinoma remains Grade I.

In squamous-cell carcinoma, Grade II (Fig. 160), the invading cell masses are, as a rule, poorly demarcated from the surrounding stroma. They may invade deeply. Keratinization is much less in evidence than in Grade I. There are only few horn pearls, and those present show incompletely keratinized centers. A fairly large number of the squamous cells are atypical.



FIG. 159 Squamous-cell carcinoma, Grade I. There is invasion of the dermis by epidermal masses. The cells of the invading epidermal masses are predominantly mature squamous cells showing relatively slight atypicality. Several horn pearls are present. The dermis shows a marked inflammatory reaction ( $\times 100$ ).

In squamous-cell carcinoma, Grade III (Fig. 161), keratinization is minimal. Horn pearls are not found. Instead, keratinization occurs in small cell groups and in individual cells ("individual cell keratinization" = "malignant dyskeratosis," see page 499). The majority of cells are atypical. Mitotic figures are conspicuous and often atypical.

In squamous-cell carcinoma, Grade IV (Fig. 162), keratinization is almost completely absent. Nearly all squamous cells are atypical and devoid of prickles. Thus, it is often difficult to arrive at the correct






FIG 160 Squamous-cell carcinoma Grade II The cell masses show much less keratinization than in Grade I There are only few horn pearls and those present show incompletely keratinized centers Atypical cells are conspicuous ( $\times 200$ )




FIG 161 Squamous-cell carcinoma Grade III No horn pearls are present Keratinization occurs only in small cell groups Many cells are atypical and devoid of prickles To the right a cell shows individual cell keratinization (I K) ( $\times 200$ )

diagnosis as long as individual fields only are studied. The tumor may suggest a malignant melanoma in some cases and a sarcoma in others. The latter diagnosis may be particularly difficult to rule out when the cells, as occasionally occurs, are spindle shaped ('spindle-cell' squamous cell carcinoma) (Brooks, Underwood, Montgomery and Broders). If, however, sections are subjected to a thorough study, the type of origin from the epidermis and the presence in a few areas



FIG 162 Squamous-cell carcinoma, Grade IV. There is no evidence of keratinization. The epithelial cells appear atypical, are devoid of prickles and elongated so that the tumor suggests a sarcoma almost more than a carcinoma. There are numerous mitotic figures (X200).

of cells showing prickles or partial keratinization usually establish the diagnosis. Squamous-cell carcinoma, Grade IV, is relatively rare in the skin. Many of the reported instances of 'spindle-cell' squamous-cell carcinoma occurred in areas of radiodermatitis. In this connection, it may be pointed out that it is not yet established fully whether or not sarcomas can develop following radiodermatitis. Most, if not all, cases reported as such in the literature represent squamous cell carcinomas of the spindle-cell type (Sims and Kirsch, Gentile) (see page 122).

Metastases are rare in squamous-cell carcinoma, Grade I, but common in the other grades. The regional lymph nodes are the first site to be invaded by metastases.

**Differential Diagnosis.** The diagnosis of squamous-cell carcinoma, although easily made in typical cases, may be difficult at times

Squamous cell carcinoma must be differentiated from senile keratosis, pseudo epitheliomatous hyperplasia and basal cell epithelioma.

The differences between squamous cell carcinoma and senile keratosis lie in the degree rather than in the type of changes. In both conditions, one finds atypicality of cells with dyskeratosis of individual cells and downward proliferation of the epidermis. However, in squamous cell carcinoma these changes are more severe, and, in addition, horn pearl formation and actual invasion of the dermis are present. No sharp line of separation exists between the two conditions, and it is not infrequent to find in a lesion which in general has the appearance of senile keratosis, on serial sections, one or several areas in which the changes have progressed to squamous cell carcinoma.

For differentiation from pseudo epitheliomatous hyperplasia, see below. For differentiation from basal cell epithelioma, see page 380.

#### PSEUDO EPITHELIOMATOUS HYPERPLASIA

Considerable thickening and irregular proliferation of the skin, which clinically, as well as histologically, may suggest carcinoma, occurs not infrequently in chronic granulomas, such as bromoderma, blastomycosis and granuloma inguinale, and at the edges of chronic ulcers, such as occur after burns, in stasis dermatitis, basal cell epithelioma, lupus vulgaris, scrofuloderma, gumma and pyoderma gangrenosum. In addition, granular cell myoblastoma is known to evoke quite frequently a pseudo epitheliomatous hyperplasia.

**Histopathology.** Histologically, one observes an epithelial hyperplasia which may closely resemble squamous cell carcinoma, Grades I or II, and therefore is referred to as pseudo-epitheliomatous hyperplasia. Although squamous cell carcinoma may develop at the edges of chronic ulcers, it is likely that some of the cases that have been regarded as such in the past were in reality pseudo epitheliomatous hyperplasia.

The histologic picture of pseudo epitheliomatous hyperplasia shows irregular invasion of the dermis by epidermal cell masses with horn pearl formation and often numerous mitotic figures (Fig. 163). The penetration may extend even below the level of the sweat glands as isolated fragments of epidermal tissue (Sommerville). However the squamous cells usually are well differentiated, and atypicalities such as individual cell keratinization and nuclear hyperplasia and hyperchromasia are absent. Furthermore in pseudo epitheliomatous hyperplasia, there often are invasion of the epithelial proliferations by leukocytes and disintegration of some of the epidermal cells, a phe-

nomenon usually not seen in squamous-cell carcinoma (Winer; Montgomery) But even when all these criteria are taken into account, it may still be difficult to differentiate between squamous cell carci-



FIG. 103 Pseudo-epitheliomatous hyperplasia in bromoderma. There is downward proliferation of the epidermis analogous to squamous-cell carcinoma, Grade I. In the field shown it is impossible to rule out carcinoma. Note, however, the permeation of the epidermis in many areas by inflammatory cells ( $\times 100$ ).

nomas and

log

diff

It is worth remembering to study the inflammation in every section in which one carcinoma, Grades I or II, or the granulomatous myc.

... such evidence is found, one may

be dealing with pseudo epitheliomatous hyperplasia rather than with squamous cell carcinoma

### "MULTIPLE, PRIMARY, SELF HEALING SQUAMOUS CELL CARCINOMATA"

In this disorder, which often is familial, there is a continuous appearance, especially on the face and the extremities, of papules which grow into nodules, ulcerate and, after a few months, heal with a depressed scar.

**Histopathology.** The histologic appearance is like that of a well differentiated, keratinizing squamous cell carcinoma, Grade I. A dense inflammatory infiltrate is present. There may or may not be ulceration, depending on the stage of the lesion.

It is generally agreed that differentiation from squamous cell carcinoma is impossible on a histologic basis and depends on the clinical data (Smith, Sommerville and Milne Witten and Zak)

The histogenesis is not clear. However, it is certain that the epidermal proliferation represents a pseudo epitheliomatous hyperplasia. Whimster assumes that all the downward prolongations of epithelium can be explained as extreme hyperplasia, probably in inflammatory origin, of sweat ducts and hair follicles.

### BOWEN'S DISEASE

Bowen's disease usually manifests itself as a single lesion. It is characterized by a dull red patch of sharp but irregular outline, showing little or no infiltration. Within the patch, there usually are areas of crusting beneath which one finds a granular and oozing surface. The patch slowly spreads by peripheral extension and shows no tendency to healing in its center.

**Histopathology.** Bowen's disease is an intra epidermal, squamous cell carcinoma, or a squamous cell carcinoma in situ, and not a "pre cancerous dermatosis," under which title it was described originally by Bowen.

The epidermis shows hyperkeratosis with parakeratosis and acanthosis. The rete ridges are elongated and thickened, often to such a degree that the papillae located between them are reduced to thin strands or are obliterated. However, the basal layer is intact, and no true invasion can be seen. Throughout the stratum malpighii, the cells lie in complete disorder; many are atypical, showing large and hyperchromatic nuclei. Multinucleated epidermal cells containing clusters of nuclei are common (Fig. 164). Some cells may show marked vacuolization simulating Paget cells. However, the intercellular




FIG 164 Bowen's disease The epidermis is thickened the basal layer is intact The cells of the stratum malpighii lie in complete disorder and many of them are atypical showing large and hyperchromatic nuclei Several multinucleated cells with clumped nuclei and numerous mitotic figures are present in the stratum malpighii ( $\times 200$ )




FIG 165 Bowen's disease In addition to the changes, as described for Figure 164 there are within the stratum malpighii three cells showing individual cell keratinization (IK) and one horn pearl (HP) ( $\times 200$ )

bridges of these cells are preserved, in contrast with Paget cells, in which they are absent

A common and if present, characteristic finding is the presence of individual cell keratinization in the stratum malpighii (Fig 165). Such keratinized cells are large and round, have a homogeneous and strongly eosinophilic cytoplasm and a large, irregularly shaped, hyperchromatic nucleus (This individual cell keratinization, which may occur not only in Bowen's disease but also in squamous cell carcinoma, is often referred to as malignant dyskeratosis in contrast with the benign dyskeratosis that underlies the formation of corps ronds and grains in Darier's disease). Occasionally, actual horn pearls may occur within the epidermis.

The upper dermis usually shows a moderate amount of inflammatory infiltration composed chiefly of lymphocytes and plasma cells.

As already stated, in true Bowen's disease the basal layer is intact. However, in some cases the basal layer ultimately is broken through and a true invasive squamous cell carcinoma results. This may occur at first in only one or a few areas. In order not to miss such areas it is advisable to examine representative sections throughout the entire tissue block. As soon as the invasion of the dermis occurs, the prognosis changes. As long as Bowen's disease remains in the true, intraepidermal stage metastases do not occur. However, when invasion of the dermis has occurred, the likelihood of metastases is rather great. This is due to the fact that if Bowen's disease changes into an invasive carcinoma, it usually is Grade II or even Grade III with considerable atypicality of the cells and little tendency to keratinization (Kuznitsky and Jacoby).

**Differential Diagnosis.** In the differential diagnosis senile keratosis and arsenical keratosis must be considered. Senile keratosis may resemble Bowen's disease closely but shows less atypicality of the squamous cells. No sharp line of distinction can be drawn between the two conditions. Arsenical keratosis may closely resemble either Bowen's disease or senile keratosis but shows, as a rule, more vacuolization of the squamous cells than these two diseases.

#### ERYTHROPLASIA OF QUEYRAT

Erythroplasia of Queyrat usually occurs on the glans penis but may be seen occasionally on the prepuce, the vulva or the oral mucosa. The lesion usually single, consists of a well defined area with a brilliant red, velvety surface and little or no infiltration.

**Histopathology.** Erythroplasia represents an intraepithelial squamous cell carcinoma of the mucous membranes and as such is analogous to Bowen's disease of the skin (Pautrier). Progression into

invading squamous-cell carcinoma usually occurs sooner than with Bowen's disease of the skin (Sulzberger and Sitenstein McDaniel and Mason Pautrier)

**Differential Diagnosis** Sachs and Sachs as well as Zoon recently have described cases in which the clinical appearance was identical with that of erythroplasia of Queyrat but in which histologic examination revealed no malignant changes in the epidermis. Instead the dermis contained an inflammatory infiltrate composed predominantly of plasma cells. Zoon has suggested the diagnostic term *balanoposthitis chronica circumscripta* for these cases. Because the two diseases are indistinguishable on clinical grounds it is evident that a diagnosis of erythroplasia of Queyrat always requires histologic confirmation.

## PAGET'S DISEASE

Paget's disease of the nipple occurs as a rule on and around the nipples of women; a few instances of its occurrence on the male breast have been described (Rubenstein). The lesion is always unilateral. Extramammary Paget's disease, which is uncommon, occurs on or near the male and the female genitals in the perianal region and in the axillae.

The lesion of Paget's disease consists of a sharply defined, slightly infiltrated area of dusky erythema showing scaling, oozing and crusting. If located on the breast, the process begins in the nipple or the areola of the breast and slowly extends to the surrounding skin. There may or may not be retraction of the nipple.

**Histopathology** For many years Paget's disease of the nipple was thought to begin in the skin as a precancerous lesion that later became malignant and then invaded the mammary gland. It is now generally accepted that Paget's disease of the nipple is a cancer from the outset and that the initial lesion is a carcinoma *in situ* arising in one or more mammary ducts near their outlets. The primary duct cancer extends from the site of origin downward to the epithelium lining the acini and upward and outward to the epidermis where it causes the cutaneous lesion. Thus the tumor cells present in the epidermis, often referred to as Paget cells, are ductal and not epidermal cells (Pautrier, Fraser, Muir, Inglis). At a later stage the cancer breaks through the wall of a duct or acinus and infiltrates the connective tissue of the breast.

A carcinoma has been found in the mammary ducts in nearly all cases of Paget's disease and in the mammary glands in most cases. In the few cases where no carcinoma was found in the breast, the underlying ductal carcinoma may have been overlooked because in the



bridges of these cells are preserved, in contrast with Paget cells, in which they are absent

A common and, if present, characteristic finding is the presence of individual cell keratinization in the stratum malpighii (Fig 165). Such keratinized cells are large and round, have a homogeneous and strongly eosinophilic cytoplasm and a large, irregularly shaped, hyperchromatic nucleus (This individual cell keratinization, which may occur not only in Bowen's disease but also in squamous cell carcinoma, is often referred to as malignant dyskeratosis in contrast with the benign dyskeratosis that underlies the formation of corps ronds and grains in Darier's disease). Occasionally, actual horn pearls may occur within the epidermis.

The upper dermis usually shows a moderate amount of inflammatory infiltration composed chiefly of lymphocytes and plasma cells.

As already stated, in true Bowen's disease the basal layer is intact. However, in some cases the basal layer ultimately is broken through and a true invasive squamous cell carcinoma results. This may occur at first in only one or a few areas. In order not to miss such areas it is advisable to examine representative sections throughout the entire tissue block. As soon as the invasion of the dermis occurs the prognosis changes. As long as Bowen's disease remains in the true, intraepidermal stage, metastases do not occur. However, when invasion of the dermis has occurred, the likelihood of metastases is rather great. This is due to the fact that if Bowen's disease changes into an invasive carcinoma, it usually is Grade II or even Grade III, with considerable atypicality of the cells and little tendency to keratinization (Kuznitzky and Jacoby).

**Differential Diagnosis.** In the differential diagnosis, senile keratosis and arsenical keratosis must be considered. Senile keratosis may resemble Bowen's disease closely but shows less atypicality of the squamous cells. No sharp line of distinction can be drawn between the two conditions. Arsenical keratosis may closely resemble either Bowen's disease or senile keratosis but shows, as a rule, more vacuolization of the squamous cells than these two diseases.

#### ERYTHROPLASIA OF QUEYRAT

Erythroplasia of Queyrat usually occurs on the glans penis but may be seen occasionally on the prepuce, the vulva or the oral mucosa. The lesion usually single consists of a well defined area with a brilliant red, velvety surface and little or no infiltration.

**Histopathology.** Erythroplasia represents an intraepithelial squamous cell carcinoma of the mucous membranes and as such is analogous to Bowen's disease of the skin (Pautrier). Progression into



Only a  
re large



FIG 168 Paget's disease of the nipple. Intraductal carcinoma is present in the mammary ducts. The carcinoma is confined within the walls of the ducts (X200)

ducts the carcinoma may undergo considerable regression with resultant sclerosis and calcification of the involved ducts (Marr)

Histologic examination of the epidermis reveals in early lesions acanthosis with elongation of the rete ridges, and in older lesions




FIG 166 Paget's disease of the nipple Low magnification The epidermis is permeated with numerous Paget cells lying singly and in groups Note that there is no invasion of the dermis by Paget cells An inflammatory infiltrate is present in the dermis ( $\times 200$ )

thinning and flattening. Paget cells are scattered through the epidermis (Fig 166). They are large, devoid of prickles and at times appear to lie in clear spaces (Fig 167). Their cytoplasm stains much lighter than that of the adjacent squamous cells. Their nuclei are large, round and pale staining. Paget cells usually are most numerous in the basal layer and may cause disorganization of this layer. Invasion of the dermis from the epidermis, however, does not occur.

## 2 TUMORS OF THE EPIDERMAL APPENDAGES

## NEVUS SEBACEUS (JADASSOHN)

Nevus sebaceus is located most commonly on the scalp or on the face as a single lesion present since birth. It consists of a circum



FIG 169 Nevus sebaceus (Jadassohn) There are hyperkeratosis and papillomatosis. Numerous mature sebaceous glands lie in the upper dermis. In the lower dermis mature apocrine glands are located (X50)

scribed slightly raised firm yellow plaque with a smooth though furrowed surface

**Histopathology** The tumor is composed of large numbers of many or many mature sebaceous glands. The overlying epidermis may be thickened. Frequently apocrine glands are present in the dermis beneath the masses of sebaceous gland lobules (Fig 169). The presence of abortive hair follicles has also been noted on several occasions.

In some cases, Paget cells are so numerous throughout the epidermis that the normal squamous cells show signs of injury. They are compressed and deformed and may form only a network, the meshes of which are filled with Paget cells (Fig 166)

The dermis shows in Paget's disease a moderately severe chronic inflammatory reaction

Histologic examination of the mammary ducts nearly always shows malignant changes in some of them. At first, the carcinoma is intraductal and the tumor cells are confined within the walls of the ducts (so called comedo carcinoma) (Fig 168). Ultimately, the tumor cells invade the connective tissue. From then on, lymphatic spread and metastases occur.

**EXTRAMAMMARY PAGET'S DISEASE**, as a rule, presents the same epidermal changes as Paget's disease of the nipple, although in lesions located in the perianal region the Paget cells often contain a considerable amount of mucin. In most reported cases, an underlying carcinoma of apocrine ducts and glands has been found (Parsons, Foraker and Miller, Zoon and Gelpke). This is in accordance with the fact that the mammary gland is a modified apocrine gland. The presence of mucin within the Paget cells of some of the cases located in the perianal region may be the result of mucous metaplasia of apocrine glands, but Whimster suggests that the primary cancer may be one of ectopic mucous glands rather than of apocrine glands.

**Differential Diagnosis.** Paget's disease of the nipple must be differentiated from Bowen's disease. In Bowen's disease, large, vacuolated epidermal cells may also occur, but, in contrast with the Paget cells, they often possess prickles. Furthermore, one observes clumping of nuclei within multinucleated epithelial giant cells and individual cell keratinization in Bowen's disease but never in Paget's disease. In cases in which the Paget cells are concentrated in the lower epidermis, the resemblance to an amelanotic junction nevus or an early malignant melanoma may be very great because in both diseases the characteristic cell is large and vacuolated (Stout). Allen believes that many cases of Paget's disease and especially of extramammary Paget's disease are wrongly thus diagnosed and in reality are junction nevi. The most important points of differentiation are first, the absence of invasion of the dermis by the tumor cells in Paget's disease, and second, the presence of minute amounts of melanin in at least some tumor cells even in apparently amelanotic junction nevi and malignant melanomas. The melanin is best demonstrated by the use of a silver stain. Carrying out of the dopa stain (see page 12) also aids in the differentiation.

## 2 TUMORS OF THE EPIDERMAL APPENDAGES

### NEVUS SEBACEUS (JADASSOHN)

Nevus sebaceus is located most commonly on the scalp or on the face as a single lesion present since birth. It consists of a circum



FIG 169 Nevus sebaceus (Jadassohn) There are hyperkeratosis and papillomatosis. Numerous mature sebaceous glands i.e. in the upper dermis. In the lower dermis mature apocrine glands are located (X50).

scribed slightly raised firm yellow plaque with a smooth though furrowed surface.

**Histopathology** The tumor type or nearly mature sebaceous or may not show hyperkeratosis. Apocrine glands have been described as occurring in nevus sebaceus (Robinson, Koch, Pantner). They are located deep in the dermis beneath the masses of sebaceous gland lobules (Fig 169). The presence of abortive hair follicles has also been noted on several occasions.

(Ostrowski). Koch and Pautrier believe that the frequent presence of apocrine glands is evidence that nevus sebaceus develops from primary epithelial germs (For a discussion of the primary epithelial germ, see pages 4 and 318.)

The presence of a basal cell epithelioma within a nevus sebaceus is not uncommon (Fig. 170) (Ostrowski; Robinson; Szodoray, Pautrier,



FIG. 170 Nevus sebaceus (Jadassohn) within which a basal cell epithelioma has arisen. The nevus sebaceus is on the right side, the basal cell epithelioma on the left ( $\times 25$ )

Savatard) In rare instances, a squamous-cell carcinoma may develop from the epidermis overlying the nevus sebaceus (Parkin).

### ADENOMA SEBACEUM (PRINGLE) (TUBEROUS SCLEROSIS)

Adenoma sebaceum (Pringle) usually represents only part of a widely disseminated tissue malformation which results in the formation of tumors in both ectodermic and mesodermic structures. Tumors are found, aside from the skin, notably in the brain, the retina, the kidney and the heart. The tumors in the brain, which are referred to as tuberous sclerosis, are gliomas. Those of the retina have the same histologic appearance as the brain tumors (Kveim). The tumors of the heart usually are rhabdomyomas (Pratt-Thomas). Those of the kidney may be angiomas, fibromas, adenomas or mixed tumors; they may become malignant (Butterworth and Wilson).

The cutaneous lesions of adenoma sebaceum (Pringle) occur on the face, particularly in the malar region. They consist of numerous small papules and nodules which are yellowish brown in color and frequently show telangiectases on their surface.




FIG 171 Adenoma sebaceum (Pringle). Instead of showing excessive development of sebaceous glands, as is usually the case, this lesion shows a large number of immature hair structures ( $\times 50$ )

**Histopathology.** The term adenoma sebaceum is a misnomer. Since the sebaceous structures are not adenomatous but fully or almost fully developed and abnormal merely by their presence in excessive number, the lesion represents an organic nevus, rather than an adenoma (see Table I).

In addition to the  
findings in the

Usually, a large number of hair structures are found. Usually, those hair structures are somewhat immature (Fig 171) (Butterworth and Wilson, Good and Garb)



(Ostrowski). Koch and Pautrier believe that the frequent presence of apocrine glands is evidence that nevus sebaceus develops from primary epithelial germs. (For a discussion of the primary epithelial germ, see pages 4 and 318.)

The presence of a basal-cell epithelioma within a nevus sebaceus is not uncommon (Fig. 170) (Ostrowski; Robinson; Szodoray, Pautrier;



FIG. 170. Nevus sebaceus (Jadassohn) within which a basal cell epithelioma has arisen. The nevus sebaceus is on the right side, the basal cell epithelioma on the left ( $\times 25$ )

Savataud) In rare instances, a squamous cell carcinoma may develop from the epidermis overlying the nevus sebaceus (Parkin).

### ADENOMA SEBACEUM (PRINGLE) (TUBEROUS SCLEROSIS)

Adenoma sebaceum (Pringle) usually represents only part of a widely disseminated tissue malformation which results in the formation of tumors in both ectodermic and mesodermic structures. Tumors are found, aside from the skin, notably in the brain, the retina, the kidney and the heart. The tumors in the brain, which are referred to as tuberous sclerosis, are gliomas. Those of the retina have the same histologic appearance as the brain tumors (Klein). The tumors of the heart usually are rhabdomyomas (Pratt-Thomas). Those of the kidney may be angiomas, fibromas, adenomas or mixed tumors; they may become malignant (Butterworth and Wilson)

## SEBACEOUS ADENOMA

An adenoma may be defined as an organoid tumor consisting of circumscribed proliferations of incompletely differentiated glandular structures. If this definition is adhered to, sebaceous adenoma is a very rare tumor. Many tumors described in the literature as sebaceous



FIG. 172. Sebaceous adenoma. The tumor is composed of lobules which are irregular in size and shape. Two types of cells compose the lobules, generative and sebaceous cells. ( $\times 100$ )

adenoma are in reality sebaceous nevi. Examples of sebaceous adenoma have been described by Reitmann, Pautrier, Woolhandler and Becker, and Lever, among others.

Sebaceous adenoma occurs as a smooth, firm, round, elevated, often slightly pedunculated, tumor. In most reported cases, the lesion was solitary, located on the face or the scalp, and measured less than 1 cm in diameter.

**Histopathology.** On histologic examination, the tumor is sharply demarcated from the surrounding tissue and is usually surrounded by a connective tissue capsule. It is composed of lobules which are irregular in size and shape (Fig. 172). Two types of cells are present

## SENILE SEBACEOUS NEVUS (SENILE SEBACEOUS HYPLERPLASIA)

Senile sebaceous nevus occurs on the face, chiefly on the forehead, in persons past middle life, and consists of either one or a few small yellowish, translucent nodules

**Histopathology.** This lesion, like nevus sebaceus (Jadassohn) and adenoma sebaceum (Pringle), is composed of large numbers of mature or nearly mature sebaceous glands. Differentiation of senile sebaceous nevus from the other two lesions on histologic grounds is therefore, often impossible. In contrast with nevus sebaceus and adenoma sebaceum (Pringle), however, ~~apocrine~~ apocrine glands or immature hair follicles are never found.

Senile sebaceous nevus is regarded by some writers as a senile hyperplasia of sebaceous glands (Gilman, Woolhandler and Becker), while others consider it as a delayed, senile organic nevus or hamartoma (Gans). The circumscribed nature of the lesions makes the latter view more likely than the former.

## FORDYCE'S DISEASE

In this condition, groups of minute, yellowish, globoid lesions are observed on the vermillion border of the lips or on the oral mucosa.

**Histopathology.** Fordyce's disease has as pathologic substrate the presence of sebaceous glands in areas where they are normally absent. It thus represents a sebaceous nevus or hamartoma.

Histologic examination reveals groups of mature sebaceous lobules located in the upper dermis. Some lobules lie free in the dermis; others lie at the end of downward proliferations of the overlying epidermis, and still others are connected with the epidermis by true sebaceous ducts (Chambers). However, ~~hairs~~ hairs are never found.

## APOCRINE GLAND NEVUS

Organic nevi composed only of apocrine glands do not occur. Apocrine gland structures, however, frequently are present in nevus sebaceus (Jadassohn) (see page 343) and occasionally in nevus verrucosus (see page 322).

## HAIR NEVUS

~~Purely~~ Purely hair nevi occur, but ~~more~~ more commonly they appear in conjunction with other nevroid lesions, for instance with nevus sebaceus (Jadassohn), nevus verrucosus and nevus pigmentosus. Some cases of adenoma sebaceum (Pringle) are hair nevi rather than sebaceous nevi (see page 345).

sebaceous lobules. In most lobules, however, the two types of cells occur in approximately equal proportions, often arranged in such a way that groups of sebaceous cells are surrounded by generative cells. Larger lobules may contain in their center cystic spaces formed by the decomposition of mature sebaceous cells.

✓ Fat stains reveal the presence of fat in the sebaceous and transitional cells and in the cystic spaces.

### SYRINGOCYSTADENOMA PAPILLIFERUM (NEVUS SYRINGOCYSTADENOMATOSUS PAPILLIFERUS)

This tumor represents an apocrine gland adenoma with differentiation predominantly toward apocrine ducts, in contrast with Hidradenoma papilliferum in which differentiation is directed predominantly toward apocrine glands.



FIG 174 Syringocystadenoma papilliferum. High magnification of Figure 173. The villi are lined by two rows of cells. The outer row is composed of small cuboidal cells (myo-epithelial cells); the inner row of high cylindrical cells (secretory cells of apocrine glands) (X200).

Clinically, syringocystadenoma papilliferum occurs not as a single verrucous papule, but as a cluster of papules. The cysts can

in the lobules. The cells of the first type are identical with the cells present at the periphery of normal sebaceous glands and resemble the cells of which basal cell epitheliomas are composed. They have



FIG 173 Syringocystadenoma papilliferum. Low magnification. A cystic invagination extends downward from the epidermis. Numerous villuslike projections extend into the lumen of the cystic invagination. A group of apocrine glands (A.P.) is present in the left lower corner. (X50)

been called generative cells by Grynfeldt (see page 17). The cells of the second type are mature sebaceous cells. They have developed from the generative cells. In addition there are cells in a transitional stage of differentiation. The distribution of the generative and the sebaceous cells within the lobules varies. Some lobules contain predominantly generative cells and thus resemble basal cell epithelioma. Other lobules contain mainly sebaceous cells and resemble mature

ence of active secretion. These cells represent secretory cells. Cellular debris is found in the lumina. Beneath the cystic invaginations deep in the dermis one finds groups of glandular lumens (Fig 175). Their secretory activity (decapitation secretion) clearly labels them as apocrine glands (Fig 176) (Tappeiner Appel Grund). Connections of the apocrine glands



FIG 176 Syringocystadenoma papilliferum. High magnification of the apocrine glands in Figure 175. The secretory cells of the apocrine glands show evidence of active secretion (decapitation secretion) ( $\times 400$ ).

with the cystic invaginations in the upper dermis can be traced without difficulty.

In most cases a dense inflammatory infiltrate is present in the upper dermis and especially in the villi. This infiltrate contains a large percentage of plasma cells and in some cases is composed almost exclusively of plasma cells.

The association of syringocystadenoma papilliferum with nevus sebaceus (Dorffel, Marcus and Wooldridge Grund) or with basal cell epithelioma (Dorffel, Reuterwall) has been noted on several occasions.

**Differential Diagnosis** For differentiation from Darier's disease in which villi are also sometimes very prominent, see page 50.

**Histopathology.** The epidermis shows acanthosis and papilloma-  
tosis. Cystic invaginations extend downward from the epidermis (Fig  
173). These invaginations represent greatly dilated apocrine ducts



FIG. 173 *Syringocystadenoma papilliferum*. Low magnification. In the upper dermis there are several cystic invaginations filled with villuslike projections. A marked inflammatory infiltrate containing many plasma cells is present around the cystic invaginations. The lower dermis contains numerous apocrine glands ( $\times 50$ ).

Numerous villuslike projections extend into the lumina of the invaginations. The cystic invaginations, as well as the villi, are lined by two rows of cells (Fig 174). The outer row is composed of small cuboidal cells with deeply staining nuclei. These cells are immature myoepithelial cells. The inner row is composed of high, cylindrical cells which have large, oval, pale-staining nuclei and may show evi

larly



Gates, Warren and Warvi have pointed out that hidradenoma papilliferum closely resembles the papillary cystadenomas of the mammary gland. It should be remembered that, phylogenetically, the mammary gland is an apocrine gland.

### SEBACEOUS EPITHELIOMA

Clinically, the tumor has no characteristic appearance. Usually, it presents itself as a solitary small nodule or plaque. It is an uncommon type of tumor. Examples have been described by Grynfeltt, Biberstein, Milian, Petin and Brunel, and Lever.

**Histopathology.** In degree of differentiation, sebaceous epithelioma stands between sebaceous adenoma, in which there are typical sebaceous lobules, and cystic basal-cell epithelioma, in which there



## HIDRADENOMA PAPILLIFERUM

This tumor occurs almost exclusively on the labia majora and on the perineum of women as a solitary intracutaneous lesion covered by normal epidermis. It usually measures only a few millimeters in diameter. Malignant degeneration does not occur.



FIG 177 Hidradenoma papilliferum. Low magnification. The tumor consists of a large cystlike lumen into which numerous interlacing villi project ( $\times 50$ ).

**Histopathology** The tumor represents an adenoma of apocrine glands (Anderson, Winer). It is located in the dermis, shows no connection with the epidermis, and is well encapsulated. It is composed of a large cystlike lumen into which numerous interlacing villi project (Fig. 177). The wall of the cyst as well as the villi are lined usually by a single layer of high cylindrical cells. These cells have a

differentiation is directed mainly toward apocrine duct cells, differentiation in syndromia is directed mainly toward apocrine gland cells and in myo epithelioma mainly toward apocrine myo epithelial cells

- 1 - in women and develops as yellowish nodules, the size



FIG 180 Syringoma. The dermis contains several small cystic ducts. The walls of most ducts are lined by two rows of epithelial cells. Colloidal material fills the lumina. Two of the ducts have comma like tails giving them the appearance of tadpoles (X200)

of a pinhead are found around the eyelids on the chest, on the abdomen and on the anterior aspects of the thighs, but occasionally also elsewhere on the skin.

**Histopathology** The dermis contains numerous, small, cystic ducts (Fig 180). The walls of the ducts are lined usually by two rows of epithelial cells. In most instances these cells are flat and compressed, evidence of ac-

is but little differentiation toward sebaceous cells (See Table 6 page 319) As Loos has put it sebaceous epithelioma grows like a basal cell epithelioma but its cells have undergone considerable differentiation toward sebaceous cells

The tumor is composed of irregularly shaped cell masses which are scattered through the upper dermis (Fig 179) The majority of cells



Fig 179 Sebaceous epithelioma The tumor is composed of irregularly shaped cell masses The majority of cells are of the same type as in basal cell epithelioma but many cells (SC) show differentiation toward sebaceous cells ( $\times 100$ )

are of the same type as the cells in basal cell epithelioma However a fairly large number are transitional cells showing beginning fatty transformation of their cytoplasm Groups of mature sebaceous cells lie in the centers of the cell masses Cysts formed by the disintegration of sebaceous cells may or may not be present in the center of some of the tumor masses

### SYRINGOMA

Three types of apocrine epitheliomas occur syringoma cylindromatous and myoeplithelioma (see Table 6) While in syringoma

Besancon Riehl Homma and Fischer Wendlberger) (2) its simultaneous occurrence with rudimentary hair structures or with trichioepithelioma a tumor of hair structures (Fischer Lever) (3) its appearance at puberty when apocrine glands first begin to function (Kyle Wendlberger) and (4) the prevalence of the lesions in areas of the skin where apocrine glands occur or used to occur phylogenetically (Wendlberger)

### CYLINDROMA

This disorder which is often hereditary is characterized by the presence of numerous rounded smooth tumors of various sizes on



Fig. 81. Tumors

apocrine gland cells ( $\times 900$ )

the scalp. Occasionally a few tumors are present also on the face and the upper trunk. The lesions begin to appear in early adulthood and increase in number and size throughout life. They vary in size from a few millimeters to several centimeters and by their arrangement in groups resemble bunches of grapes or tomatoes. The tumor

man and Besancon; Riehl; Homma and Escher; Wendtberger). The lumina of the ducts are filled with a colloidal material. Some of the cystic ducts possess small, comma-like tails of epithelial cells, giving them the appearance of tadpoles. In addition, there are solid strands of epithelial cells independent of the ducts. The cells composing the




FIG. 181 Cylindroma. Low magnification. The tumor is composed of various sized islands of epithelial cells. The islands are surrounded by a hyaline membrane. Several of the islands in the lower quadrant contain droplets of hyalin ( $\times 75$ ).

solid strands have the same appearance as the cells in basal cell epithelioma. Rudimentary hair structures are found occasionally in the lesions (Fischer).

Several authors have reported the simultaneous occurrence of syringoma and trichoepithelioma (epithelioma adenoides cysticum) in the same patient (McDonagh; Weidman and Besancon) and even in the same lesion (Ingels; Lever).

The reasons given by various authors for the apocrine genesis of syringoma are: (1) the presence of active secretion (Weidman and

Besancon Riehl Homma and Escher Wendlberger) (2) its simultaneous occurrence with rudimentary hair structures or with trichoepithelioma a tumor of hair structures (Fischer Lever) (3) its appearance at puberty when apocrine glands first begin to function (Kyrle Wendlberger) and (4) the prevalence of the lesions in areas of the skin where apocrine glands occur or used to occur phylogenetically (Wendlberger)

### CYLINDROMA

This disorder which is often hereditary is characterized by the presence of numerous rounded smooth tumors of various sizes on




FIG. 189 Cylindroma High magnification of Fig. 181 T

... of the inner row show active secretion like apocrine gland cells ( $\times 900$ )

the scalp. Occasionally a few tumors are present also on the face and the upper trunk. The lesions begin to appear in early adulthood and increase in number and size throughout life. They vary in size from a few millimeters to several centimeters and by their arrangement in groups resemble bunches of grapes or tomatoes. The tumors

man and Besincon, Riehl, Homma and Escher, Wendlberger) The lumina of the ducts are filled with a colloidal material. Some of the cystic ducts possess small, comma like tails of epithelial cells, giving them the appearance of tadpoles. In addition, there are solid strands of epithelial cells independent of the ducts. The cells composing the




FIG 181 Cylindroma low magnification. The tumor is composed of various sized islands of epithelial cells. The islands in the lower quadrant contain droplets of hyaline. (×75)

solid strands have the same appearance as the cells in basal cell epithelioma. Rudimentary hair structures are found occasionally in the lesions (Fischer).


Several authors have reported the simultaneous occurrence of syringoma and trichioepithelioma (*epithelioma adenoides cysticum*) in the same patient (McDonagh, Weidman and Besincon) and even in the same lesion (Ingels, Lever).

The reasons given by various authors for the apocrine genesis of syringoma are (1) the presence of active secretion (Weidman and

and (3) its simultaneous occurrence with tricho epithelioma a tumor of hair structures

## MYO EPITHELIOMA (MYO EPITHELIAL SWEAT GLAND TUMOR)

Myo epithelioma occurs as a rule as a solitary tumor occasionally however several lesions are present. The tumors present themselves



as firm intracutaneous nodules and usually measure between 0.5 and 2 cm in diameter although they may be larger. The overlying skin is either normal or shows slight erythema. Myo-epithelioma is a rather uncommon tumor. Examples have been described by Sheldon, Balogh, Hartz and Lever.

**Histopathology.** Histologic examination reveals a sharply circumscribed usually encapsulated essentially solid tumor. The epithelial cells of the tumor are arranged usually in intertwining bands but occasionally in lobular masses. The same two types of cells are present as in cylindroma namely secretory and myo-epithelial cells but



may cover the entire scalp like a turban. For this reason, they are referred to occasionally as turban tumors.

**Histopathology.** The tumors are composed of numerous, variously sized islands of epithelial cells. These islands are surrounded by a hyaline membrane. In addition, droplets of hyalin are present in some of the islands (Fig. 181). Two types of cells compose the islands (Stillians, Savatard, Zakon, Lever). Cells with large, oval, light staining nuclei lie in the centers of the islands, while cells with small, round, dark staining nuclei are present in palisade arrangement at the periphery of the islands and also scattered between the cells of the first type (Fig. 182). The cells of the first type may be regarded as cells differentiating toward secretory cells, those of the second type as cells differentiating toward myoepithelial cells of apocrine glands. The secretory cells always outnumber the myoepithelial cells. The hyalin which is present around and within the islands appears to be produced by the myoepithelial cells, since it is always found in apposition to them. It is particularly well demonstrated by staining with methylene blue.

In some cases, a thorough inspection of the histologic sections will reveal the presence of a few glandular lumina lined by two layers of cells: an inner actively secreting layer and an outer myoepithelial layer (Stillians, Lever) (Fig. 182). These glandular structures strongly resemble apocrine glands.

Several authors have observed connections of the tumor masses with hair follicles (Watanabe, Stillians, Schlammadinger). The simultaneous presence of cylindroma and trichoepithelioma (epithelioma adenoides cysticum) in the same patient has been observed repeatedly (Watanabe, Schlammadinger, Schuermann and Weber, Savatard). Cylindromas neverly always remain benign. Two cases, however, are on record in which malignant degeneration of the tumors occurred with metastases to the lymph nodes in one case (Lausecker) and to internal organs as well in the other (Luger). The areas of malignant degeneration showed in the cell lobules polymorphism of the nuclei, numerous mitotic figures, loss of the hyaline sheath, loss of palisading at the periphery and invasion into the surrounding stroma.

The histogenesis of cylindroma is not yet clearly established. Some authors (Coenen, Davies) regard it as a sweat gland tumor, others (Watanabe, Stillians, Schlammadinger) as a hair follicle tumor and still others (Balog, Schuermann and Weber) as an apocrine tumor. In favor of the apocrine genesis of cylindroma are (1) the presence of two types of cells, namely, secretory and myoepithelial cells, (2) the occasional presence of actively secreting glandular lumina.

and (3) its simultaneous occurrence with tricho-epithelioma, a tumor of hair structures

### MYO EPITHELIOMA (MYO EPITHELIAL SWEAT GLAND TUMOR)

Myo epithelioma occurs as a rule as a solitary tumor occasionally, however several lesions are present. The tumors present themselves




FIG 189 Myo-epithelioma. The epithelial cells are arranged in interlacing bands. Two types of cells can be seen. Secretory cells lie in the center of the bands and tend to be arranged around lumina. Myo-epithelial cells lie at the periphery of the bands. ( $\times 400$ )

14 Form

**Histopathology** Histologic examination reveals a sharply circumscribed usually encapsulated essentially solid tumor. The epithelial cells of the tumor are arranged usually in interlacing bands but occasionally in lobular masses. The same two types of cells are present as in cylindroma, namely, secretory and myo epithelial cells.

in contrast with cylindroma, the majority of cells are <sup>↑</sup>myo epithelial cells. The secretory cells lie in groups and frequently are arranged around lumina which are usually small.

In myo epitheliomas with bandlike arrangement of the epithelial cells, the secretory cells are located in the center and the myo epi




FIG 184 Myo-epithelioma. The epithelial cells are arranged in lobules. The lobule shown is composed largely of myo epithelial cells but lumina lined with secretory cells are scattered through the lobule ( $\times 200$ ).

thelial cells at the periphery of either tubules or solid bands (Fig 183). The myo epithelial cells may be seen proliferating irregularly into the stroma. In myo epitheliomas with lobular arrangement, the lobules are composed largely of myo epithelial cells, but throughout the lobules one finds lumina lined with secretory cells (Fig 181).

Some of the tumors possess a considerable amount of stroma of hyaline appearance and thus resemble the so called mixed tumors of the salivary gland type.

#### MIXED TUMOR OF THE SKIN

Mixed tumors of the salivary gland type occasionally have been described as occurring in the skin (Hirsch, Highman, Morehead,

Lennox Pearse and Richards) Their clinical appearance is the same as that of myo epithelioma

Histopathology Mixed tumors of the skin show epithelial cells arranged in nests and strands as well as around small lumina (Fig 185) Either one or two layers of epithelial cells are present around the lumina In addition epithelial cells may be seen proliferating irregularly from the lumina into the stroma The majority of epi




FIG 185 Mixed tumor of the skin Epithelial cells lie arranged around small tubular lumina and scattered through the stroma The stroma shows mucoid and hyaline degeneration ( $\times 100$ )

thelial cells are spindle shaped and resemble myo-epithelial cells (Morehead) Those located around lumina have the appearance of glandular cells The stroma usually is abundant and shows hyaline and mucoid degeneration

Some authors regard the mixed tumors of the skin as of sweat gland origin (Hirsch Lennox Pearse and Richards) It is the author's belief that most if not all mixed tumors of the skin represent myo epitheliomas They differ from myo epithelioma only by showing a greater amount of stroma and more marked degenerative changes in the stroma.

#### CLEAR CELL MYO EPITHELIOMA

The clinical appearance of this tumor is that of a solitary nodule usually covered by intact skin but occasionally discharging serous material




FIG 186 Clear cell myoepithelioma The tumor is composed of two types of cells, uniform myoepithelial cells and suboidal clear cells. (X200)



FIG 187 Clear-cell myoepithelioma There are numerous clear cells and in addition, tubular lumina lined by a single layer of secretory cells (X200)

**Histopathology** The tumor has a lobular structure. Two types of epithelial cells can be recognized. Uniform myo-epithelial cells with deeply basophilic nuclei and clear cells which are cuboidal have a distinct cellular membrane, very clear cytoplasm and a round nucleus (Fig 186). The clear cells seem to develop from the myo-epithelial cells (Lever and Castleman). Varying amounts of glycogen are present in the clear cells. In some tumors there are groups of keratinized cells, cysts and glandular structures. The glandular structures are tubular and their lining cells often show decapitation secretion as observed in apocrine glands (Fig 187).

While Lever and Castleman regard this type of tumor as a variant of myo-epithelioma, Liu because of the presence of glycogen thought that it was derived from the outer hair sheath which normally contains abundant amounts of glycogen. However glycogen is found so commonly in young cells and in tumors that its presence cannot be considered as a reliable criterion for the histogenetic classification of neoplasms.

### TRICHO EPITHELIOMA (EPITHELIOMA ADENOIDES CYSTICUM, MULTIPLE BENIGN CYSTIC EPITHELIOMA)

The name trichoepithelioma is preferable to the other names listed above because it indicates that differentiation in this tumor is directed toward hair structures.

The disorder begins as a rule at the age of puberty and is frequently hereditary. It is characterized by the presence of numerous pinhead to pea-sized rounded yellowish or pink nodules on the face and occasionally on the upper trunk. A few telangiectatic vessels are often present on the surface of the larger lesions. Occasionally one or several lesions become ulcerated because of change into a basal-cell epithelioma.

**Histopathology** On histologic examination trichoepithelioma appears as a well-circumscribed tumor. Horn cysts represent the characteristic lesion. They consist of a fully keratinized center surrounded by a shell of flattened basal cells without prickles (Fig 188). The keratinization in the horn cysts is abrupt and complete not radial and incomplete as in the horn pearls of squamous-cell carcinoma. This process corresponds to the abrupt development of the horn cells of the hair from the hair matrix cells (which are also cells without prickles). It may be concluded therefore that the cells surrounding the horn cysts are hair matrix cells and that the horn cysts represent attempts at hair shaft formation (Lever). Occasionally one sees prickles around some of the horn cysts. They represent outer hair sheath cells. Since the outer hair sheath develops from

prickle cells at a time when the hair germ has already advanced to a rather high stage of differentiation the presence of prickle cells around the horn cysts is evidence of rather high differentiation

In addition to horn cysts irregularly shaped islands and intertwining strands composed of basal cells are present. Such areas are indistinguishable from basal-cell epithelioma (Goldman). Abortive



FIG. 188 Trichoepithelioma. The tumor contains numerous horn cysts. In the center near the epidermis two rudimentary hairs can be seen ( $\times 100$ ).

hair papillae and hair shafts are seen occasionally. Since hair papillae contain a high concentration of alkaline phosphatase (Hardy) their presence can be well demonstrated by the use of the Gomori stain for alkaline phosphatase. Calcification of the horn cysts may occur and evoke a foreign body giant cell reaction in the adjacent connective tissue.

Tumors histologically in an intermediate stage between trichoepithelioma and basal cell epithelioma occur (Summerill and Hurton Traenkle). As just stated a typical trichoepithelioma is characterized by circumscribed growth and by the presence of many horn cysts and of basal cell masses; however basal cell epithelioma may

also show horn cysts (Fig. 189) (see page 379) and may be circumscribed in growth. Since thus no sharp line of demarcation exists between tricho-epithelioma and basal-cell epithelioma histologically, it may be necessary, in order to arrive at a diagnosis in a given case, to have knowledge of clinical data such as the number and the distribution of the lesions and the age at which the lesions first ap-




FIG. 189. Basal-cell epithelioma with horn cysts. Histologically, this tumor is in an intermediate stage of differentiation between basal-cell epithelioma and tricho-epithelioma. Clinically the lesion was a basal-cell epithelioma ( $\times 200$ ).

peared. The close relation between the two types of tumors is attested further by reports of cases in which one or several of the lesions of tricho-epithelioma after having persisted as such for many years, developed into ulcers with the histologic picture of basal-cell epithelioma (Adamson, Little, Savatard). This close relationship of the two types of tumors can be explained best by assuming that they have a common genesis from primary epithelial germs (Lever) and that they differ only in the degree of maturity of their cells (Adamson). Since cells of various maturity may occur in the same lesion, one may find in tricho-epithelioma areas consistent with the histologic picture of basal-cell epithelioma and *vice versa*, also, if active growth occurs in a lesion of tricho-epithelioma, the newly formed cells may



be more embryonal than the older cells and the lesion may grow as a basal cell epithelioma.

A close relation exists not only between tricho epithelioma and basal cell epithelioma, but also between tricho epithelioma and other types of benign epitheliomas such as syringoma and cylindroma. Tricho epithelioma may occur with syringoma or cylindroma in the same patient (see pages 356, 358).

### CALCIFYING EPITHELIOMA (MALHERBL)

Calcifying epithelioma is a solitary hard, deep seated tumor which is covered by normal skin. It occurs most frequently on the face and



FIG 190 Calcifying epithelioma (Malherbe) The tumor consists of lobules embedded in a connective tissue stroma. Two types of cells compose the lobules: basophilic cells and shadow cells. The basophilic cells resemble the cells of basal cell epithelioma. The shadow cells show a central unstained shadow at the site of the nucleus. In the center of the field one can see transformation of the basophilic cells into shadow cells. The stroma contains numerous foreign body giant cells ( $\times 100$ ).

the upper extremities. The size is from 0.5 to 3 or even 5 cm in diameter.

**Histopathology** Calcifying epithelioma is a sharply demarcated, often encapsulated tumor and usually located in the lower dermis.

or in the subcutaneous fat. The histologic appearance is characteristic. Embedded in a connective tissue stroma variously shaped masses of epithelial cells are present. As a rule two types of cells—basophilic cells and shadow cells—compose these masses (Fig 190). Occasionally basophilic cells are absent. The basophilic cells greatly resemble the cells of basal cell epithelioma. They possess



FIG 191. Calcifying epithelioma (Malherbe). Small and large areas of calcification are present within the lobules of shadow cells (X100).

large, round or elongated, deeply basophilic nuclei and only little cytoplasm, so that the nuclei are densely packed. The cellular borders are often indistinct so that it appears as if the nuclei were embedded in a symplasmic mass. The shadow cells stain pale pink with hematoxylin-eosin. They have a distinct border but take no nuclear staining. Instead they show a central unstained shadow at the site of the nucleus. In some areas one can see clearly that the shadow cells develop from the basophilic cells (Fig 190). In tumors of recent origin numerous areas of basophilic cells are usually present. As the lesion ages the number of basophilic cells decreases due to their development into shadow cells, and in tumors of long standing few or no basophilic cells remain. In addition to basophilic and shadow

cells, areas of cornification are present. The cornification may occur as sheets of horn cells or as small, round centers of cornification. Sheets of horn cells are found only within the masses of shadow cells while the small, round centers of cornification may lie within areas of basophilic cells as well as within the masses of shadow cells. In occasional instances, melanin is present within the tumor. The melanin may be located in either the basophilic cells, the shadow cells or the stroma (Turhan and Krammer, Lever and Griesemer).

The stroma of the tumor usually shows a considerable foreign body giant cell reaction adjacent to the masses of shadow cells. A frequent but not constant feature is calcification (Fig. 191). The calcium may be present in large sheets replacing the shadow cells or may be present as fine granules within the cytoplasm of shadow cells. Occasionally, ossification of calcified areas takes place (Nicholson, Highman and Ogden). Malignant degeneration does not occur.

**Histogenesis.** The histogenesis of calcifying epithelioma is not established fully. Originally described by Malherbe and Chenantais as a calcified epithelioma of sebaceous glands, it has been variously regarded as a tumor of misplaced rests of sebaceous gland epithelium (Sutton and Sutton), as a basal cell epithelioma with degeneration instead of differentiation of cells (Fink, Muehlton) or as a growth intermediary between epidermal cyst and basal-cell epithelioma (Côte). Turhan and Krammer expressed the opinion that calcifying epithelioma arises from hair matrix cells. They believe that the basophilic cells are hair matrix cells and that the formation of shadow cells represents a form of keratinization, an attempt at hair shaft formation. In support of their theory, these authors point out that the basophilic cells resemble hair matrix cells, that the small, round centers of keratinization observed in the epithelial formations of these tumors resemble cross sections of hair shafts and that the melanin in some of these tumors is analogous to the melanin found in normal hair. Highman and Ogden have arrived at similar conclusions.

Lever and Griesemer regard the basophilic cells in calcifying epithelioma as primary epithelial germ cells with a tendency to differentiation into keratotic hair cells. The basophilic cells, thus, are assumed to have a function akin to that of hair matrix cells, but, being less mature than hair matrix cells, they produce not hair but irregular masses of immature hair cells (shadow cells). Accordingly, in a classification of primary epithelial germ tumors, calcifying epithelioma of Malherbe can be placed under those with differentiation toward hair structures (see Table 6, page 319).

**Differential Diagnosis.** Epidermal and sebaceous cysts may resemble calcifying epithelioma of Malherbe if their contents has un-

undergone partial calcification and if they have ruptured and due to the resulting foreign body giant-cell reaction, have undergone partial disintegration. They then share with calcifying epithelioma the presence of irregular islands of epithelial cells of areas of calcification and of a foreign body giant cell reaction. However, basophilic cells and shadow cells are not found. It should be remembered that old lesions of calcifying epithelioma may show no more basophilic cells so that then the presence of shadow cells alone is the decisive factor in favor of calcifying epithelioma.

Differentiation of calcifying epithelioma from sebaceous cyst with secondary basal cell epithelioma also rests largely on the presence of shadow cells since the areas of basophilic cells in calcifying epithelioma resemble basal-cell epithelioma, however, in calcifying epithelioma the epithelial structures never show keratinizing of the peripheral-cell layer and often there are areas in which one can observe the transformation of basophilic into shadow cells.

### BASAL-CELL EPITHELIOMA

Basal-cell epithelioma may occur anywhere on the skin except on the palms and the soles. However, the face—particularly the periorbital region—is by far the most common site of location. The mucous membranes are never affected. Although basal-cell epithelioma occurs usually as a single lesion, multiple lesions are not infrequent.

present in

types of

epithelioma including rodent ulcer, (2) pigmented basal-cell epithelioma, (3) morphea like or fibrosing basal-cell epithelioma and (4)

first as

telangi

also in size and number

secondary ulceration

ing ulcer sur

rodent ulcer

Pigmented basal-cell epithelioma differs from the nodulo ulcerative type only by the dark pigmentation of the lesion.

Morphea like or fibrosing basal-cell epithelioma manifests itself as an only slightly elevated firm yellowish plaque with ill-defined borders over which the skin remains intact for a long time before finally ulceration occurs.

threadlike, pearly border The patches usually show small areas of superficial ulceration and crusting, in addition, their center may show smooth, atrophic scarring

Basal cell epitheliomas cannot be produced experimentally like squamous cell carcinomas However, they can develop following prolonged administration of inorganic arsenic (Anderson) (see page 155) and, in rare instances, in areas of radiodermatitis (Anderson and Anderson) (see page 122) Furthermore, basal cell epitheliomas of the face occur much more frequently in areas with much sunshine, as in Australia and in the south of the United States, than in less sunny regions, such as England and the north of the United States

Basal cell epitheliomas do not metastasize, as a rule It seems, however, that exceptions to this rule occur (Montgomery, 1928, Foot Amersbach, Lattes and Kessler) Lattes and Kessler found a total of 20 cases of metastasizing basal cell epitheliomas reported in the literature Yet many authors doubt the occurrence of metastases in basal cell epithelioma Walther expressed the belief that cases reported as metastasizing basal cell epithelioma represent wrong diagnoses

**Histogenesis.** Several theories exist regarding the origin of basal cell epithelioma Krompecher, the original describer of basal cell epithelioma stated in 1903 that he regarded these tumors as carcinomas of the basal cells of the epidermis He believed that those tumors which showed a tendency to gland formation imitated the embryonal gland formation of the basal cells Krompecher's view is still adhered to by many (Montgomery 1940, Teloh and Wheelock) However, Kramz believed that only those basal cells which develop into glandular cells give rise to basal cell carcinoma Geschickter and Koehler shared this view They suggested the designation appendage cell carcinoma Mallory and Haythorn held the opinion that basal cell epitheliomas were carcinomas of hair matrix cells

Foot expressed the view that basal cell epitheliomas were carcinomas which develop from distorted primordia of dermal adnexa rather than from ordinary epidermal basal cells He stated that the tumors took origin from any or all three types of adnexal primordia i.e., hair, sebaceous gland and sweat gland and imitated their embryonal development

The first author to express doubts that basal cell epitheliomas were carcinomas was Adamson who in 1911 stated that, in his opinion basal cell epitheliomas were nevroid tumors originating from latent embryonic foci aroused from their dormant state at a later period of life He believed that the latent embryonic foci usually were embryonic pilosebaceous follicles but occasionally were embryonic sweat ducts Several authors have since reached sim

cently Wallace and Halpert have suggested that basal-cell epitheliomas are benign tumors either of the hair matrix or of the hair invagination and have proposed the term trichoma for them.

It is the author's belief that basal-cell epitheliomas are not carcinomas and are not derived from basal cells but instead are nevusoid

entiated anaplastic cells. In accordance with the potentiality inherent in the primary epithelial germ to differentiate into sebaceous glands, apocrine glands and hair, differentiation in basal cell epitheliomas can be either toward sebaceous gland, apocrine gland or hair structures. (See histogenesis and classification of epidermal tumors, page 317 and Table 6, page 319).

Pinkus recently has suggested that basal cell epitheliomas arise not from an embryonal rest such as the primary epithelial germ but from pluripotential cells which have formed during adult life and can like the primary epithelial germ differentiate in the direction of sebaceous glands, apocrine glands and hair. The fact that basal cell epithelioma occasionally arises in areas of radiodermatitis and following the ingestion of arsenic is cited by Pinkus as evidence (although it could as well be assumed that these agents merely stimulate dormant primary epithelial germs to active growth).

Meuck has pointed out that in basal cell epitheliomas the connective tissue stroma always shows a close relationship to the tumor and proliferates with it just as it does in benign fibroepithelial tumors and in adenomas. On the other hand, in carcinomas the connective tissue usually is stretched then tears until it finally disintegrates. He therefore regards basal-cell epithelioma not as a carcinoma but as a solid adenoma.

**Histopathology.** The characteristic cell of basal-cell epithelioma has a large oval or elongated deeply basophilic nucleus and little cytoplasm. The cytoplasm is often defined poorly so that it may appear as if the nuclei were embedded in a sympylasmic mass. The nuclei resemble those of the basal cells of the epidermis very closely but the cells of basal cell epithelioma have no connection with the epidermis. They are by not

true carcinoma. A carcinoma is an invasive tumor in most instances. Scattered islands of tumor cells are often found away from the main tumor deep in the dermis and even in the subcutaneous tissue.

latter showing differentiation toward primary epithelial germ structures—i.e. toward sebaceous glands, apocrine glands or hair. A sharp dividing line between the two groups cannot be drawn because many undifferentiated basal cell epitheliomas show differentiation in some areas and most differentiated basal cell epitheliomas show areas lacking differentiation. Correlating the clinical with the histologic classification it can be stated that the nodulo ulcerative type of basal-cell epithelioma may be differentiated or undifferentiated whereas pigmented basal-cell epithelioma, fibrosing basal cell epithelioma and superficial basal cell epithelioma usually are undifferentiated. If the nodulo ulcerative basal cell epithelioma shows no differentiation it is called solid basal cell epithelioma if it shows differentiation it may be either a cystic or adenoid or a keratotic basal cell epithelioma.

**I UNDIFFERENTIATED BASAL CELL EPITHELIOMAS** The group of undifferentiated basal cell epitheliomas includes the solid basal cell epithelioma, the pigmented basal cell epithelioma, the morphia-like or fibrosing basal cell epithelioma and the superficial basal cell epithelioma.

**A SOLID BASAL CELL EPITHELIOMA** (primordial type of basal cell epithelioma Foot) In this form variously sized and shaped masses of tumor cells are embedded in the dermis (Fig. 192). Some of the masses originate from the surface epidermis and have no connections with it even on serial sections. Others see a mass of tumor cells originate from a hair follicle. In the masses of tumor cells the nuclei of cells often show palisade arrangement whereas the nuclei of the cells inside lie in haphazard fashion.

A mild or moderately severe inflammatory reaction may be present in the dermis particularly in the more rapidly growing tumors but it may be entirely lacking. The connective tissue frequently proliferates with the tumor and is arranged in parallel bundles around the tumor masses so that a definite mutual relationship seems to exist between the parenchyma of the tumor and its stroma (Hueck Pinkus). The connective tissue nearest to the tumor masses often undergoes mucinous degeneration. Since mucin shrinks during fixation the stroma frequently retracts from the tumor islands so that in prepared sections the tumor islands seem to lie free in cavities (Fig. 201). Although this retraction represents merely an artefact caused by shrinkage during fixation it is quite typical of basal cell epithelioma and aids in its recognition. In squamous cell carcinoma the stroma often is atrophic. (In basal cell epithelioma and in squamous cell carcinoma the stroma is often atrophic.)

✓ Some basal-cell epitheliomas with little or no structural differentiation toward the cutaneous appendages show evidence nevertheless of cellular differentiation by presenting two types of cells. One type of cell has small, elongated, deeply basophilic nuclei, the other has




Fig. 10. Some basal-cell epithelioma (undifferentiated or primordial basal-cell epithelioma).

large round cells

to the secretory cells of either sebaceous or apocrine glands

**B. PIGMENTED BASAL-CELL EPITHELIOMA** Although the presence of melanin can be demonstrated by silver-stains in many basal-cell epitheliomas (Becker) large amounts are encountered only rarely. The



presence of melanin in basal cell epitheliomas can be explained by the fact that melanin producing melanocytes are present not only in the surface epidermis but also in the primary epithelial germ (As the primary epithelial germ matures into the hair these melanocytes arrange themselves among the hair matrix cells overlying the hair papilla and form the melanin of the hair (See Fig 9)




FIG. 133 Basal cell epithelioma with differentiation into two types of cells. One type of cell has large oval pale nuclei, the other type has small elongated dark nuclei ( $\times 400$ )

Basal cell epitheliomas with large amounts of pigment contain interspersed between the tumor cells numerous melanin laden melanocytes (Bloch). The connective tissue of these tumors contains numerous melanophores (Eller and Anderson, Becker, Foot).

C. MORPHEA LIKE OR FIBROSING BASAL CELL EPITHELIOMA. In this variant connective tissue proliferation is much greater than in the other types of basal cell epithelioma. Imbedded in a dense fibrous stroma one observes innumerable groups of closely packed tumor cells arranged in elongated strands (Fig 194) (Caro and Howell). Most of the strands are small but others are larger and show branching. The groups of tumor cells often extend deeply into the dermis.

D. SUPERFICIAL BASAL CELL EPITHELIOMA. This type of basal cell epithelioma originates from the epidermis in multiple foci (Fig 195).

The peripheral-cell layer of the budlike proliferations usually shows palisade arrangement. There is little penetration into the dermis. It may be pointed out that the epidermal buds often show great resemblance to the primary epithelial germ buds as seen in the em



FIG. 191 Morpheiform or fibrosing basal cell epithelioma. Innumerable small groups of closely packed tumor cells, many of them arranged in elongated strands, are embedded in a dense fibrous stroma ( $\times 100$ ).

brional skin (see Figure 1). The overlying epidermis usually shows atrophy, but may show acanthosis and papillomatosis. A mild to moderate amount of inflammatory infiltrate composed of lymphocytes and plasma cells is present in the upper dermis (Wise-Montgomery, 1929).

Some superficial basal-cell epitheliomas, after having persisted as such for various lengths of time, become true invasive basal-cell epi-

presence of melanin in basal cell epitheliomas can be explained by the fact that melanin producing melanocytes are present not only in the surface epidermis but also in the primary epithelial germ (As the primary epithelial germ matures into the hair these melanocytes arrange themselves among the hair matrix cells overlying the hair papilla and form the melanin of the hair (See Fig 9)




FIG 193 Basal cell epithelioma with differentiation into two types of cells. One type of cell has large oval pale nuclei the other type has small elongated dark nuclei ( $\times 400$ )

Basal cell epitheliomas with large amounts of pigment contain interspersed between the tumor cells numerous melanin laden melanocytes (Bloch). The connective tissue of these tumors contains numerous melanophores (Eller and Anderson, Becker Foot).

**C MORPHEA LIKE OR FIBROSING BASAL CELL EPITHELIOMA** In this variant connective tissue proliferation is much greater than in the other types of basal cell epithelioma. Embedded in a dense fibrous stroma one observes innumerable groups of closely packed tumor cells arranged in elongated strands (Fig 194) (Carr and Howell). Most of the strands are small but others are larger and show branching. The groups of tumor cells often extend deeply into the dermis.

**D SUPERFICIAL BASAL CELL EPITHELIOMA** This type of basal-cell epithelioma originates from the epidermis in multiple foci (Fig 195)

The peripheral-cell layer of the budlike proliferations usually shows palisade arrangement. There is little penetration into the dermis. It may be pointed out that the epidermal buds often show great resemblance to the primary epithelial germ buds as seen in the em



FIG. 123. Morphea like or fibrosing basal cell epithelioma. Innumerable small groups of closely packed tumor cells, many of them arranged in elongated strands, are embedded in a dense fibrous stroma. (X100).

horizontal skin (see Figure 1). The overlying epidermis usually shows acropia, but may show acanthosis and papillomatosis. A mild to moderate amount of inflammatory infiltrate composed of lymphocytes and plasma cells is present in the upper dermis (Wise, Montgomery 1929).

Some superficial basal-cell epitheliomas, after having persisted as such for various lengths of time, become true invasive basal cell epi

theliomas Since this change may at first be limited to a few areas, representative sections throughout the entire tissue block should be examined

A rare type of superficial basal cell epithelioma is the intra epidermal type In this type, the epidermis shows acanthosis Embedded in the broadened epidermis one finds multiple, sharply demarcated foci of basal cell epithelioma The tumor cells stand out clearly from the surrounding epidermis by their strongly basophilic staining (Montgomery, 1929, Sims and Parker)



FIG 195 Superficial basal cell epithelioma The tumor shows multiple points of origin from the epidermis Note the similarity between the tumor buds in this illustration and the primary epithelial germ buds in the embryonal skin shown in Figure 1 ( $\times 100$ )

**2 DIFFERENTIATED BASAL CELL EPITHELIOMAS** Differentiation in basal cell epitheliomas may proceed in three directions toward sebaceous glands, apocrine glands or hair (Foot, Lever) Those with differentiation toward sebaceous glands are called cystic, those with differentiation toward apocrine glands are called adenoid, and those with differentiation toward hair are called keratotic basal-cell epitheliomas In most differentiated basal cell epitheliomas, differentiation is directed toward more than one of the three structures For example, areas of keratinization may be found in a tumor which also shows glandular structures

**A CYSTIC BASAL CELL EPITHELIOMA** In this type, cystic spaces are present in the center of the tumor masses The cysts may form in three ways by degeneration of stroma that has become enclosed into the tumor, by degeneration of the center of the tumor masses, and

by differentiation of the cells in the center of the tumor masses to

the vicinity of the cyst are apt to be vacuolated (Foot) or even foamy (Pierard and Dupont) (Fig 196)




FIG 196 Cystic basal-cell epithelioma. The cyst has formed by disintegration of cells with sebaceous differentiation ( $\times 50$ )

**B. ADENOID BASAL CELL EPITHELIOMA** This type of tumor shows formations suggesting tubular glandlike structures. The cells are arranged in intertwining strands and radially around islands of connective tissue resulting in a honeycomb patterning of the tumor (Fig 197). In some tumors one may find lumina surrounded by cells which have a tendency to

**C. KERATOTIC BASAL-CELL EPITHELIOMA (pilar type Foot)** In this type cells with elongated nuclei and slightly eosinophilic cytoplasm are seen traversing the tumor as short bands or are arranged in concentric layers around parakeratotic or keratotic centers. As pointed out by Krompecher who first described this type of basal-cell epi



FIG 197 Adenoid basal cell epithelioma The strands of epithelial cells show a lacelike pattern The stroma shows mucoid degeneration ( $\times 200$ )




FIG 198 Adenoid basal-cell epithelioma The tumor contains lumina surrounded by cells that have the appearance of glandular cells ( $\times 200$ )

thelioma the elongated cells cornify abruptly so that the parakeratotic or keratotic centers are sharply demarcated against the surrounding cells (Fig 199)

otic or parakeratotic centers in keratotic basal-cell epithelioma develop from hair matrix cells and represent attempts at hair shaft formation (Foot) The keratinization in keratotic basal cell epithelioma thus is one of hair matrix cells which are cells without prickles and not one of squamous cells

Some keratotic basal-cell epitheliomas possess large centers of keratinization so-called horn cysts (Fig 200) They are like those observed in tricho-epithelioma (page 363) These horn cysts must not be confused with the horn pearls that occur in squamous cell carcinoma (see below under differential diagnosis)

**BASAL SQUAMOUS CELL EPITHELIOMA** Several authors (Darier and Ferrand Montgomery 1908 Juon) have described basal-cell epitheliomas with features of squamous-cell carcinoma According to Montgomery they represent metamorphosis of basal-cell to squamous-cell epitheliomas He stated that from 15 to 20 per cent of all basal-cell epitheliomas present such changes

Two types of basal squamous cell epithelioma are recognized by the authors named above a mixed and an intermediary type The mixed type is described as showing partial horn pearl formation with a peribornous rather than a horny center The intermediary type is described as showing two kinds of cells those with small elongated deeply basophilic nuclei regarded as basal cells and those with large round pale staining nuclei regarded as intermediary in character between basal and squamous cells

The existence of basal squamous-cell epithelioma is not generally accepted (Lever Welton Elliott and Himmelstiel Lennox and Wells) It is likely that the mixed type of basal squamous cell epithelioma represents a keratotic basal-cell epithelioma (Fig 199) and the intermediary type a solid basal-cell epithelioma in which the cells with large round pale staining nuclei show differentiation to keratinizing cells of either sebaceous or apocrine glands (Fig 193) (see page 373)

**MIXED CARCINOMA** These tumors show





FIG 197 Adenoid basal cell epithelioma The strands of epithelial cells show a lacelike pattern The stroma shows mucoid degeneration ( $\times 200$ )




FIG 198 Adenoid basal cell epithelioma The tumor contains lumina surrounded by cells that have the appearance of glandular cells ( $\times 200$ )

However, they  
 ng much more  
 it to remember

that keratinization is not a prerogative of squamous cell carcinoma



tains unusually large horn cysts. ( $\times 50$ )

but occurs also in basal-cell epitheliomas with differentiation toward hair structures (See "Keratotic Basal Cell Epithelioma," page 377.) Keratinization in basal-cell epitheliomas may be partial ("parakeratotic centers") or complete ("horn cysts"). The keratinization differs from that seen in the horn pearls of squamous-cell carcinomas by occurring abruptly without the interposition of gradually keratinizing squamous cells. The fairly common presence in basal-cell epithelioma of areas of retraction of the tumor cell masses from the sur-

lesions (for instance, lupus vulgaris, gumma and burns), basal cell epithelioma may stimulate the development of a squamous cell carcinoma. Before making a diagnosis of "mixed carcinoma," one must rule out the possibility of pseudo epitheliomatous hyperplasia occurring in a basal-cell epithelioma (see page 334).




FIG 199 Keratotic basal cell epithelioma (pilar type of basal cell epithelioma) The cell masses in this tumor are composed largely of elongated cells lying in concentric arrangement. Several of the cell masses contain a horn cyst in their center. The elongated cells contain no prickles; they represent hair matrix cells. Note the abrupt and complete keratinization in the horn cysts ( $\times 100$ ).

**Differential Diagnosis.** Differentiation of basal cell epithelioma from squamous cell carcinoma may be difficult at times—so difficult that many authors have decided that intermediary forms ("basal squamous cell epithelioma") occur. As a rule, however, differentiation is fairly easy. One of the best points of differentiation is that the cells of basal cell epithelioma stain deeply basophilic, whereas most cells of squamous cell carcinoma at least in Grades I and II, have an eosinophilic tint due to partial or complete keratinization. In squamous cell carcinoma, Grades III and IV, the cells, because of

noid type (Hookey) The former type is more common than the latter. The latter shows considerable acanthosis. The acanthosis is due entirely to upward growth of epithelial cells. Thus, the lower base of the tumor lies above a straight line that may be drawn from the normal epidermis at one end of the tumor to the normal epidermis at the other end and the lesion appears as though tacked on the sur-



FIG. 202 Basal-cell papilloma, keratotic type. Low magnification. The superficial location of the tumor is apparent. The lesion appears as if tacked on the epidermis. (X25)

tracts of epithelial cells surrounding islands of connective tissue. The horny layer invaginates in plues deep into the lesion. Because of the tortuosity of these invaginations, cystic inclusions of horny material result (Fig. 203). They are surrounded by a shell of squamous cells in small amounts. It is located in the clear cells") are prominent in the pigmented portions.

The nodular type of basal-cell papilloma shows, originating from the epidermis, numerous thin tracts composed of a double row of cells. These cells resemble those of basal-cell epitheliomas. The tracts

rounding connective tissue (see page 372) also aids in differentiating it from squamous cell carcinoma, since such areas of retraction are not found in the latter.

The differential diagnosis between basal cell epithelioma and tricho epithelioma already has been discussed (see page 364)



FIG 201 "Mixed carcinoma" A basal cell epithelioma (left) and a squamous cell carcinoma (right) lie side by side ( $\times 50$ )

### BASAL CELL PAPILLOMA (VERRUCA SENILIS, KERATOSIS SEBORRHEICA)

Basal cell papillomas develop, often in large numbers, on the trunk, the face and the arms in persons past middle life. They are sharply circumscribed, slightly raised, verrucous, more or less pigmented lesions which often look as if "stuck on" the surface of the skin. The verrucous covering of the lesions tends to have a soft, greasy consistency. Although most lesions measure only a few millimeters in diameter, an occasional lesion may reach a size of several centimeters.

Histopathology. This tumor represents a squamous cell papilloma in which there are areas of proliferation of cells of the same type as seen in basal-cell papilloma.

Basal cell papillomas occur in two types a keratotic and an ade

noid type (Hookey) The former type is more common than the latter. The same lesion is shows considerable The acanthosis is due entirely to upward growth of epithelial cells. Thus the lower base of the tumor lies above a straight line that may be drawn from the normal epidermis at one end of the tumor to the normal epidermis at the other end and the lesion appears as though tacked on the sur-



FIG. 202. Basal-cell papilloma keratotic type. Low magnification. The superficial location of the tumor is apparent. The lesion appears as if tacked on the epidermis. (X25)

face of the skin (Fig. 202). The acanthosis is due in part to proliferation of the basal cells.

These cells resemble those of basal-cell epitheliomas. The tracts of epithelial cells surrounding islands of connective tissue. The horny layer invaginates in places deep into the lesion. Because of the tortuosity of these invaginations cystic inclusions of horny material result (Fig. 203). They are surrounded by a shell of squamous cells. Melanin is present in small amounts but is located in the basal layer.

These cells resemble those of basal-cell epitheliomas. The tracts of epithelial cells surrounding islands of connective tissue. The horny layer invaginates in places deep into the lesion. Because of the tortuosity of these invaginations cystic inclusions of horny material result (Fig. 203). They are surrounded by a shell of squamous cells. Melanin is present in small amounts but is located in the basal layer.

rounding connective tissue (see page 372) also aids in differentiating it from squamous-cell carcinoma, since such areas of retraction are not found in the latter.

The differential diagnosis between basal-cell epithelioma and tricho epithelioma already has been discussed (see page 364)




FIG 201 "Mixed carcinoma" A basal cell epithelioma (left) and a squamous cell carcinoma (right) lie side by side ( $\times 50$ )

### BASAL-CELL PAPILLOMA (VERRUCA SENILIS, KERATOSIS SEBORRHEICA)

Basal cell papillomas develop, often in large numbers, on the trunk, the face and the arms in persons past middle life. They are sharply circumscribed, slightly raised, verrucous, more or less pigmented lesions which often look as if "stuck on" the surface of the skin. The verrucous covering of the lesions tends to have a soft, greasy consistency. Although most lesions measure only a few millimeters in diameter, an occasional lesion may reach a size of several centimeters.

**Histopathology.** This tumor represents a squamous cell papilloma in which there are areas of proliferation of cells of the same type as seen in basal-cell papilloma

Basal-cell papillomas occur in two types, a keratotic and an ade-

branch and interweave (Fig 204) Considerable amounts of melanin usually are present in the tracts Cystic inclusions of horny material are absent The demarcation at the lower border is sharp just as in the keratotic type of basal-cell papilloma

Although basal-cell papilloma shows no tendency to invasion of the dermis it may be regarded as related to basal-cell epithelioma because a large proportion of the cells are of the same type as found in basal-cell epithelioma However actual change of basal cell papilloma into a frank basal cell epithelioma with deep invasion and ulceration is rare (Eller and Ryan Pinkus)

(For differential diagnosis from nevus verrucosus see page 322 and from keratosis senilis see page 328)

### DERMATOSIS PAVULOSA NIGRA

This condition is frequent among Negroes especially females The lesions are located on the face especially in the malar regions and consist of minute soft rounded usually hyperpigmented papules

Histopathology The histologic changes are the same as in basal cell papilloma Melanin pigmentation of the basal layer is pronounced (Michael and Seale)

### CARCINOMA OF SEBACEOUS GLANDS

Carcinomas of sebaceous glands have been described occasionally in the literature Some authors are more inclined to make this diagnosis than others Warren and Warvi for instance reported 28 personally observed cases while most other authors have reported only single instances (Beach and Severance)

Carcinomas of sebaceous glands occur most frequently on the eyelids originating from meibomian glands which are modified

nywhere

picture

ion consists

<sup>12</sup> <sup>1</sup> <sup>2</sup> <sup>3</sup> <sup>4</sup> <sup>5</sup> <sup>6</sup> <sup>7</sup> <sup>8</sup> <sup>9</sup> <sup>10</sup> <sup>11</sup> <sup>12</sup> <sup>13</sup> <sup>14</sup> <sup>15</sup> <sup>16</sup> <sup>17</sup> <sup>18</sup> <sup>19</sup> <sup>20</sup> <sup>21</sup> <sup>22</sup> <sup>23</sup> <sup>24</sup> <sup>25</sup> <sup>26</sup> <sup>27</sup> <sup>28</sup> <sup>29</sup> <sup>30</sup> <sup>31</sup> <sup>32</sup> <sup>33</sup> <sup>34</sup> <sup>35</sup> <sup>36</sup> <sup>37</sup> <sup>38</sup> <sup>39</sup> <sup>40</sup> <sup>41</sup> <sup>42</sup> <sup>43</sup> <sup>44</sup> <sup>45</sup> <sup>46</sup> <sup>47</sup> <sup>48</sup> <sup>49</sup> <sup>50</sup> <sup>51</sup> <sup>52</sup> <sup>53</sup> <sup>54</sup> <sup>55</sup> <sup>56</sup> <sup>57</sup> <sup>58</sup> <sup>59</sup> <sup>60</sup> <sup>61</sup> <sup>62</sup> <sup>63</sup> <sup>64</sup> <sup>65</sup> <sup>66</sup> <sup>67</sup> <sup>68</sup> <sup>69</sup> <sup>70</sup> <sup>71</sup> <sup>72</sup> <sup>73</sup> <sup>74</sup> <sup>75</sup> <sup>76</sup> <sup>77</sup> <sup>78</sup> <sup>79</sup> <sup>80</sup> <sup>81</sup> <sup>82</sup> <sup>83</sup> <sup>84</sup> <sup>85</sup> <sup>86</sup> <sup>87</sup> <sup>88</sup> <sup>89</sup> <sup>90</sup> <sup>91</sup> <sup>92</sup> <sup>93</sup> <sup>94</sup> <sup>95</sup> <sup>96</sup> <sup>97</sup> <sup>98</sup> <sup>99</sup> <sup>100</sup> <sup>101</sup> <sup>102</sup> <sup>103</sup> <sup>104</sup> <sup>105</sup> <sup>106</sup> <sup>107</sup> <sup>108</sup> <sup>109</sup> <sup>110</sup> <sup>111</sup> <sup>112</sup> <sup>113</sup> <sup>114</sup> <sup>115</sup> <sup>116</sup> <sup>117</sup> <sup>118</sup> <sup>119</sup> <sup>120</sup> <sup>121</sup> <sup>122</sup> <sup>123</sup> <sup>124</sup> <sup>125</sup> <sup>126</sup> <sup>127</sup> <sup>128</sup> <sup>129</sup> <sup>130</sup> <sup>131</sup> <sup>132</sup> <sup>133</sup> <sup>134</sup> <sup>135</sup> <sup>136</sup> <sup>137</sup> <sup>138</sup> <sup>139</sup> <sup>140</sup> <sup>141</sup> <sup>142</sup> <sup>143</sup> <sup>144</sup> <sup>145</sup> <sup>146</sup> <sup>147</sup> <sup>148</sup> <sup>149</sup> <sup>150</sup> <sup>151</sup> <sup>152</sup> <sup>153</sup> <sup>154</sup> <sup>155</sup> <sup>156</sup> <sup>157</sup> <sup>158</sup> <sup>159</sup> <sup>160</sup> <sup>161</sup> <sup>162</sup> <sup>163</sup> <sup>164</sup> <sup>165</sup> <sup>166</sup> <sup>167</sup> <sup>168</sup> <sup>169</sup> <sup>170</sup> <sup>171</sup> <sup>172</sup> <sup>173</sup> <sup>174</sup> <sup>175</sup> <sup>176</sup> <sup>177</sup> <sup>178</sup> <sup>179</sup> <sup>180</sup> <sup>181</sup> <sup>182</sup> <sup>183</sup> <sup>184</sup> <sup>185</sup> <sup>186</sup> <sup>187</sup> <sup>188</sup> <sup>189</sup> <sup>190</sup> <sup>191</sup> <sup>192</sup> <sup>193</sup> <sup>194</sup> <sup>195</sup> <sup>196</sup> <sup>197</sup> <sup>198</sup> <sup>199</sup> <sup>200</sup> <sup>201</sup> <sup>202</sup> <sup>203</sup> <sup>204</sup> <sup>205</sup> <sup>206</sup> <sup>207</sup> <sup>208</sup> <sup>209</sup> <sup>210</sup> <sup>211</sup> <sup>212</sup> <sup>213</sup> <sup>214</sup> <sup>215</sup> <sup>216</sup> <sup>217</sup> <sup>218</sup> <sup>219</sup> <sup>220</sup> <sup>221</sup> <sup>222</sup> <sup>223</sup> <sup>224</sup> <sup>225</sup> <sup>226</sup> <sup>227</sup> <sup>228</sup> <sup>229</sup> <sup>230</sup> <sup>231</sup> <sup>232</sup> <sup>233</sup> <sup>234</sup> <sup>235</sup> <sup>236</sup> <sup>237</sup> <sup>238</sup> <sup>239</sup> <sup>240</sup> <sup>241</sup> <sup>242</sup> <sup>243</sup> <sup>244</sup> <sup>245</sup> <sup>246</sup> <sup>247</sup> <sup>248</sup> <sup>249</sup> <sup>250</sup> <sup>251</sup> <sup>252</sup> <sup>253</sup> <sup>254</sup> <sup>255</sup> <sup>256</sup> <sup>257</sup> <sup>258</sup> <sup>259</sup> <sup>260</sup> <sup>261</sup> <sup>262</sup> <sup>263</sup> <sup>264</sup> <sup>265</sup> <sup>266</sup> <sup>267</sup> <sup>268</sup> <sup>269</sup> <sup>270</sup> <sup>271</sup> <sup>272</sup> <sup>273</sup> <sup>274</sup> <sup>275</sup> <sup>276</sup> <sup>277</sup> <sup>278</sup> <sup>279</sup> <sup>280</sup> <sup>281</sup> <sup>282</sup> <sup>283</sup> <sup>284</sup> <sup>285</sup> <sup>286</sup> <sup>287</sup> <sup>288</sup> <sup>289</sup> <sup>290</sup> <sup>291</sup> <sup>292</sup> <sup>293</sup> <sup>294</sup> <sup>295</sup> <sup>296</sup> <sup>297</sup> <sup>298</sup> <sup>299</sup> <sup>300</sup> <sup>301</sup> <sup>302</sup> <sup>303</sup> <sup>304</sup> <sup>305</sup> <sup>306</sup> <sup>307</sup> <sup>308</sup> <sup>309</sup> <sup>310</sup> <sup>311</sup> <sup>312</sup> <sup>313</sup> <sup>314</sup> <sup>315</sup> <sup>316</sup> <sup>317</sup> <sup>318</sup> <sup>319</sup> <sup>320</sup> <sup>321</sup> <sup>322</sup> <sup>323</sup> <sup>324</sup> <sup>325</sup> <sup>326</sup> <sup>327</sup> <sup>328</sup> <sup>329</sup> <sup>330</sup> <sup>331</sup> <sup>332</sup> <sup>333</sup> <sup>334</sup> <sup>335</sup> <sup>336</sup> <sup>337</sup> <sup>338</sup> <sup>339</sup> <sup>340</sup> <sup>341</sup> <sup>342</sup> <sup>343</sup> <sup>344</sup> <sup>345</sup> <sup>346</sup> <sup>347</sup> <sup>348</sup> <sup>349</sup> <sup>350</sup> <sup>351</sup> <sup>352</sup> <sup>353</sup> <sup>354</sup> <sup>355</sup> <sup>356</sup> <sup>357</sup> <sup>358</sup> <sup>359</sup> <sup>360</sup> <sup>361</sup> <sup>362</sup> <sup>363</sup> <sup>364</sup> <sup>365</sup> <sup>366</sup> <sup>367</sup> <sup>368</sup> <sup>369</sup> <sup>370</sup> <sup>371</sup> <sup>372</sup> <sup>373</sup> <sup>374</sup> <sup>375</sup> <sup>376</sup> <sup>377</sup> <sup>378</sup> <sup>379</sup> <sup>380</sup> <sup>381</sup> <sup>382</sup> <sup>383</sup> <sup>384</sup> <sup>385</sup> <sup>386</sup> <sup>387</sup> <sup>388</sup> <sup>389</sup> <sup>390</sup> <sup>391</sup> <sup>392</sup> <sup>393</sup> <sup>394</sup> <sup>395</sup> <sup>396</sup> <sup>397</sup> <sup>398</sup> <sup>399</sup> <sup>400</sup> <sup>401</sup> <sup>402</sup> <sup>403</sup> <sup>404</sup> <sup>405</sup> <sup>406</sup> <sup>407</sup> <sup>408</sup> <sup>409</sup> <sup>410</sup> <sup>411</sup> <sup>412</sup> <sup>413</sup> <sup>414</sup> <sup>415</sup> <sup>416</sup> <sup>417</sup> <sup>418</sup> <sup>419</sup> <sup>420</sup> <sup>421</sup> <sup>422</sup> <sup>423</sup> <sup>424</sup> <sup>425</sup> <sup>426</sup> <sup>427</sup> <sup>428</sup> <sup>429</sup> <sup>430</sup> <sup>431</sup> <sup>432</sup> <sup>433</sup> <sup>434</sup> <sup>435</sup> <sup>436</sup> <sup>437</sup> <sup>438</sup> <sup>439</sup> <sup>440</sup> <sup>441</sup> <sup>442</sup> <sup>443</sup> <sup>444</sup> <sup>445</sup> <sup>446</sup> <sup>447</sup> <sup>448</sup> <sup>449</sup> <sup>450</sup> <sup>451</sup> <sup>452</sup> <sup>453</sup> <sup>454</sup> <sup>455</sup> <sup>456</sup> <sup>457</sup> <sup>458</sup> <sup>459</sup> <sup>460</sup> <sup>461</sup> <sup>462</sup> <sup>463</sup> <sup>464</sup> <sup>465</sup> <sup>466</sup> <sup>467</sup> <sup>468</sup> <sup>469</sup> <sup>470</sup> <sup>471</sup> <sup>472</sup> <sup>473</sup> <sup>474</sup> <sup>475</sup> <sup>476</sup> <sup>477</sup> <sup>478</sup> <sup>479</sup> <sup>480</sup> <sup>481</sup> <sup>482</sup> <sup>483</sup> <sup>484</sup> <sup>485</sup> <sup>486</sup> <sup>487</sup> <sup>488</sup> <sup>489</sup> <sup>490</sup> <sup>491</sup> <sup>492</sup> <sup>493</sup> <sup>494</sup> <sup>495</sup> <sup>496</sup> <sup>497</sup> <sup>498</sup> <sup>499</sup> <sup>500</sup> <sup>501</sup> <sup>502</sup> <sup>503</sup> <sup>504</sup> <sup>505</sup> <sup>506</sup> <sup>507</sup> <sup>508</sup> <sup>509</sup> <sup>510</sup> <sup>511</sup> <sup>512</sup> <sup>513</sup> <sup>514</sup> <sup>515</sup> <sup>516</sup> <sup>517</sup> <sup>518</sup> <sup>519</sup> <sup>520</sup> <sup>521</sup> <sup>522</sup> <sup>523</sup> <sup>524</sup> <sup>525</sup> <sup>526</sup> <sup>527</sup> <sup>528</sup> <sup>529</sup> <sup>530</sup> <sup>531</sup> <sup>532</sup> <sup>533</sup> <sup>534</sup> <sup>535</sup> <sup>536</sup> <sup>537</sup> <sup>538</sup> <sup>539</sup> <sup>540</sup> <sup>541</sup> <sup>542</sup> <sup>543</sup> <sup>544</sup> <sup>545</sup> <sup>546</sup> <sup>547</sup> <sup>548</sup> <sup>549</sup> <sup>550</sup> <sup>551</sup> <sup>552</sup> <sup>553</sup> <sup>554</sup> <sup>555</sup> <sup>556</sup> <sup>557</sup> <sup>558</sup> <sup>559</sup> <sup>560</sup> <sup>561</sup> <sup>562</sup> <sup>563</sup> <sup>564</sup> <sup>565</sup> <sup>566</sup> <sup>567</sup> <sup>568</sup> <sup>569</sup> <sup>570</sup> <sup>571</sup> <sup>572</sup> <sup>573</sup> <sup>574</sup> <sup>575</sup> <sup>576</sup> <sup>577</sup> <sup>578</sup> <sup>579</sup> <sup>580</sup> <sup>581</sup> <sup>582</sup> <sup>583</sup> <sup>584</sup> <sup>585</sup> <sup>586</sup> <sup>587</sup> <sup>588</sup> <sup>589</sup> <sup>590</sup> <sup>591</sup> <sup>592</sup> <sup>593</sup> <sup>594</sup> <sup>595</sup> <sup>596</sup> <sup>597</sup> <sup>598</sup> <sup>599</sup> <sup>600</sup> <sup>601</sup> <sup>602</sup> <sup>603</sup> <sup>604</sup> <sup>605</sup> <sup>606</sup> <sup>607</sup> <sup>608</sup> <sup>609</sup> <sup>610</sup> <sup>611</sup> <sup>612</sup> <sup>613</sup> <sup>614</sup> <sup>615</sup> <sup>616</sup> <sup>617</sup> <sup>618</sup> <sup>619</sup> <sup>620</sup> <sup>621</sup> <sup>622</sup> <sup>623</sup> <sup>624</sup> <sup>625</sup> <sup>626</sup> <sup>627</sup> <sup>628</sup> <sup>629</sup> <sup>630</sup> <sup>631</sup> <sup>632</sup> <sup>633</sup> <sup>634</sup> <sup>635</sup> <sup>636</sup> <sup>637</sup> <sup>638</sup> <sup>639</sup> <sup>640</sup> <sup>641</sup> <sup>642</sup> <sup>643</sup> <sup>644</sup> <sup>645</sup> <sup>646</sup> <sup>647</sup> <sup>648</sup> <sup>649</sup> <sup>650</sup> <sup>651</sup> <sup>652</sup> <sup>653</sup> <sup>654</sup> <sup>655</sup> <sup>656</sup> <sup>657</sup> <sup>658</sup> <sup>659</sup> <sup>660</sup> <sup>661</sup> <sup>662</sup> <sup>663</sup> <sup>664</sup> <sup>665</sup> <sup>666</sup> <sup>667</sup> <sup>668</sup> <sup>669</sup> <sup>670</sup> <sup>671</sup> <sup>672</sup> <sup>673</sup> <sup>674</sup> <sup>675</sup> <sup>676</sup> <sup>677</sup> <sup>678</sup> <sup>679</sup> <sup>680</sup> <sup>681</sup> <sup>682</sup> <sup>683</sup> <sup>684</sup> <sup>685</sup> <sup>686</sup> <sup>687</sup> <sup>688</sup> <sup>689</sup> <sup>690</sup> <sup>691</sup> <sup>692</sup> <sup>693</sup> <sup>694</sup> <sup>695</sup> <sup>696</sup> <sup>697</sup> <sup>698</sup> <sup>699</sup> <sup>700</sup> <sup>701</sup> <sup>702</sup> <sup>703</sup> <sup>704</sup> <sup>705</sup> <sup>706</sup> <sup>707</sup> <sup>708</sup> <sup>709</sup> <sup>710</sup> <sup>711</sup> <sup>712</sup> <sup>713</sup> <sup>714</sup> <sup>715</sup> <sup>716</sup> <sup>717</sup> <sup>718</sup> <sup>719</sup> <sup>720</sup> <sup>721</sup> <sup>722</sup> <sup>723</sup> <sup>724</sup> <sup>725</sup> <sup>726</sup> <sup>727</sup> <sup>728</sup> <sup>729</sup> <sup>730</sup> <sup>731</sup> <sup>732</sup> <sup>733</sup> <sup>734</sup> <sup>735</sup> <sup>736</sup> <sup>737</sup> <sup>738</sup> <sup>739</sup> <sup>740</sup> <sup>741</sup> <sup>742</sup> <sup>743</sup> <sup>744</sup> <sup>745</sup> <sup>746</sup> <sup>747</sup> <sup>748</sup> <sup>749</sup> <sup>750</sup> <sup>751</sup> <sup>752</sup> <sup>753</sup> <sup>754</sup> <sup>755</sup> <sup>756</sup> <sup>757</sup> <sup>758</sup> <sup>759</sup> <sup>760</sup> <sup>761</sup> <sup>762</sup> <sup>763</sup> <sup>764</sup> <sup>765</sup> <sup>766</sup> <sup>767</sup> <sup>768</sup> <sup>769</sup> <sup>770</sup> <sup>771</sup> <sup>772</sup> <sup>773</sup> <sup>774</sup> <sup>775</sup> <sup>776</sup> <sup>777</sup> <sup>778</sup> <sup>779</sup> <sup>780</sup> <sup>781</sup> <sup>782</sup> <sup>783</sup> <sup>784</sup> <sup>785</sup> <sup>786</sup> <sup>787</sup> <sup>788</sup> <sup>789</sup> <sup>790</sup> <sup>791</sup> <sup>792</sup> <sup>793</sup> <sup>794</sup> <sup>795</sup> <sup>796</sup> <sup>797</sup> <sup>798</sup> <sup>799</sup> <sup>800</sup> <sup>801</sup> <sup>802</sup> <sup>803</sup> <sup>804</sup> <sup>805</sup> <sup>806</sup> <sup>807</sup> <sup>808</sup> <sup>809</sup> <sup>810</sup> <sup>811</sup> <sup>812</sup> <sup>813</sup> <sup>814</sup> <sup>815</sup> <sup>816</sup> <sup>817</sup> <sup>818</sup> <sup>819</sup> <sup>820</sup> <sup>821</sup> <sup>822</sup> <sup>823</sup> <sup>824</sup> <sup>825</sup> <sup>826</sup> <sup>827</sup> <sup>828</sup> <sup>829</sup> <sup>830</sup> <sup>831</sup> <sup>832</sup> <sup>833</sup> <sup>834</sup> <sup>835</sup> <sup>836</sup> <sup>837</sup> <sup>838</sup> <sup>839</sup> <sup>840</sup> <sup>841</sup> <sup>842</sup> <sup>843</sup> <sup>844</sup> <sup>845</sup> <sup>846</sup> <sup>847</sup> <sup>848</sup> <sup>849</sup> <sup>850</sup> <sup>851</sup> <sup>852</sup> <sup>853</sup> <sup>854</sup> <sup>855</sup> <sup>856</sup> <sup>857</sup> <sup>858</sup> <sup>859</sup> <sup>860</sup> <sup>861</sup> <sup>862</sup> <sup>863</sup> <sup>864</sup> <sup>865</sup> <sup>866</sup> <sup>867</sup> <sup>868</sup> <sup>869</sup> <sup>870</sup> <sup>871</sup> <sup>872</sup> <sup>873</sup> <sup>874</sup> <sup>875</sup> <sup>876</sup> <sup>877</sup> <sup>878</sup> <sup>879</sup> <sup>880</sup> <sup>881</sup> <sup>882</sup> <sup>883</sup> <sup>884</sup> <sup>885</sup> <sup>886</sup> <sup>887</sup> <sup>888</sup> <sup>889</sup> <sup>890</sup> <sup>891</sup> <sup>892</sup> <sup>893</sup> <sup>894</sup> <sup>895</sup> <sup>896</sup> <sup>897</sup> <sup>898</sup> <sup>899</sup> <sup>900</sup> <sup>901</sup> <sup>902</sup> <sup>903</sup> <sup>904</sup> <sup>905</sup> <sup>906</sup> <sup>907</sup> <sup>908</sup> <sup>909</sup> <sup>910</sup> <sup>911</sup> <sup>912</sup> <sup>913</sup> <sup>914</sup> <sup>915</sup> <sup>916</sup> <sup>917</sup> <sup>918</sup> <sup>919</sup> <sup>920</sup> <sup>921</sup> <sup>922</sup> <sup>923</sup> <sup>924</sup> <sup>925</sup> <sup>926</sup> <sup>927</sup> <sup>928</sup> <sup>929</sup> <sup>930</sup> <sup>931</sup> <sup>932</sup> <sup>933</sup> <sup>934</sup> <sup>935</sup> <sup>936</sup> <sup>937</sup> <sup>938</sup> <sup>939</sup> <sup>940</sup> <sup>941</sup> <sup>942</sup> <sup>943</sup> <sup>944</sup> <sup>945</sup> <sup>946</sup> <sup>947</sup> <sup>948</sup> <sup>949</sup> <sup>950</sup> <sup>951</sup> <sup>952</sup> <sup>953</sup> <sup>954</sup> <sup>955</sup> <sup>956</sup> <sup>957</sup> <sup>958</sup> <sup>959</sup> <sup>960</sup> <sup>961</sup> <sup>962</sup> <sup>963</sup> <sup>964</sup> <sup>965</sup> <sup>966</sup> <sup>967</sup> <sup>968</sup> <sup>969</sup> <sup>970</sup> <sup>971</sup> <sup>972</sup> <sup>973</sup> <sup>974</sup> <sup>975</sup> <sup>976</sup> <sup>977</sup> <sup>978</sup> <sup>979</sup> <sup>980</sup> <sup>981</sup> <sup>982</sup> <sup>983</sup> <sup>984</sup> <sup>985</sup> <sup>986</sup> <sup>987</sup> <sup>988</sup> <sup>989</sup> <sup>990</sup> <sup>991</sup> <sup>992</sup> <sup>993</sup> <sup>994</sup> <sup>995</sup> <sup>996</sup> <sup>997</sup> <sup>998</sup> <sup>999</sup> <sup>1000</sup> <sup>1001</sup> <sup>1002</sup> <sup>1003</sup> <sup>1004</sup> <sup>1005</sup> <sup>1006</sup> <sup>1007</sup> <sup>1008</sup> <sup>1009</sup> <sup>1010</sup> <sup>1011</sup> <sup>1012</sup> <sup>1013</sup> <sup>1014</sup> <sup>1015</sup> <sup>1016</sup> <sup>1017</sup> <sup>1018</sup> <sup>1019</sup> <sup>1020</sup> <sup>1021</sup> <sup>1022</sup> <sup>1023</sup> <sup>1024</sup> <sup>1025</sup> <sup>1026</sup> <sup>1027</sup> <sup>1028</sup> <sup>1029</sup> <sup>1030</sup> <sup>1031</sup> <sup>1032</sup> <sup>1033</sup> <sup>1034</sup> <sup>1035</sup> <sup>1036</sup> <sup>1037</sup> <sup>1038</sup> <sup>1039</sup> <sup>1040</sup> <sup>1041</sup> <sup>1042</sup> <sup>1043</sup> <sup>1044</sup> <sup>1045</sup> <sup>1046</sup> <sup>1047</sup> <sup>1048</sup> <sup>1049</sup> <sup>1050</sup> <sup>1051</sup> <sup>1052</sup> <sup>1053</sup> <sup>1054</sup> <sup>1055</sup> <sup>1056</sup> <sup>1057</sup> <sup>1058</sup> <sup>1059</sup> <sup>1060</sup> <sup>1061</sup> <sup>1062</sup> <sup>1063</sup> <sup>1064</sup> <sup>1065</sup> <sup>1066</sup> <sup>1067</sup> <sup>1068</sup> <sup>1069</sup> <sup>1070</sup> <sup>1071</sup> <sup>1072</sup> <sup>1073</sup> <sup>1074</sup> <sup>1075</sup> <sup>1076</sup> <sup>1077</sup> <sup>1078</sup> <sup>1079</sup> <sup>1080</sup> <sup>1081</sup> <sup>1082</sup> <sup>1083</sup> <sup>1084</sup> <sup>1085</sup> <sup>1086</sup> <sup>1087</sup> <sup>1088</sup> <sup>1089</sup> <sup>1090</sup> <sup>1091</sup> <sup>1092</sup> <sup>1093</sup> <sup>1094</sup> <sup>1095</sup> <sup>1096</sup> <sup>1097</sup> <sup>1098</sup> <sup>1099</sup> <sup>1100</sup> <sup>1101</sup> <sup>1102</sup> <sup>1103</sup> <sup>1104</sup> <sup>1105</sup> <sup>1106</sup> <sup>1107</sup> <sup>1108</sup> <sup>1109</sup> <sup>1110</sup> <sup>1111</sup> <sup>1112</sup> <sup>1113</sup> <sup>1114</sup> <sup>1115</sup> <sup>1116</sup> <sup>1117</sup> <sup>1118</sup> <sup>1119</sup> <sup>1120</sup> <sup>1121</sup> <sup>1122</sup> <sup>1123</sup> <sup>1124</sup> <sup>1125</sup> <sup>1126</sup> <sup>1127</sup> <sup>1128</sup> <sup>1129</sup> <sup>1130</sup> <sup>1131</sup> <sup>1132</sup> <sup>1133</sup> <sup>1134</sup> <sup>1135</sup> <sup>1136</sup> <sup>1137</sup> <sup>1138</sup> <sup>1139</sup> <sup>1140</sup> <sup>1141</sup> <sup>1142</sup> <sup>1143</sup> <sup>1144</sup> <sup>1145</sup> <sup>1146</sup> <sup>1147</sup> <sup>1148</sup> <sup>1149</sup> <sup>1150</sup> <sup>1151</sup> <sup>1152</sup> <sup>1153</sup> <sup>1154</sup> <sup>1155</sup> <sup>1156</sup> <sup>1157</sup> <sup>1158</sup> <sup>1159</sup> <sup>1160</sup> <sup>1161</sup> <sup>1162</sup> <sup>1163</sup> <sup>1164</sup> <sup>1165</sup> <sup>1166</sup> <sup>1167</sup> <sup>1168</sup> <sup>1169</sup> <sup>1170</sup> <sup>1171</sup> <sup>1172</sup> <sup>1173</sup> <sup>1174</sup> <sup>1175</sup> <sup>1176</sup> <sup>1177</sup> <sup>1178</sup> <sup>1179</sup> <sup>1180</sup> <sup>1181</sup> <sup>1182</sup> <sup>1183</sup> <sup>1184</sup> <sup>1185</sup> <sup>1186</sup> <sup>1187</sup> <sup>1188</sup> <sup>1189</sup> <sup>1190</sup> <sup>1191</sup> <sup>1192</sup> <sup>1193</sup> <sup>1194</sup> <sup>1195</sup> <sup>1196</sup> <sup>1197</sup> <sup>1198</sup> <sup>1199</sup> <sup>1200</sup> <sup>1201</sup> <sup>1202</sup> <sup>1203</sup> <sup>1204</sup> <sup>1205</sup> <sup>1206</sup> <sup>1207</sup> <sup>1208</sup> <sup>1209</sup> <sup>1210</sup> <sup>1211</sup> <sup>1212</sup> <sup>1213</sup> <sup>1214</sup> <sup>1215</sup> <sup>1216</sup> <sup>1217</sup> <sup>1218</sup> <sup>1219</sup> <sup>1220</sup> <sup>1221</sup> <sup>1222</sup> <sup>1223</sup> <sup>1224</sup> <sup>1225</sup> <sup>1226</sup> <sup>1227</sup> <sup>1228</sup> <sup>1229</sup> <sup>1230</sup> <sup>1231</sup> <sup>1232</sup> <sup>1233</sup> <sup>1234</sup> <sup>1235</sup> <sup>1236</sup> <sup>1237</sup> <sup>1238</sup> <sup>1239</sup> <sup>1240</sup> <sup>1241</sup> <sup>1242</sup> <sup>1243</sup> <sup>1244</sup> <sup>1245</sup> <sup>1246</sup> <sup>1247</sup> <sup>1248</sup> <sup>1249</sup> <sup>1250</sup> <sup>1251</sup> <sup>1252</sup> <sup>1253</sup> <sup>1254</sup> <sup>1255</sup> <sup>1256</sup> <sup>1257</sup> <sup>1258</sup> <sup>1259</sup> <sup>1260</sup> <sup>1261</sup> <sup>1262</sup>





FIG 203 Basal cell papilloma keratotic type High magnification Thick interoven tracts of basal cells compose the tumor Interspersed are cystic inclusions of horny material caused by invaginations of the horny layer ( $\times 100$ )




FIG 201 Basal cell papilloma adenoid type Thin interoven tracts composed of a double row of basal cells compose the tumor No cystic inclusions of horny material are present ( $\times 100$ )

branch and interweave (Fig 204). Considerable amounts of melanin usually are present in the tracts. Cystic inclusions of horny material are absent. The demarcation at the lower border is sharp just as in the keratotic type of basal cell papilloma.

Although basal-cell papilloma shows no tendency to invasion of the dermis it may be regarded as related to basal cell epithelioma because a large proportion of the cells are of the same type as found in basal cell epithelioma. However actual change of basal cell papilloma into a frank basal-cell epithelioma with deep invasion and ulceration is rare (Filler and Ryan Pinkus).

(For differential diagnosis from nevus verrucosus see page 322 and from keratosis senilis see page 328.)

### DERMATOSIS PAPULOSA NIGRA

This condition is frequent among Negroes especially females. The lesions are located on the face especially in the malar regions and consist of minute soft rounded usually hyperpigmented papules.

Histopathology. The histologic changes are the same as in basal cell papilloma. Melanin pigmentation of the basal layer is pronounced (Michael and Seale).

### CARCINOMA OF SEBACEOUS GLANDS

Carcinomas of sebaceous glands have been described occasionally in the literature. Some authors are more inclined to make this diagnosis than others. Warren and Warren for instance reported 28 personally observed cases while most other authors have reported only single instances (Beach and Severance).

Carcinomas of sebaceous glands occur most frequently on the eyelids, originating from meibomian glands which are modified sebaceous glands (Hagedoorn) they may however occur anywhere on the skin (Beach and Severance). No characteristic

conserves definite lobular formations (Fig 205). Although many cells are undifferentiated distinct sebaceous cells are present. Mitotic figures are numerous. Lipid material can be demonstrated not only in the sebaceous cells but also as fine globules in many other cells of the tumor. Beach and Severance state that the undifferentiated cells of sebaceous carcinoma differ from those of basal cell epithelioma by showing greater variation in size and shape and by having a more acidophilic cytoplasm and a lighter staining nucleus.

It seems that many cases reported as sebaceous carcinoma may be regarded as sebaceous adenoma (see page 347) or as sebaceous epithelioma, which is a basal cell epithelioma with considerable differentiation of the cells toward sebaceous gland cells (page 353). For in



FIG. 205. Carcinoma of sebaceous glands. Closely packed lobular formations are present in the dermis. They are composed of sebaceous and undifferentiated cells. Some of the latter are atypical ( $\times 200$ ).

stance, the five cases of sebaceous carcinoma reported by Savataid as arising in cases of nevus sebaceus are best classified as instances of sebaceous epithelioma.

### CARCINOMA OF ECCRINE SWEAT GLANDS

Carcinomas of the eccrine sweat glands are rare, but their existence seems to be established definitely. They do not possess a characteristic appearance, either clinically or histologically.

**Histopathology.** It is difficult to recognize carcinoma of eccrine sweat glands and to differentiate it from other cutaneous tumors,


carcinomas of sweat glands  
cystic papillary and solid

forms. He pointed out that combinations of these forms were frequent. He recognized the occasional occurrence of metaplasia of glandular cells into squamous and horn cells.

#### ADENOCANTHOMA OF SWEAT GLANDS

Carcinomas of sweat glands with glandular as well as epidermal elements have recently been reported under the designation adenocanthoma of sweat glands (Lever).

Clinically, these tumors resemble squamous-cell carcinoma. They show shallow central ulceration and may have a verrucous surface. Their site of predilection is the face, especially the ears.



ADENOCANTHOMA OF SWEAT GLANDS

... the alveolar formations, an island of tumor tissue and several atypical sweat glands or ducts are present. (X50)

**Histopathology.** Adenocanthomas show tubular and alveolar lumina lined with one or several layers of epithelium (Fig 206). In areas where the lumina are lined with a single layer of epithelium the epithelial cells have the appearance of glandular cells, but in areas with several layers of epithelium, squamous and partially keratinized cells usually form the inner layers. The lumina are filled with



FIG 207 Adenocanthoma of sweat glands. High magnification of the atypical sweat glands or ducts seen in Figure 206. Papillary tufts protrude into their lumina. These sweat glands approach in appearance the glandular structures belonging to the tumor ( $\times 200$ ).

desquamated cells, many of which are partially or completely keratinized. In addition, there are solid areas which have the appearance of squamous cell carcinoma. Atypical eccrine sweat glands and sweat ducts are present within or at the periphery of these tumors (Fig 207).

**Histogenesis.** The author originally regarded these tumors as sweat gland carcinomas composed of glandular and epidermal elements and explained the presence of these two structures by the fact that sweat ducts are composed of squamous cells in their epidermal portion, and of glandular cells in the dermis. It is possible, however, that these tumors are squamous cell carcinomas of alveolar growth in which there is considerable individual cell keratinization resulting in acantholysis in the center of the alveolar formations.

Borelli regards these tumors as dyskeratotic squamous-cell carcinomas which arise from the hair follicle and tend to imitate the structure of glands which develop embryologically together with the follicle

### CARCINOMA OF APOCRINE GLANDS

Extramammary Paget's disease represents a carcinoma of apocrine glands (see page 339). No other form of carcinoma of apocrine glands has so far been established

### METASTATIC CARCINOMA OF THE SKIN

Cutaneous metastasis of carcinoma is rare with the exception of carcinoma of the breast. Next to carcinoma of the breast metastases




FIG. 208 Metastatic carcinoma of the skin from carcinoma of the breast. Inflammatory carcinoma. The dermal lymphatics are filled with clusters of tumor cells ( $\times 100$ )

to the skin are observed most frequently in carcinoma of the stomach, the uterus, the lungs, the large intestines and the kidneys in that order (Gates). In occasional instances cancer of the

Dissemination may take place through the lymphatics or through the blood stream. In carcinoma of the breast cutaneous metastases

**Histopathology** Adenocanthomas show tubular and alveolar lumina lined with one or several layers of epithelium (Fig 206). In areas where the lumina are lined with a single layer of epithelium the epithelial cells have the appearance of glandular cells but in areas with several layers of epithelium squamous and partially keratinized cells usually form the inner layers. The lumina are filled with




FIG 207 Adenocanthoma of sweat glands. High magnification of the atypical sweat glands or ducts seen in Figure 206. Epillary tufts protrude into their lumen. These sweat glands approach in appearance the glandular structures belonging to the tumor ( $\times 200$ ).

desquamated cells many of which are partially or completely keratinized. In addition there are solid areas which have the appearance of squamous cell carcinoma. Atypical eccrine sweat glands and sweat ducts are present within or at the periphery of these tumors (Fig 207).

**Histogenesis** The author originally regarded these tumors as sweat gland carcinomas composed of glandular and epidermal elements and explained the presence of these two structures by the fact that sweat ducts are composed of squamous cells in their epidermal portion and of glandular cells in the dermis. It is possible however that these tumors are squamous-cell carcinomas of alveolar growth in which there is considerable individual cell keratinization resulting in acantholysis in the center of the alveolar formations.

In inflammatory carcinoma the skin of the affected breast and the adjoining areas present erythema and diffuse edema simulating erysipelas. In telangiectatic carcinoma the skin contains numerous purplish papules and hemorrhagic pseudovesicles resembling hemolymphangioma. In cancer en cuirasse the skin of the breast affected by the carcinoma and often also the surrounding skin shows dif-



FIG 210 Metastatic carcinoma of the skin from carcinoma of the breast Cancer en cuirasse, in ulcerated area. Only few tumor cells are present. They are embedded between collagen bundles in single rows. Because of their small number and size they may be overlooked easily. ( $\times 400$ )

fuse brownish induration interspersed with nodules and punched out ulcerations.

**Histopathology** In inflammatory carcinoma histologic examination of the skin reveals extensive invasion of the dermal lymphatics—especially of the subepidermal lymphatics—by groups and cords of tumor cells (Fig 208) (Taylor and Meltzer). The tumor cells are similar to those of the primary growth atypical in character with large rounded deeply staining nuclei and a moderate amount of cytoplasm. Occasional mitoses are seen in these cells. There is marked capillary congestion (which is the reason for the inflammatory appearance clinically). In addition one observes edema and a slight perivascular lymphocytic infiltrate in the dermis but no fibrosis (Pfahler and Case).



usually are caused by way of the lymphatics. In other carcinomas, cutaneous metastases, as a rule, are caused by dissemination through the blood stream, although occasionally dissemination takes place through lymphatics secondary to local lymph node involvement (Gates)



FIG. 209 Metastatic carcinoma of the skin from carcinoma of the breast. *Cancer en cuirasse* nodular lesion. There is considerable fibrosis of the dermis with small scattered islands of tumor cells. Some of the islands show a suggestive glandular arrangement of the tumor cells. ( $\times 200$ )

#### CUTANEOUS METASTASIS FROM CARCINOMA OF THE BREAST

Three types of cutaneous metastases may occur in carcinoma of the breast: inflammatory carcinoma, telangiectatic carcinoma, and cancer en cuirasse. Two or all three of these types may be present in the same patient. If dissemination of metastases through the lymphatics proceeds rapidly, inflammatory carcinoma results in most instances, and telangiectatic carcinoma in rare instances. If dissemination proceeds slowly, cancer en cuirasse eventuates (Taylor and Meltzer).

**Histopathology** Large and small groups of tumor cells are present throughout the dermis. As a rule no tumor cells are found within capillaries or lymphatics. A certain amount of fibrosis is usually present. Signs of inflammation are absent (Gates).

The metastatic carcinoma readily can be classified as either an adenocarcinoma (Fig. 211), squamous-cell carcinoma or an undiffer-




FIG. 212 Metastatic carcinoma of the skin originating from the stomach. Many of the tumor cells are so-called signet ring cells in which on account of the presence of mucin in the cytoplasm the nucleus is pressed against the cell wall ( $\times 400$ ).

entiated carcinoma. However, it is only occasionally possible to draw conclusions from the histologic appearance of the metastasis as to the site of the primary tumor. For instance, if the tumor cells of the cutaneous metastasis contain mucin the primary carcinoma most likely resides in the gastro-intestinal tract. The mucin-containing cells present in the metastasis appear just as in the primary tumor as so-called signet ring cells—namely, as large round cells filled with mucin which presses the nucleus against the cellular wall (Fig. 212).

In melangiectatic carcinoma, the dilated lymphatics contain in addition to groups of tumor cells, varying amounts of erythrocytes (Freeman and Lynch, Leavell and Tillotson) The frequent location of these lymphatics immediately beneath the epidermis causes the clinical resemblance of the lesions to vesicles

In cancer en cuirasse, the nodular areas usually contain large and small groups as well as sheets of tumor cells lying outside of lymphatics in the dermis and surrounded by fibrosis (Fig 209) However, the indurated areas often contain only few tumor cells which therefore, may be easily overlooked The tumor cells are small angular and deeply basophilic and lie as single cells in single row lines or small groups between thickened collagen bundles The arrangement in single row lines, like Indians in a file, is of particular diagnostic importance (Fig 210)

#### CUTANEOUS METASTASIS FROM CARCINOMAS OTHER THAN BREAST CARCINOMA

The cutaneous metastases caused by carcinomas of organs other than the breast occur clinically as circumscribed nodules or tumors The nodules are usually discrete and moderately firm Their number may vary from one to several hundred




FIG. 211 Metastatic adenocarcinoma of the skin. Lumina & lumina are present ( $\times 100$ )

**Histopathology.** Large and small groups of tumor cells are present throughout the dermis. As a rule, no tumor cells are found within capillaries or lymphatics. A certain amount of fibrosis is usually present. Signs of inflammation are absent (Gates).

The metastatic carcinoma readily can be classified as either an adenocarcinoma (Fig 211), a squamous-cell carcinoma or an undiffer-




FIG 212. Metastatic carcinoma of the skin originating from the stomach. Many of the tumor cells are so-called signet ring cells in which on account of the presence of mucin in the cytoplasm the nucleus is pressed against the cell wall ( $\times 400$ ).

entiated carcinoma. However, it is only occasionally possible to draw conclusions from the histologic appearance of the metastasis as to the site of the primary tumor. For instance, if the tumor cells of the cutaneous metastasis contain mucin, the primary carcinoma most likely resides in the gastrointestinal tract. The mucin-containing cells present in the metastasis appear, just as in the primary, as so-called signet ring cells, in which the nucleus is pushed against the cell wall by the mucin.

## BIBLIOGRAPHY

## Histogenesis and Classification of Epidermal Tumors

- Albrecht I Ueber Hamartome Verhandl d Dtsch path Gesellsch 7 153 1901  
 Becker S W Benign epidermal neoplasms Arch Dermat & Syph 26 838 1932  
 Foulds L The histological analysis of tumors Am J Cancer 39 1 1940  
 Hansemann D Ueber asymmetrische Zelltheilung in Epithelkrebsen und deren biologische Bedeutung Virchows Arch 119 299 1890  
 Jadassohn J Die benignen Epitheliome Arch f Dermat u Syph 117 577 705 833 1913 1914  
 Lever W F Pathogenesis of benign tumors of cutaneous appendages and of basal cell epithelioma Arch Dermat & Syph 57 679 1948  
 Pinkus H Premalignant fibroepithelial tumors of the skin Arch Dermat & Syph 67 598 1953  
 Van der Valk J W Adénome sébacé ou épithéliome sébacé primaire Bull de l'Assn franç pour l'étude du cancer 13 465 1924

## Nevus Verrucosus

- Binzer E The generalized keratotic nevus Arch Dermat & Syph 21 413 1930

## Epidermal Cyst (Wen) Sebaceous Cyst Milium Dermoid Cyst

- Love W R and Montgomery H Epithelial cysts Arch Dermat & Syph 47 185 1913  
 New G B and Erich J B Dermoid cysts of the head and neck Surg Gynec & Obst 65 48 1937  
 Peden J C Carcinoma developing in sebaceous cysts Ann Surg 128 1136 1948  
 Warr W N and Gates O Epithelial cysts and cystic tumors of the skin Am J Path 14 765 1943

## Keratosis Senilis

- Freudenthal W Verruca senilis und Keratoma senile Arch f Dermat u Syph 152 505 1926  
 Halter K Ueber ein wenig beachtetes histologisches Kennzeichen des Keratoma senile Hautarzt 3 215 1952  
 Montgomery H and Dorfel J Verruca senilis und Keratoma senile Arch f Dermat u Syph 166 286 1932  
 Ruiter M An unusual case of keratoma senile Arch Dermat & Syph 33 807 1936  
 Szodoray L Histologic characteristics of the so called precancerous processes of the skin Arch Dermat & Syph 36 552 1937

## Leukoplakia

- Montgomery H Councillor V S and Craig W M Kraurosis leukoplakia and pruritus vulvae Arch Dermat & Syph 30 80 1934

## Squamous Cell Carcinoma and Pseudoepitheliomatous Hyperplasia

- Broders A C Squamous cell epithelioma of the skin Ann Surg 73 141 1921  
 (Good review)

- Broders A C Practical points on the microscopic grading of carcinoma New York State J Med 32 667 1932
- Brooks S M Carcinoma which resembles sarcoma Arch Path 36 141 1913
- Edmundson W F Microscopic grading of cancer and its practical implication  
the skin Acta dermat venereol

- G
- MacKee G M and Lipollaro A C Cutaneous Cancer and Precancer A Practical Monograph p 130 New York Am J Cancer 1937
- Montgomery H in discussion of Winer L H Pseudoepitheliomatous hyperplasia Arch Dermat & Syph 42 806 1910
- Sims C F and Kirsch N Spindle cell epidermoid epithelioma simulating sarcoma in chronic radiodermatitis Arch Dermat & Syph 57 63 1918
- Smith J F Multiple primary self healing squamous epithelioma of the skin Brit J Dermat 60 315 1918
- Sommerville J Pseudoepitheliomatous hyperplasia Acta dermat venereol 33 936 1953
- and Milne J A Familial primary self healing squamous epithelioma of the skin Brit J Dermat 62 185 1950
- Underwood L J Montgomery H and Broders A C Squamous cell epithelioma that simulates sarcoma Arch Dermat & Syph 64 149 1921
- Whimster I W in discussion of Musso L Spontaneous resolution of molluscum sebaceum Brit J Dermat 63 151 1921
- White C and Weidman F D Pseudo-epitheliomatous hyperplasia at the margin of cutaneous ulcers, J A M A 88 1979 1927
- Winer L H Pseudoepitheliomatous hyperplasia Arch Dermat & Syph 42 806 1910
- Witten V H and Zak F G Multiple primary self healing prickle cell epithelioma of the skin Cancer 5 939 1952

#### Bowen's Disease and Erythroplasia of Queyrat

- Bowen J T Precancerous dermatoses J Cutan Dis 30 241 1912 33 787 1915
- Kurnitzky E and Jacoby H Bowen's disease with metastases Arch Dermat & Syph 38 191 1938
- McDaniel W E and Mason L M Malignant dyskeratosis erythroplasia of Queyrat type Arch Dermat & Syph 60 419 1919
- Madsen A Contribution à la question de la nature neoplasique de la maladie de Bowen Acta dermat venereol 22 24 1931
- Montgomery H Precancerous dermatosis and epithelioma in situ Arch Dermat & Syph 39 387 1939
- Pautrier L M Erythroplasie du gland (Maladie de Bowen des muqueuses) Dermatolog 87 169 1943
- Sachs W and Sachs P Erythroplasia of Queyrat Arch Dermat & Syph 58 184 1918
- Sulzberger M B and Satenstein D L Erythroplasia of Queyrat Arch Dermat & Syph 28 798 1933
- Zoon J J Balanoposthène chronique circonscrite bénigne à plasmocytes Dermatolog 105 1 1952

#### Paget's Disease

- Allen A C A reorientation on the histogenesis and clinical significance of cutaneous nevi and melanomas Cancer 2 28 1919

- Foraker A G and Miller, C J Extramammary Paget's disease of perianal skin *Cancer* 2 141 1949
- Fraser J F Bowen's disease and Paget's disease of the nipple *Arch Dermat & Syph* 18 809 1928
- Inglis A Paget's disease of the nipple *Am J Path* 22 1 1946 (Good review)
- Marx R Significance of ductal sclerosis in Paget's disease *Arch Path* 47 401 1949
- Muir R Further observations on Paget's disease of the nipple *J Path & Bact* 49 299 1939
- Pirson L Extramammary Paget's disease *Arch Path* 36 324 1913
- Pautrier L M Paget's disease of the nipple *Arch Dermat & Syph* 17 767 1928
- Rubenstein M W Paget's disease of the male nipple and areola *Arch Dermat & Syph* 22 281 1930
- Stout A P The relationship of malignant melanotic melanoma (melanocarcinoma) to extramammary Paget's disease *Am J Cancer* 33 106 1935
- Whimster J W Ueber schleimsernierende Carcinome der Perianalregion *Hautarzt* 140 1953
- Zoon J J and Gelpke A S Findings in a case of extramammary Paget's disease *Dermatolog* 101 153 1950

#### Nevus Sebaceus (Jadassohn)

- Koch T Ueber das Vorkommen von apokrinen Drüsen in Trichomei *Arch Dermat u Syph* 174 126 1936
- Ostrowski S Beitrag zur Histologie des Nevus (epitheliomatosis) unilaterialis capitis *Acta dermat venercol* 8 235 1927
- Parkin T Naevus sebaceus (Jadassohn) with squamous cell epithelioma *Brit J Dermat* 62 167 1950
- Pautrier L M Le nevus sébacé de la face et du cuir chevelu l'épithélioma sébacé *Ann de dermat et de syph* 7 897 1936
- Robinson S S Naevus sebaceus (Jadassohn) *Arch Dermat & Syph* 26 663 1932
- Seward L Naevus sebaceus and sebaceous carcinoma *Brit J Dermat* 33 214 1941
- Szodory L Relations between nevus sebaceus and epithelioma *Acta dermat venercol* 13 1 1932

#### Adenoma Sebaceum (Pringle)

- Butterworth T and Wilson M Dermatologic aspects of tuberous sclerosis *Arch Dermat & Syph* 43 1 1941
- Good C B and Garb J Symmetric nevus of the face tuberous sclerosis epilepsy and fibromatous growth on scalp *Arch Dermat & Syph* 47 197 1943
- Kveim A Ueber Adenoma sebaceum (morbis Pringle) *Acta dermat venercol* 18 637 1937
- Moolten S F Hamman's nature of the tuberous sclerosis complex and its bearing on the tumor problem *Arch Int Med* 69 83 1942
- Pratt Thomas H R Tuberous sclerosis with congenital tumors of heart and kidney *Am J Path* 23 169 1917

#### Senile Sebaceous Nevus

- Gans O Histologie der Hautkrankheiten Vol 2 p 261 Berlin Julius Springer 1928

- Gilman R L Adenomatoid sebaceous tumors with particular reference to adenomatoid hyperplasia Arch Dermat & Syph 35 633 1937
- Woolhandler H W and Becker W S Adenoma of sebaceous glands (adenoma sebaceum) with consideration of keratotic adenoma sebaceum and true adenoma of sebaceous glands Arch Dermat & Syph 45 734 1942

#### Fordyce's Disease

- Chambers S O The structure of Fordyce's disease as demonstrated by wax reconstruction Arch Dermat & Syph 18 666 1908
- Margolies A and Weidman F Statistical and histologic studies of Fordyce's disease Arch Dermat & Syph 3 723 1921

#### Sebaceous Adenoma

- Grusfelt E Un cas d'épithélioma baso-sébacé Bull de l'Assoc Franç p l'étude du cancer 13 474 1921
- Lever W F Sebaceous adenoma review of the literature and report of a case Arch Dermat & Syph 57 102 1918
- Pautrier L M Le naevus sébacé de la face et du cuir chevelu l'épithélioma sébacé Ann de dermat et de syph 7 897 1936
- Reitmann K Zur Kenntnis der Talgdrüsen und der von ihnen ausgehenden Wucherungs- und Neubildungsprozesse Arch f Dermat u Syph 99 120 1909
- Woolhandler H W and Becker W S Adenoma of sebaceous glands (adenoma sebaceum) Arch Dermat & Syph 45 734 1942

#### Syringocystadenoma Papilliferum

- Appel B Nevus syringadenomatosus papilliferus Arch Dermat & Syph 61 311 1900
- D rffel J Zur Histogenese des Naevus syringocystadenomatosus papilliferus Dermat Wchnschr 101 851 1935
- Grund J L Syringocystadenoma papilliferum and nevus sebaceus (Jadassohn) occurring as a single tumor Arch Dermat. & Syph 65 340 1950
- Lever W F Pathogenesis of benign tumors of cutaneous appendages and of basal cell epithelioma Arch Dermat & Syph 57 679 1948
- Marcus M D and Woodruffe W E Sebaceous nevus and nevus syringo-adenomatosus papilliferus occurring as a mixed form Arch. Dermat & Syph 61 100 1900
- Reutersall O Naevus syringocystadenomatosus papilliferus and its relation to malignancy Acta path et microbiol scand 1933 supp 16 p 376
- Sachs W and Lewis G M Naevus syringadenomatosus papilliferus Arch Dermat. & Syph 36 1202 1937
- Tappeiner S Ueber das Vorkommen apokriner Drüsen in Naevusformationen Arch f Dermat u Syph 179 144 1939

#### Hydradenoma Papilliferum

- Anderson N P Hydradenoma of the vulva Arch Dermat & Syph 62 873 1900
- Danforth W C Hydradenoma vulvae Am J Obst & Gynec. 45 329 1943
- Cates O Warren S and Warr W A Tumors of the sweat glands Am J Path. 19 291 1943
- McDonald J R Apocrine sweat gland carcinoma of the vulva Am J Clin Path 11 890 1941



Winer L H in discussion of Anderson N P Hidradenoma of the vulva Arch Dermat & Syph 62 873 1950

### Sebaceous Epithelioma

Biberstein H Talgdrüsenkrebs und Epitheliom Arch f Dermat u Syph 117 177 1921

Grynfeltt L Un cas d'épithélioma sébacé Bull de l'Assoc franç pour l'étude du cancer 13 174 1921

Lever W F Pathogenesis of benign tumors of cutaneous appendages and of basal cell epithelioma Arch Dermat & Syph 57 679 1918

Ioos H O Die Carcinome der Anhangsgebilde der Haut Arch f Dermat u Syph 171 165 1936

Milhan Perin and Brunel L'épithélioma sébacé Bull Soc franç de dermat et syph 40 553 1933

### Syringoma

Ischer H Zur Genese epitheliärer Hauttumoren Dermat Wehnschr 83 1509 1926

Homma H and Ischer D H I Genesis of syringoma Arch Dermat & Syph 33 700 1936

Ingels A C Epithelioma adenoides cysticum with features of syringoma Arch Dermat & Syph 32 75 1935

Kyrle J Vorlesungen über Histo Biologie der menschlichen Haut und ihrer Erkrankungen vol I p 245 Berlin Julius Springer 1925

Lever W F Pathogenesis of benign tumors of cutaneous appendages and of basal cell epithelioma Arch Dermat & Syph 57 679 1918

McDonagh J L R Mixed tumor (trichoepithelioma papulosum and syringo cystadenoma) Brit J Dermat 22 20 1910

Riehl G Jr Zur Histogenese des Syringocystadenoms Wien Klin Wehnschr 48 209 1930

in a syringoma of

ch f Dermat u

### Cylindroma

Bilog P Beitrag zur Frage des Schweißdrüsenkrebs Dermat Wehnschr 82 841 1926

Cocern H Das Hidradenoma cylindromatosum der Kopfschwarte Beitr z klin Chir 93 205 1914 1915

Dwies J H T A contribution to the histology of the cylindromatous tumors of the scalp Brit J Dermat 40 211 1923

Lauecker H Beitrag zu den Nievo-epitheliomen Arch f Dermat u Syph 194 639 1952

Lever W F Pathogenesis of benign tumors of cutaneous appendages and of basal cell epithelioma Arch Dermat & Syph 57 679 1918

Luger A Das Cylindrom der Haut und seine maligne Degeneration Arch f Dermat u Syph 188 155 1919

Savard L Epithelioma adenoides cysticum Brit J Dermat 50 333 1938

Schlamadinger J Cylindrom und Trichoepithelioma papulosum multiplex Arch f Dermat u Syph 171 26 1935

Schuermann H and Weber K Beitrag zur Kenntnis der Spieglerischen Tumoren (Cylindrome) nebst einigen Bemerkungen zum Epithelioma adenoides cysticum Arch f Dermat u Syph 173 682 1937

- Stillians A W Naevus-epithelioma adenoides (cylindroma) of the scalp Arch Dermat & Syph 27 481 1933  
 Watanabe J Ueber das Cylindrom und das Epithelioma adenoides cysticum Arch f Dermat u Syph 110 208 1927  
 Zakon S J Naevus-epithelioma adenoides (cylindroma) of the scalp Arch Dermat. & Syph 40 945 1939

#### Myo-epithelioma Clear Cell Myo-epithelioma and Mixed Tumor of the Skin

- Balog P Beitrag zur Frage der Schweißdrüsenadenome Adenoma hydradenoides parenchymatosa cum lymphostasi Acta dermat venerol 22 187 1941  
 Hartz P H Adenomyoepithelioma of sweat gland Report of a case Am J Clin Path 16 383 1946  
 Highman B A mixed tumor of the salivary gland type on the left hand Arch Path 57 387 1941  
 Hirsch E F A subcutaneous mixed tumor (salivary gland type) of the left thigh Arch Path 16 494 1933  
 Lennox B Pearse A G E and Richards H G H Mucin secreting tumours of the skin and subcutaneous tissue

- Liu V The histogenesis of clear cell papillary carcinoma of the skin Am J Path 25 93 1949  
 Morehead R P Mixed tumors of the skin Arch Path 40 107 1945 (Good review)  
 Sheldon W B The myoepithelium in sweat gland tumors Arch Path 51 326 1941

#### Tricho-epithelioma

- Adamson H G On the nature of rodent ulcer Its relationship to epithelioma adenoides cysticum of Brooke and to other tricho-epitheliomata of benign nevus character its distinction from malignant carcinoma Lancet 1 810 1914  
 Goldman H J Multiple benign cystic epithelioma J.A.M.A 115 2253 1940  
 Hardy M H The histochemistry of hair follicles in the mouse Am J Anat 90 283 1930  
 Lever W F Pathogenesis of benign tumors of cutaneous appendages and of basal cell epithelioma Arch Dermat & Syph 57 679 1918  
 Little E C G Epithelioma adenoides cysticum of Brooke with rodent ulceration Brit J Dermat 51 193 1919  
 Savatard L Epithelioma adenoides cysticum Brit J Dermat 50 333 1938  
 Summerill F and Hutton J G Multiple benign cystic epithelioma (epithelioma adenoides cysticum) of the skin and basal cell cancer

#### Calcifying Epithelioma

- Cité F H Benign calcified epithelioma of the skin J Path & Bact 43 573 1936  
 Frick W Die verkalkenden Epitheliome der Haut und ihre Beziehungen zu Organisationsvorgängen in Atheromen Virchows Arch 289 527 1933

- Winer L H in discussion of Anderson N P Hidradenoma of the vulva Arch. Dermat & Syph 62 873 1950

### Sebaceous Epithelioma

- Biberstein H. Talgdrüsennevi und Epitheliom Arch f Dermat u Syph  
147 177 1924

- Grynfeldt E. Un cas d'epithélioma sébacé Bull de l'Assoc franç pour l'étude  
du cancer 13 474 1924

- Lever W F Pathogenesis of benign tumors of cutaneous appendages and of basal cell epithelioma *Arch Dermat & Syph* 57:679 1948

- Loos H O Die Carcinome der Anhangsgebilde der Haut Arch f Dermat u Syph 174 465 1936

- Milian Perin and Brunel Epithélioma sébacé Bull Soc franç de dermat et  
syph 40 553 1933

### Syringoma

- Fischer H Zur Genese epithelialer Hauttumoren Dermat Wchnschr 83 1909

- Homma H and Escher D H E. Genesis of syringoma Arch Dermat & Syph  
33 700 1936

- Ingels A E Epithelioma adenoides cysticum with features of syringoma Arch Dermat & Syph 32:75 1935

- Kytle J. Vorlesungen über Histo Biologie der menschlichen Haut und ihrer  
Erkrankungen vol 1 p 245 Berlin Julius Springer 1925

- Lever W F Pathogenesis of benign tumors of cutaneous appendages and of basal cell epithelioma Arch Dermat & Syph 57 679 1948

- McDonagh J E R Mixed tumor (trichoeplithelioma papulosum and syringo  
cystadenoma) Brit J Dermat 22:20 1910

- Riehl G Jr Zur Histogenese des Syringocystadenoms Wien klin Wchnschr  
48 209 1930

- Weidman F D and Besancon J H Histologic differences in a syringoma of the face and shoulder Arch Dermat & Syph 21:279 1930

- Wendlberger J. Beitrag zur Histogenese des Syringoms Arch f Dermat u Syph 176 467 1938

### Cylindroma

- Balog P Beitrag zur Frage des Schweißdrüsennaevus Dermat Wechschr  
82 841 1926

- Coenen H Das Hidradenoma cylindromatosum der Kopfschwarte Beitr z  
 Klin Chir 95 205 1914 1915

- Davies J H T A contribution to the histology of the cylindromatous tumors of the scalp Brit J Dermat 40 241 1923

- Lausecker H. Beitrag zu den Naevopigmentomen. Arch f Dermat u Syph  
194 639 1952

- Lever W F Pathogenesis of benign tumors of cutaneous appendages and of basal cell epithelioma Arch Dermat & Syph 57 679 1948

- Luger A Das Cylindrom der Haut und seine maligne Degeneration Arch f  
Dermat u Syph 188 155 1919

- Savatard L.  
Schlammadin

- Arch f Des a a Weber K Beitrag zur Kenntniss der Spieglerischen  
Schuermann H and Weber K Beitrag zur Kenntniss der Spieglerischen

- Tumoren (Cylindrome) nebst einigen Bemerkungen zum Epitheliom der  
noides cysticum Arch f Dermat u Syph 175 682 1937

- Lattes R and Kessler R W Metastasizing basal cell epithelioma of skin  
Cancer 4866 1951
- Lennox B and Wells A L Differentiation in the rodent ulcer group of  
tumours Brit J Cancer 5 195 1951
- Lever W F Pathogenesis of benign tumors of cutaneous appendages and of  
basal cell epithelioma Arch Dermat & Syph 57 679 1948
- Mallory F B Recent progress in the microscopic anatomy and differentiation  
of cancer JAMA 55 1513 1910
- Montgomery H Basal squamous cell epithelioma Arch Dermat & Syph 18 50  
1908
- Superficial epitheliomatosis Arch Dermat & Syph 20 339 1929
- in discussion of Goldman H J Multiple benign cystic epithelioma  
JAMA 115 2253 1940
- Nasbet T W Multiple basal cell epitheliomas originating from congenital pig  
pig
- Non

- Pautrier L M Epitheliomas naeviques multiples de la face et du tronc Ann de  
dermat et de syph 75 1917
- Lierard J and Dupont A Nodular epithelioma Brit J Dermat 60 50 1948
- Pinkus H Premalignant fibroepithelial tumors of the skin Arch Dermat &  
Syph 67 598 1953
- Sims C F and Parker R L Intraepidermal basal cell epithelioma Arch  
Dermat & Syph 59 45 1949
- Teloh H A and Wheelock M C Histogenesis of basal cell carcinoma Arch  
Path 48 447 1949
- Wallace S A and Halpert B Trichoma tumor of hair anlage Arch Path  
30 199 1930
- Walther H F Untersuchungen über Krebsmetastasen Schweiz med Wchnschr  
71 998 1941
- Welton D G Elliott J A and Kimmelsiel P Epithelioma Arch Dermat &  
Syph 60 277 1949
- Wise F Multiple superficial benign epithelioma of the skin Arch Dermat &  
Syph 19 1 1909

#### Basal Cell Papilloma and Dermatitis Papulosa Nigra

- Becker S W Seborrheic keratosis and verruca with special reference to the  
melanotic variety Arch Dermat & Syph 63 338 1951
- Eller J J and Ryan V J Senile keratoses and seborrheic keratoses Arch  
Dermat & Syph 22 1013 1930
- Hookey J A Keratoma senile and verruca senilis Arch Dermat & Syph  
23 916 1931
- Michael J C and Seale E R Dermatitis papulosa nigra Arch Dermat &  
Syph 20 609 1929
- Montgomery H and Driffel J Verruca senilis und Keratoma senile Arch f  
Dermat u Syph 166 286 1932
- Pinkus H Ueber einen Fall von Basaliomenepitheliom auf Verruca senilis  
Arch f Dermat u Syph 169 58 1933 1934

#### Carcinoma of Sebaceous Glands

- Beach A and Severance A O Sebaceous gland carcinoma Ann Surg 115 238  
1912

- Highman B and Ogden G E Calcified epithelioma Arch Path 37 169 1944  
 King L S Mummified epidermal cysts Am J Path 23 29 1947  
 Lever W F and Griesemer R D Calcifying epithelioma of Malherbe Arch Dermat & Syph 59 506 1949  
 Malherbe A and Chenavants J Note sur l'épithéliome calcifié des glandes sébacées Bull Soc anat de Paris 5 169 1880  
 Muehlton W F Zur Kenntnis des Malherbeschen Epitheliom Schweiz Ztschr f Path u Bakt 5 53 1912  
 Nicholson G W The formation of bone in a calcified epithelioma of the skin with some remarks on metaplasia J Path & Bact 21 287 1916 1917  
 Sutton R I and Sutton R L Jr Calcifying epithelioma Arch Dermat & Syph 31 18 1935  
 Turhan B and Krüner I Bemerkungen über die sogenannten verkalenden Epitheliome der Haut und ihre Genese Dermatolog 85 73 1942

### Basal Cell Epithelioma

- Adams H G On the nature of rodent ulcer Its relationship to epithelioma adenoides cysticum of Brooke and to other tricho epitheliomata of benign nevoid character its distinction from malignant carcinoma Lancet 1839 1911  
 Amersbach J C Metastatic basal cell epithelioma Arch Dermat & Syph 56 172 1917  
 Anderson N P Bowen's precancerous dermatosis and multiple benign superficial epithelioma evidence of arsenic as an etiologic agent Arch Dermat & Syph 26 1052 1932  
 ——— and Anderson H E Development of basal cell epithelioma as a consequence of radiodermatitis Arch Dermat & Syph 63 586 1951  
 Becker S W Pigmented epitheliomas Arch Dermat & Syph 27 981 1933  
 Bloch B Ueber benigne nicht nevoides Melanoepitheliome der Haut nebst Bemerkungen über das Wesen und die Genese der Dendritenzellen Arch f Dermat u Syph 153 20 1927  
 Ciro M R and Howell J B Morphic like epithelioma Arch Dermat & Syph 63 53 1951  
 Darier J and Ferrand M L'épithéliome pavimenteux mixte et intermédiaire Ann de dermat et de syph 3 385 1922  
 Eller J J and Anderson N P Basal cell epitheliomas with excessive pigment formation Arch Dermat & Syph 27 277 1933  
 Foot N C Adnexal carcinoma of the skin Am J Path 23 1 1947 (Good discussion of histogenesis of basal cell epithelioma)  
 Geschickter C F and Koehler H P Ectodermal tumors of the skin Am J Cancer 23 804 1935  
 Haythorn S R Studies on the histogenesis of the so called basal cell carcinoma Am J Cancer 15 1969 1931  
 Hueck W Zur Morphologie der epithelialen Tumoren insbesondere der Basiliome Virchows Arch 314 137 1947  
 Juon M Ueber die metatypischen Formen der Hautepitheliome Arch f Dermat u Syph 157 81 1929  
 Krantz W Ueber drüsenartige Bilder bei Basalzellenkrebs Dermat Wchnschr 73 1297 1921  
 Krompecher E Der Basalzellenkrebs Jena G Fischer 1903

## Mesodermal Tumors

The mesodermal tumors occurring in the skin may be divided into (1) tumors of fibrous tissue (2) tumors of mucoid tissue (3) tumors of fatty tissue (4) tumors of nerve tissue, (5) tumors of vascular tissue (6) tumors of muscular tissue and (7) tumors of osseous tissue. Inclusion of tumors of nerve tissue under mesodermal tumors is of course not entirely correct but appears justified because mesodermal nerve sheath cells seem to represent an important component of most of the cutaneous tumors of nerve tissue.

Benign and malignant tumors occur. The malignant tumors of mesodermal tissue are called sarcomas. They are almost invariably single tumors. Involvement of other organs takes place by way of metastasis.

Old classifications formerly recognized spindle cell sarcoma and round cell sarcoma. The term spindle-cell sarcoma is synonymous with fibrosarcoma. Round cell sarcomas are no longer classified as such but as lymphomas (i.e. stem-cell reticulum-cell lymphoblastic and lymphocytic lymphoma; see page 471). Lymphomas differ from sarcomas by their potentially systemic nature which is demonstrated by the fact that frequently they arise in multiple foci. Also the term melanosarcoma is no longer in use since the mother cell of the tumors formerly so designated is not a mesodermal cell. The preferred term is malignant melanoma.

### I TUMORS OF FIBROUS TISSUE

#### DERMATOFIBROMA HISTIOCYTOMA (SCLEROSING HEMANGIOMA NODULAR SUBEPIDERMAL FIBROSIS)

This lesion occurs in the skin as firm indolent single or multiple nodules. The nodules usually arise in adults and are situated most commonly on the extremities but occasionally elsewhere. Although they are as a rule not larger than a few millimeters in diameter they may measure several centimeters in size. Most lesions have a reddish color; others are yellowish brown (because of the presence of large amounts of lipid) or bluish black (because of the presence

- Hagedoorn A    Adenocarcinoma of a meibomian gland Arch Ophth 12 850 1934
- Savittard L    Nævus sebaceus and sebaceous carcinoma Brit J Dermat 53 214 1911
- Warren S and Warren N W    Tumors of sebaceous glands Am J Path 19 411 1913

#### Carcinoma of Sweat Glands and Adenocanthoma of Sweat Glands

- Borelli D    Aspetti pseudoglandolari nell'epitelioma discheratosico Adenocanthoma of sweat glands di Lever Dermatolog 97 193 1948
- Gates O Warren S and Warren N W    Tumors of the sweat glands Am J Path 19 591 1913
- Lever W F    Adenocanthoma of sweat glands Arch Dermat & Syph 56 157 1917
- Loos H O    Die Carcinome der Anhangsgebilde der Haut Arch f Dermat u Syph 171 465 1936

#### Metastatic Carcinoma of the Skin

- Edelstein J M    Pancreatic carcinoma with unusual metastasis to the skin and subcutaneous tissue simulating cellulitis New England J Med 242 779 1950
- Freeman C D and Lynch F W    Carcinoma of the breast with peculiar cutaneous metastases Arch Dermat & Syph 35 643 1937
- Gates O    Cutaneous metastases of malignant disease Am J Cancer 30 718 1937 (Good review)
- Levell U W Jr and Tillotson F W    Metastatic cutaneous carcinoma from the breast Arch Dermat & Syph 64 774 1951
- McDonald J H Heckel N J and Kretschmer H I    Cutaneous metastases secondary to carcinoma of urinary bladder Arch Dermat & Syph 61 276 1950
- Murrell T W and Pepple A W    Tumors of the testicle with dermatologic sequelae Arch Dermat & Syph 57 930 1918
- Pfihler G E and Case E A    Erysipelas carcinomatousum resembling radio-dermatitis Am J Roentgenol 35 801 1936
- Ronchese F    Metastases of the scalp simulating turban tumors Arch Dermat & Syph 41 639 1940
- Savittard L    Cancer en cuirasse Brit J Dermat 53 31 1913
- Taylor G W and Meltzer A    Inflammatory carcinoma of the breast Am J Cancer 33 33 1938
- Urbach E Waldow I and Stamm C J    Diffuse cutaneous metastatic lesions from an ovarian carcinoma Arch Dermat & Syph 43 962 1911

the rete ridges. In some instances, this proliferation results in budding formations resembling superficial basal-cell epithelioma. The subcutaneous fat is not invaded (Rentiers and Montgomery). Evidence of phagocytic activity is lacking and staining for iron and lipid gives negative results.



116-115 Dermatofibroma. Low magnification. The collagen shows disorderly arrangement. Much of the collagen is young and instead of staining bright red with hematoxylin-eosin stains pale blue and instead of being assembled in firm bundles, lies in individual fibers. A moderate number of fibroblasts and of capillaries are present ( $\times 50$ ).

**HISTIOCYTOMA** In this lesion, in addition to fibroblasts, there are varying numbers of histiocytes. They tend to lie in necks. Occasionally the great majority of cells are histiocytes. The histiocytes are larger cells than the fibroblasts and possess ample amounts of pale cytoplasm and ovoid rather than round nuclei. Numerous newly formed capillaries may be present. The

amounts of lipid or



of large amounts of hemosiderin) In the latter case, the clinical appearance resembles that of a malignant melanoma

**Histopathology.** Histologically, the nodules can be divided into two types those composed entirely of fibroblasts (dermatofibroma) and those containing, in addition to fibroblasts, varying amounts of histiocytes (histiocytoma)

Various theories exist concerning the histogenesis of this lesion Originally described as dermatofibroma, Woringer, in 1932, proved the presence of phagocytic histiocytes in many dermatofibromas and proposed the term histiocytoma for those containing histiocytes He regarded histiocytomas as young dermatofibromas because of his belief that the histiocyte represents the parent cell of the fibroblast and may develop into a fibroblast (see page 33) Woringer's observations were supported by the studies of Senechal and Caro who showed by vital staining with colloidal iron that some lesions which had histologically the appearance of dermatofibromas contained cells with phagocytic properties and thus were histiocytomas rather than dermatofibromas Gross and Wolbach, on the other hand, have emphasized the presence of blood vessels in these lesions and have expressed their belief that they represent sclerosing hemangiomas They regard the cells present in these lesions as proliferating endothelial cells which attempt to form new blood vessels but do not always succeed and instead become engulfed by regressive fibrosis It would seem however that the development of collagen in these tumors does not represent a regressive process but rather a progressive differentiation A still different view has been taken by Michelson and by Runtz and Montgomery These authors believe that histiocytomas are not tumors at all but represent a chronic inflammatory proliferation of fibroblasts that may follow trauma They refer therefore to dermatofibroma and histiocytoma as nodular subepidermal fibrosis The presence of histiocytes is explained by them as a response to hemorrhage and local tissue destruction

**DERMATOFIBROMA** The nodule contains varying numbers of cells all of which are fibroblasts with spindle shaped nuclei Much of the collagen produced by these fibroblasts is young and instead of staining bright red with hematoxylin eosin it stains a pale blue and instead of being assembled in firm bundles it lies in individual fibers (see Plate 3) The fibroblasts and the collagen are arranged in parallel bands (Fig 213) The lesion is encapsulated, and nearly always is separated from the overlying epidermis by a narrow band of normal collagen The epidermis may be normal or atrophic but commonly shows acanthosis with irregular downward proliferation of

the rete ridges. In some instances this proliferation results in budding formations resembling superficial basal cell epithelioma. The subcutaneous fat is not invaded (Rentiers and Montgomery). Evidence of phagocytic activity is lacking and staining for iron and lipid gives negative results.



collagen shows disorderly arrangement. Much of the collagen is young and instead of staining bright red with hematoxylin is stained in a number of places.

#### HISTIOCYTOMA In situ

varying numbers of histiocytes. The histiocytes are usually the great majority. They are larger cells than the fibroblasts and possess ample amounts of pale cytoplasm and ovoid rather than spindle shaped nuclei (Fig. 214). Numerous newly formed capillaries with prominent endothelial cells may be present. The histiocytes contain varying amounts of lipid or

hemosiderin or both. These substances can often be recognized without special staining but are much better visualized when special stains for iron and lipid are employed. The lipid usually is doubly refractile. Some tumors contain a fairly large number of true foam cells and even Touton giant cells. Occasionally, one may find large foreign body giant cells which have developed from either histio-




FIG. 214 Histiocytoma. The lesion is composed mainly of histiocytes. There are numerous large and small blood vessels lined by prominent endothelial cells. Several foam cells are present, two particularly large foam cells (F.C.) can be seen in the upper left corner. (X400)

cytes or endothelial cells (see below under Nevus xantho endothelialis, page 407).

**Differential Diagnosis** In rare instances, dermatofibromas show a considerable number of nuclei. In such cases differentiation from fibrosarcoma, especially dermatofibrosarcoma protuberans (see page 410) may cause difficulties. However, the circumscribed nature of the lesion, especially the absence of invasion into the subcutaneous fat, the lack of atypicality in the appearance of the nuclei and the absence of mitotic figures rule out sarcoma (Michelson).

Histiocytoma with many fat cells closely resembles xanthoma tuberosum in a regressive fibrosing stage and may even be indistinguishable from it (Arnold and Tilden). In such cases clinical data, such as the number and the location of the lesions and the

presence or the absence of hypercholesterolemia are necessary to arrive at the correct diagnosis

### NEVO XANTHO-ENDOTHELIOMA

Nevo xantho endothelioma is characterized clinically by the presence of a group or groups of yellowish brown nodules arising in early




FIG. 915 Nevo-xantho-endothelioma. The lesion is composed mainly of histiocytes. Several foam cells (FC) and foreign body giant cells (G.C.) are present. This lesion probably represents a form of histiocytoma. (X400)

childhood usually on the extremities. The lesions involute spontaneously within a few years.

**Histopathology.** Nevo xantho-endothelioma probably represents a young histiocytoma. As in histiocytoma, histiocytes predominate; many histiocytes have a pale, vacuolated cytoplasm and, on staining for fat, are shown to contain lipid. Typical foam cells and Touton giant cells are present. There are many capillaries showing proliferation of their endothelium. In addition, large foreign body giant cells

are present which originate either from histiocytes or from endothelial cells (Fig. 215)

Senear and Caro and Montgomery and Osterberg regard nevo xantho endothelioma as a variant of xanthoma. However, since in patients with nevo xantho endothelioma the clinical and the blood chemical findings do not support a diagnosis of xanthomatosis, and since the histologic picture does not differ essentially from that of histiocytoma, it seems best to regard the lesion as a histiocytoma. Nevo xantho endothelioma differs from histiocytoma only by showing foam cells and giant cells of the foreign body type in larger number.

#### RETICULO HISTIOCYTOMA

This rare condition occurs as solitary or multiple, fairly large cutaneous nodules in adults. The lesions may remain stationary or may involute.

**Histopathology.** The histologic picture is characterized by the presence of large, bizarre shaped, multinucleated giant cells with abundant, pale cytoplasm and vesicular nuclei. They are separated from one another by fibrous connective tissue (Fig. 216). Occasionally, it is possible to demonstrate the presence of fat within the cytoplasm of these cells. Vital staining with colloidal iron indicates that the large cells have phagocytic properties and are reticulo endothelial in nature (Caro and Senear).

It is likely that reticulo histiocytoma is an unusual type of histiocytoma. Caro and Senear assume that it is not a true neoplasm but actually a granuloma.

#### KELOID

Keloids represent a post traumatic tissue proliferation. They are red raised and firm and have a smooth, shiny surface. Occasionally clawlike projections radiate from the edges of the lesions.

**Histopathology.** Keloids cannot be differentiated from a dermoid fibroma on a histologic basis. In an early keloid one finds, in addition to thick, intertwining bundles of collagen, a moderate number of fibroblasts (Fig. 216). An old keloid may show but few cellular elements.

#### FIBROSARCOMA

Fibrosarcoma occurs in the skin in two forms, as true fibrosarcoma and as dermatofibrosarcoma protuberans. However, there is justifiable doubt that the latter represents a real sarcoma.

#### TRUE FIBROSARCOMA

True fibrosarcoma starts only rarely in the dermis, more commonly, it starts in the subcutaneous fat (Broders, Hargrave and

Meyerding) The tumor usually feels firm and irregular on palpation and at first is covered by normal skin. It usually grows quite rapidly and as it increases in size the overlying skin at first shows purplish discoloration and finally ulceration. Satellite lesions frequently develop. Metastases occur sooner or later. They usually spread by way of the blood stream, especially to the lungs. The regional lymph nodes are involved only rarely (Genetele).




FIG. 216. Keloid. The lesion is composed of thick, intertwining bundles of collagen. The collagen is entirely mature and stains eosinophilic throughout. A moderate number of nuclei are present; therefore, this keloid is of relatively recent genesis. ( $\times 200$ )

**Histopathology.** This tumor is very cellular. The nuclei vary greatly in size, shape, and staining qualities. Although most nuclei are spindle shaped, others appear round or oval. Some nuclei stain very dark, while others appear vesicular with irregular chromatin structure. Typical as well as atypical mitotic figures are present, often in large numbers. The nuclei tend to lie in bundles that extend in various directions but do not form whorls. In some areas the nuclei lie in dense clusters. Thus, fibrosarcomas present an extremely disordered arrangement of the nuclei. Usually these tumors contain some collagenous tissue in which the nuclei appear to be embedded.

but highly malignant fibrosarcomas may possess little or no collagen. Instead, on staining with a reticulum stain, they are seen to contain numerous reticulum fibers (Foot)

Because of the invasive nature of these tumors, ulceration at the surface and infiltration into the subcutaneous tissue and underlying structures frequently are observed.

**Differential Diagnosis.** A highly malignant fibrosarcoma with no production of collagen may be difficult to differentiate from squamous cell carcinoma, Grade IV, amelanotic malignant melanoma, stem cell lymphoma or reticulum cell lymphoma. On thorough inspection, however, one will usually find, in squamous-cell carcinoma Grade IV, some tendency to keratinization and connections of the tumor with the epidermis (page 331), in malignant melanoma, one will find "junction activity" at the epidermo dermal border (page 458), while in stem cell and reticulum cell lymphoma spindle shaped nuclear forms are absent, since all cells are 'round cells' (see pages 474, 475)

Spindle celled malignant tumors may form in areas of radiodermitis. Although they have the morphologic appearance of a fibrosarcoma, it is likely that most, if not all, represent Grade IV spindle celled squamous cell carcinoma (see page 333) (Gentele, Blom Ides)

#### DERMATOFIBROSARCOMA PROTUBERANS

Dermatofibrosarcoma protuberans represents a slowly growing tumor which has its origin in the dermis. It begins with one or several closely set, small, hard nodules which are reddish or bluish. As the nodules coalesce, they form one or several plaques. On these plaques protruding tumors subsequently develop and may ulcerate.

Although this tumor is locally invasive as a rule, it does not give rise to metastases. In exceptional cases, metastases may occur many years after the appearance of the tumor. Binkley, for instance, reported a case in which metastases caused the death of the patient 38 years after the tumor had first appeared. Coster has expressed the belief that dermatofibrosarcoma protuberans was not truly malignant because of its slow rate of growth and the absence of metastases in most cases. He regards the tumor as intermediary between dermatofibroma and true fibrosarcoma partaking of the nature of both. He prefers to call it progressive recurrent dermatofibroma.

**Histopathology.** The histologic appearance of the tumor does not differ materially from that of true fibrosarcoma (Binkley). A differentiation of the two tumors on a histologic basis therefore is not always possible. Just as in true fibrosarcoma the nuclei lie in irregu-

degeneration of the collagen is observed commonly in parts of the tumor (Binkley). Ulceration at the surface and invasion into the subcutaneous fat occur.




Fig. 11. *Dermatofibrosarcoma protuberans*. The nuclei of the fibroblasts are arranged in irregular strands and whorls. They show a slight degree of atypicity. In contrast with malignant fibrosarcoma, formation of collagen is well in evidence. ( $\times 200$ )

**Differential Diagnosis** For differentiation from dermatofibroma, see page 406.

## 2 TUMORS OF MUCOID TISSUE

### MYXOMA

Although fibromas and neurofibromas may undergo more or less complete mucoid degeneration and then present in such degenerated areas the aspect of myxoma, myxomas may arise as such.

Clinically myxomas present themselves as fairly well circumscribed rather soft, usually intracutaneous tumors over which the epidermis is normal.



but highly malignant fibrosarcomas may possess little or no collagen. Instead on staining with a reticulum stain they are seen to contain numerous reticulum fibers (Foot)

Because of the invasive nature of these tumors ulceration at the surface and infiltration into the subcutaneous tissue and underlying structures frequently are observed.

**Differential Diagnosis** A highly malignant fibrosarcoma with no production of collagen may be difficult to differentiate from squamous cell carcinoma Grade IV, amelanotic malignant melanoma, sarcoma cell lymphoma or reticulum cell lymphoma. On thorough inspection however one will usually find in squamous cell carcinoma Grade IV some tendency to keratinization and connections of the tumor with the epidermis (page 331) in amelanotic melanoma one will find junction activity at the epidermo-dermal border (page 458) while in sarcoma cell and reticulum cell lymphoma spindle shaped nuclear forms are absent since all cells are round cells (see pages 171-175).

Spindle celled malignant tumors may form in areas of radiodermatitis. Although they have the morphologic appearance of a fibrosarcoma it is likely that most if not all represent Grade II spindle celled squamous cell carcinoma (see page 333) (Gentile Blom Ides).

#### DERMATOFIBROSARCOMA PROTUBERANS

Dermatofibrosarcoma protuberans represents a slowly growing tumor which has its origin in the dermis. It begins with one or several closely set small hard nodules which are reddish or bluish. As the nodules coalesce they form one or several plaques. On these plaques protruding tumors subsequently develop and may ulcerate.

Although this tumor is locally invasive as a rule it does not give rise to metastases. In exceptional cases metastases may occur many years after the appearance of the tumor. Binkley for instance reported a case in which metastases caused the death of the patient 38 years after the tumor had first appeared. Costa has expressed the belief that dermatofibrosarcoma protuberans was not truly malignant because of its slow rate of growth and the absence of metastases in most cases. He regards the tumor as intermediary between dermatofibroma and true fibrosarcoma partaking of the nature of both. He prefers to call it progressive recurrent dermatofibroma.

**Histopathology** The histologic appearance of the tumor does not differ materially from that of true fibrosarcoma (Binkley). A differentiation of the two tumors on a histologic basis therefore is not always possible. Just as in true fibrosarcoma the nuclei lie in irregu-

atypicality of the cells favor a diagnosis of myxosarcoma. The cells tend to be stellate and multipolar.

### SYNOVIAL CYST OF THE SKIN (MYXOMATOUS DEGENERATION CYST OF THE SKIN)

This lesion occurs most commonly on the hands particularly near the terminal interphalangeal joints. The lesion consists of a small semiglobular translucent tumor surrounded by erythema. When punctured a clear mucinous fluid exudes.

**Histopathology** Histologic examination reveals a cystic cavity in the dermis or the subcutaneous tissue. The lining wall is made up of fibrous tissue and shows no endothelial or epithelial lining. The cyst is filled with mucoid material.

The histogenesis is not clear. Mackee and Andrews regarded the lesion as a synovial cyst. Savatard held that the lesion formed because of mucoid degeneration of a fibroma. Woodburne stated that he had never seen any evidence of a previously existing solid tumor. He expressed the belief that the lesion arises because of myxomatous degeneration of fibrous tissue of the dermis or of other fibrous tissue.

## 3 TUMORS OF FATTY TISSUE

### LIPOMA

Lipomas occur as single or multiple subcutaneous soft rounded lobulated growths which may or may not be movable against the overlying skin. In rare instances multiple lipomas have been reported as present not only in the subcutaneous but also in the visceral fat depots. This condition is known as systemic multicentric lipoblastosis (Tedeschi).

**Histopathology** Lipomas are composed of fat cells and may or may not be surrounded by a connective tissue capsule. The fat cells in lipomas usually do not differ from normal fat cells. Occasionally they are slightly larger. In some lipomas there is more in others less of a connective tissue framework than in normal subcutaneous fat. Those containing a considerable proportion of connective tissue are called fibrolipomas.

Systemic multicentric lipoblastosis shows not only adult fat cells as lipomas do but also embryonal fat cells and undifferentiated mesenchymal cells with all intermediate stages. These lesions are hamartomas analogous to the lesions of von Recklinghausen's disease and not sarcomas. They show no cellular disorder and no mitotic figures (Tedeschi).

**Histopathology.** Myxomas contain embryonal fibroblasts that possess the ability to produce mucin in a manner similar to that of the embryonal fibroblasts of the umbilical cord. The number of these cells in myxomas is usually small. Those present are spindle shaped or stellate cells possessing multipolar processes. The stroma appears homogeneous and gelatinous. It stains pale blue with hematoxylin



FIG 218 Myxoma. The stroma appears homogeneous and in some places gelatinous. The empty spaces are caused by shrinkage of the mucin in the process of fixation ( $\times 200$ )

and eosin, and red with Best's mucicarmine stain. Because of shrinkage in the process of fixation empty cleftlike spaces frequently are observed in the stroma (Fig 218).

Many myxomas are not pure but contain other mesodermal elements. Thus fibromyxomas and lipomyxomas occur.

### MYXOSARCOMA

Clinically, myxosarcomas are as a rule primarily subcutaneous tumors which, as they grow in size, may cause ulceration of the skin.

**Histopathology.** A decision whether a tumor represents a myxoma or a myxosarcoma is often difficult. A high degree of cellularity and

cytoplasm (Fig 219) There is a loose meshwork of connective tissue which usually but not always is mucoid (Stout)

A highly malignant liposarcoma contains highly atypical lipoblasts with bizarre nuclei The amount of fat in them is often small There are areas which resemble fibrosarcoma The stroma usually is mucoid




Fig 219 Liposarcoma The tumor is composed of

^ ^

### MALIGNANT HIBERNOMA

The clinical appearance of malignant hibernoma is the same as that of other liposarcomas

**Histopathology** The cells comprising this tumor are rounded, have a centrally placed nucleus and contain numerous small and large vacuoles of fat in their cytoplasm Those containing large vacuoles are called mulberry cells. Some of the cells are of very large size and contain multiple bizarre nuclei (Fig 220) The tumor is

Com

## HIBERNOMA

Hibernoma is a rare, benign, solitary, soft, subcutaneous tumor which may slowly increase in size. Clinically, it is indistinguishable from lipoma.

**Histopathology.** The tumor is composed largely of multilocular fat cells which resemble those occurring as brown fat in the hibernating gland of hibernating animals. However, since multilocular fat cells resembling those of brown fat may occur as a transitory phase in the maturation of ordinary, yellow adipose tissue, it is possible that hibernoma merely represents a tumor of embryonal yellow fat cells (Sutherland).

Histologic examination reveals a lobulated tumor composed predominantly of closely spaced, round or oval cells with a distinct cell membrane and centrally placed nucleus. The cells contain either granules or small vacuoles or locules which stain positive with fat stains. This fat is in contrast with mature fat, doubly refractile to polarized light (Sutherland). In some multilocular cells coalescence of smaller into larger locules can be seen, with eccentric displacement of the nucleus. There are, in addition, a few unilocular, normal mature fat cells which are larger than the granular and multilocular cells. There thus seems to be evidence of a transition from granular to multilocular and even to unilocular fat cells (Brines and Johnson).

**Differential Diagnosis.** The tumor can be differentiated from malignant hibernoma by its regular architecture and the absence of mitotic figures and multinucleated cells.

## LIPOSARCOMA

Liposarcomas as a rule arise *de novo* but they may develop within a pre-existing lipoma (Sternberg). They not only occur in the subcutaneous fat but also may be found wherever there is fatty tissue. Those located in the subcutaneous fat present themselves as diffuse nodular infiltrations. *Metastases are common, especially to the lungs and the liver.*

**Histopathology.** The histologic picture varies somewhat with the degree of malignancy. Moderately malignant tumors are easily recognizable as fatty tumors, while very malignant ones may not be identifiable as such unless fat stains are employed.

A moderately malignant liposarcoma consists of adult fat cells and moderately atypical immature fat cells (lipoblasts). The latter have a spindle shaped or stellate nucleus and contain fat droplets in their

appear cut in transverse and others in longitudinal direction (Fig 221) Each nerve bundle is surrounded by fibrous tissue



FIG 221 Neurofibroma Numerous thick bundles of medullated nerves are present Each nerve bundle is surrounded by fibrous tissue ( $\times 900$ )

### NEUROFIBROMATOSIS (VON RECKLINGHAUSEN'S DISEASE)

Neurofibromatosis is characterized by the presence of multiple cutaneous tumors which possess a characteristic soft consistency and usually are flesh-colored but may be brownish or violaceous. The lesions are semiglobular or pedunculated and vary considerably in size. Occasionally large pendulous fleshy masses weighing a pound or more are encountered. In most cases in addition to the tumors the skin shows yellowish brown pigmented macules of various size and shape—so-called café au lait spots.

**Histogenesis** Neurofibromas are tumors of nerve sheaths. Normally each neurite or axon whether myelinated or not, is surrounded by a neuro-ectodermal sheath the schwannian sheath, and

Stout regards this as evidence that ordinary adipose tissue and brown fat come from the same ancestral lipoblastic cell




FIG. 290 Malignant hibernoma. The fat cells contain bizarre nuclei and are filled with small and large vacuoles of fat. Those containing large vacuoles are called mulberry cells. There is hardly any stroma ( $\times 200$ )

#### 4 TUMORS OF NERVE TISSUE

Three types of nerve tumors may occur in the skin: neuroma, neurofibroma, and neurolemmoma.

##### NEUROMA

Neuromas nearly always occur as single tumors secondary to an injury of a nerve. In rare instances they occur as single (Duemling) or multiple (Ludy) tumors apparently without preceding nerve injury. Neuromas are small, reddish nodules which may or may not be painful.

**Histopathology** In their histologic appearance, neuromas resemble the amputation neuromas, which represent a hyperplasia of nerves and are not tumors. For this reason, Ludy preferred to regard also the nontraumatic neuromas of the skin as neuromatoid hyperplasia.

Neuromas show numerous thick bundles of medullated nerves in the dermis, extending in different directions. Thus, some bundles

It is unfortunate that the schwannian cell and the endoneurial fibroblast cannot be differentiated by present histologic methods. Since the schwannian cell can also produce reticulum fibers and collagen (Masson, Murry and Stout) the presence of these structures in neurofibromas does not decide the issue. Murry and Stout believe that their findings in tissue cultures favor the schwannian origin of



FIG. 223 Neurofibroma (von Recklinghausen's disease) In this tumor mucoid degeneration of the collagen has taken place. This is a not uncommon occurrence in lesions of neurofibroma (X200)

neurofibroma because the mode of outgrowth of schwannian cells from normal nerves and of cells in neurofibroma is very similar.

**Histopathology** Histologic examination of the cutaneous tumors shows them to be well circumscribed but not encapsulated. The tumor masses often extend into the subcutaneous fat. They are composed of wavy fibrils which lie in loose strands and tend to form eddies and whorls. The loose textured wavy arrangement of the fibrils is characteristic of neurofibroma (Fig. 222). With hematoxylin

finds a fairly large number of nuclei which are oval or spindle shaped,



a mesodermal sheath, the endoneurium (see page 22). It is still a matter of controversy whether the cells composing neurofibromas represent neuro ectodermal schwannian cells or mesodermal endoneurial cells. Whereas von Recklinghausen originally regarded the tumors as derived from the connective tissue of nerves, Verocay, in 1910, first suggested that the cells of the tumors were immature cells




FIG 222 Neur . . . . . The tumor is composed of . . . . . together and tend . . . . . pale blue with hematoxylin eosin ( $\times 200$ )

of the sheath of Schwann and thus neuro ectodermal cells. He suggested that the tumors be called neurinomas. Among more recent writers on the subject, some (Murray and Stout) subscribe to Verocay's view. Others (Penfield, Tarlov) still adhere to the view that the tumors develop from the perineural connective tissue. Many recent authors however, feel that probably both ectodermal

ery) Masson, who observed that neural elements were common in young tumors but sparse or absent in old tumors concluded that neurofibromas basically were neural tumors. However, with aging, the neural elements gradually degenerated and were replaced by connective tissue proliferation.

nervous systems and of the bones, however, it is a not uncommon occurrence. Hosoi, in 1931, found 65 instances of malignant degeneration reported in the literature; he stated that this represented 13 per cent of all the reported cases of neurofibromas. The conclusion drawn by Charache that "sarcomatous transformation is present in 13 per cent of all cases of multiple neurofibroma" is not justified, however, because in neurofibromatosis, as in other common diseases,

fibroma. In some, however, the pattern of a neurofibroma is preserved: there is still a wavy arrangement of the collagen fibrils, but the nuclei are increased in number and are atypical (Fig. 224). Nerve fibers are never found (Stout, Wachstein and Wolf).

### NEUROLEMOMA

Neurolemoma occurs as either solitary or multiple tumors but not as a systematized disease like neurofibromatosis. The lesions which are asymptomatic are found in the dermis or in the subcutaneous tissue. They can reach a diameter of several centimeters and undergo




FIG. 225. Neurolemoma. Numerous elongated nuclei are arranged in a streaming fashion. In the center of the field lies a so-called Verocay body formed by a double palisade of nuclei enclosing a space nearly devoid of nuclei (X200).

uniform in size and rather pale staining. ~~E~~lastic fibers are absent in the tumors. On staining with Foot's stain numerous wavy reticulum fibers are seen. In most neurofibromas a few nonmedullated thin long nerve fibers can be recognized on staining with special nerve stains, such as Bodian's stain (McNairy and Montgomery).

Not infrequently mucoid degeneration of the collagen is observed in parts of a tumor or in an entire tumor. In such cases the nuclei




FIG. 224 Neurofibrosarcoma (von Recklinghausen's disease). The wavy pattern characteristic of neurofibroma is preserved but the nuclei are increased in number and are atypical ( $\times 200$ ).

are embedded in a homogeneous pale blue ground substance (Fig. 223). One must be familiar with this change because it results in a histologic picture quite different from that usually associated with neurofibroma.

The pigmented macular lesions seen in neurofibromatosis, the so-called café au lait spots, merely show increased melanin pigmentation in the basal layer.

**Systemic Lesions.** There may be tumors of the central and peripheral nervous systems of the viscera (Grill and Kuzma) and of the bones. Involvement of a bone may lead either to extensive destruction or to hypertrophy of bony tissue (Westcott and Ackerman).

**Malignant Degeneration.** Malignant degeneration of neurofibromas of the skin is rare. In tumors of the central and the peripheral

Nevus ~~flammeus~~ is characterized by one or several dull red or bluish red patches of irregular outline, not elevated above the level of the skin.

Nevus ~~vasculosus~~ is a raised, bright red, soft, often lobulated tumor. When lobulated, its appearance resembles that of a strawberry.

~~Cavernous~~ hemangioma consists of a large, soft, subcutaneous mass. If, as is often the case, the lesion is located on the face, consid-



lated capillaries with considerable proliferation of their endothelial cells (X100)

erable deformity may result. The overlying skin may be normal.

as shown (Fig 22b)

NEVUS VASCULOSUS, which represents a capillary hemangioma, in addition to the normal capillaries, consists of numerous dilated capillaries with considerable proliferation of their endothelial cells.

around capillary lumina. In addition, they form solid strands and masses with little evidence of vascular lumina.

CAVERNOUS HEMANGIOMA shows in the lower dermis and in the subcutaneous tissue large, irregular spaces filled with blood. They

partial cystic degeneration. Malignant degeneration does not occur, however.

**Histopathology.** The tumor arises from the sheath of Schwann of a peripheral nerve and is composed of schwannian cells. It is well encapsulated. One observes numerous bands of closely spaced, elongated nuclei arranged in a twisting, streaming fashion. Here and there, the nuclei lie in two rows or palisades enclosing a space nearly devoid of nuclei. Such formations are called Verocay bodies (Fig 225). In addition to these areas of streaming and palisading there are areas in which the schwannian cells are embedded in a loose mesh work of fine connective tissue fibers. Small cysts may be present which by coalescence may form gross cystic spaces (Stout). No nerve fibers are found in neurolemioma. Mast cells often are present in conspicuous numbers.

## 5. TUMORS OF VASCULAR TISSUE

### HEMANGIOMA

Hemangiomas may be divided into three clinical types (1) nevus flammeus (port wine nevus), (2) nevus vasculosus (strawberry mark) and (3) angioma cavernosum.



FIG 226 Hemangioma (nevus flammeus). The capillaries are increased in number. They are dilated, engorged with blood and lined by only a single layer of endothelial cells ( $\times 100$ ).

Nevus flammeus is characterized by the presence of bluish red patches of irregular outline, sometimes above the eye of the skin.

Nevus vasculosus is a plasma truncus and well often lobulated tumor. When lobulated, its appearance resembles that of a strawberry.

Cavernous hemangioma consists of a large and irregular mass. If as is often the case, the lesion is located in the face, it may be called strawberry nevus.



FIG. 227 Hemangioma (nevus vasculosus). There are numerous dilated capillaries with considerable proliferation of their endothelial cells. (X100)

able deformity may result. The overgrowth of the

11

NEVUS VASCULOSUS which represents a capillary hemangioma.

When they form solid strands and

CAVERNOUS HEMANGIOMA shows in the lower dermis and in the subcutaneous tissue large, irregular spaces filled with blood. They

are lined by a single layer of thin endothelial cells and by thick walls (Fig. 228). The thickening is produced mainly by overgrowth of adventitial cells

✓ Sclerosing hemangioma, since it is a dermatofibroma rather than an angioma, is discussed under "Dermatofibroma" (see page 403)



FIG. 228 Hemangioma (hemangioma cavernosum). The blood vessels show considerable thickening of their walls produced by overgrowth of adventitial cells ( $\times 100$ )

### GRANULOMA PYOGENICUM

This lesion, which usually is single, consists of a dull red, soft or fleshy, raised, more or less pedunculated nodule. Its size varies from 0.5 to 2 cm. in diameter. The surface may show a smooth, atrophic epidermis but often is covered by crusts. The lesion bleeds easily when traumatized.

**Histopathology.** On histologic examination, one finds a circumscribed, raised, pedunculated lesion covered by a flattened epidermis and containing numerous newly formed capillaries showing varying

degrees of dilatation (Fig 229). A slight to moderate amount of endothelial proliferation is usually present. The capillaries are embedded in a loose, edematous occasionally mucoid connective tissue. At the neck of the pedunculated lesion the epidermis usually shows acanthotic inward growth thus forming a so called epidermal collar.



Granuloma pyogenicum. The lesion is pedunculated. It is composed of numerous capillaries embedded in a loose edematous stroma. No inflammatory reaction is present. ( $\times 50$ )

In early lesions one finds ~~no~~ inflammatory reaction (Freund). In older lesions because of the fact that the thinned epidermis usually erodes secondary inflammatory changes are often present in the stroma and may give the tumor a granulomatous appearance.

There is no evidence of inflammation. It is caused by a tumor factor. As has been pointed out that, even when the lesion shows a marked granulomatous reaction the center of the lesion retains the same appearance.

Differential diagnosis which a



of the raised pedunculated growth of the lesion, the presence of edema or mucoid degeneration in the stroma, the thinning of the epidermis over the tumor and the collarette formation of the epidermis at the neck of the tumor

### ANGIOKERATOMA MIBELLI, ANGIOKERATOMA CORPORA DIFFUSUM

In angiokeratoma Mibelli, one observes on the dorsum of the fingers, on the toes and on the nees from pinhead to bean sized dark-red, vascular papules with a verrucous surface. The disease arises in young subjects.

In angiokeratoma corporis diffusum, dark red, vascular papules are present in great number over the entire trunk. They are smaller and show less hyperkeratosis than those of angiokeratoma Mibelli. Angiokeratoma corporis diffusum may be associated with internal lesions such as swelling and vacuolation of the muscle fibers of the blood vessels and of the heart (Ruiter).

**Histopathology.** Histologic examination of the cutaneous lesions reveals in both diseases melanectasias of superficial location associated with changes in the epidermis consisting of hyperkeratosis, acanthosis, irregular proliferation of the rete ridges and papillomatosis. Greatly dilated capillaries lined by a thin layer of endothelial cells lie in the papillae and are partly or completely surrounded by the hypertrophic stratum malpighii. If completely surrounded, the dilated capillaries have the appearance of intraepidermal "blood cysts". Some of the blood cysts may have lost their endothelial lining. Atrophy of the stratum malpighii may occur directly over blood cysts because of the pressure which the cysts exert on the overlying epidermis.

### SENILE HEMANGIOMA

Senile hemangiomas are small, raised soft, dark red nodules, measuring usually between 1 and 3 mm in diameter. They may be present in large numbers in persons past middle life. Their sites of predilection are the face and the upper trunk.

**Histopathology.** Senile hemangiomas, within a circumscribed area of the upper dermis show numerous dilated capillaries lined by a single layer of endothelium. The epidermis shows flattening of the rete ridges (Beck). Occasionally however they may show acanthosis as well as hyperkeratosis and even inclusion of dilated capillaries into the epidermis. In that case histologic distinction from angiokeratoma may be impossible (Traub and Tolmach).

## NEVUS ARANEUS

Nevus araneus or spider nevus is characterized by a central slightly elevated red dot from which fine blood vessels radiate. Occasionally pulsation can be observed. Spider nevi are common on the upper half of the face.

**Histopathology** According to Walsh and Becker, nevus araneus represents a small arteriovenous anastomosis. They concluded from serial sections that in this lesion an arteriole ascends high into the dermis where it changes directly into a vein. The vein divides into a network of smaller venules. The latter give the lesion its spiderlike appearance clinically.

However, others differ in their composition of vascular arrangements. The central vessel of the spider nevus is to be an artery which branched successively into arterioles and capillaries. In the other type, the so-called glomus type, they found the afferent artery to be connected with the central vessel of the spider nevus by a short junction which had the histologic appearance of the Sucquet-Hoyer canal as seen in the cutaneous glomus. In contrast with the usual glomus, however, the central vessel of the spider nevus was not a collecting vein but continued into capillaries.

## OSLER'S DISEASE (FAMILIAL HEMORRHAGIC TELANGIECTASIA)

This familial disease is characterized by the presence of numerous telangiectases on the skin and the mucous membranes. The presence of telangiectases on the mucous membranes may result in hemorrhages from the nose, the mouth, the stomach, the kidney, the rectum or the vagina.

**Histopathology** Scattered greatly dilated capillaries are present in the upper dermis. Fingerland and Janousek noted that the venules in the lower dermis showed narrowing of their lumina and an increase in the number of adventitial cells.

## LYMPHANGIOMA

A superficial and a deep variety of lymphangioma exist. The superficial variety, lymphangioma circumscriptum, is characterized by the presence of groups of small, thick-walled vesicles resembling frog spawn. Some of the vesicles may show a verrucous surface.

The deep variety, lymphangioma cavernosum, causes diffuse en-

of the raised pedunculated growth of the lesion, the presence of edema or mucoid degeneration in the stroma, the thinning of the epidermis over the tumor and the collarette formation of the epidermis at the neck of the tumor

### ANGIOKERATOMA MIBELLI, ANGIOKERATOMA CORPORIS DIFFUSUM

In angiokeratoma Mibelli, one observes on the dorsum of the fingers, on the toes and on the knees from pinhead to bean sized dark red, vascular papules with a verrucous surface. The disease arises in young subjects.

In angiokeratoma corporis diffusum, dark red, vascular papules are present in great number over the entire trunk. They are smaller and show less hyperkeratosis than those of angiokeratoma Mibelli. Angiokeratoma corporis diffusum may be associated with internal lesions such as swelling and vacuolation of the muscle fibers of the blood vessels and of the heart (Ruiter).

**Histopathology.** Histologic examination of the cutaneous lesions reveals in both diseases telangiectasias of superficial location associated with changes in the epidermis consisting of hyperkeratosis, acanthosis, irregular proliferation of the rete ridges and papillomatosis. Greatly dilated capillaries lined by a thin layer of endothelial cells lie in the papillae and are partly or completely surrounded by the hypertrophic stratum malpighii. If completely surrounded by the dilated capillaries have the appearance of intra epidermal blood cysts. Some of the blood cysts may have lost their endothelial lining. Atrophy of the stratum malpighii may occur directly over blood cysts because of the pressure which the cysts exert on the overlying epidermis.

### SENILE HEMANGIOMA

Senile hemangiomas are small raised soft, dark red nodules, measuring usually between 1 and 3 mm in diameter. They may be present in large numbers in persons past middle life. Their sites of predilection are the face and the upper trunk.

**Histopathology.** Senile hemangiomas within a circumscribed area of the upper dermis show numerous dilated capillaries lined by a single layer of endothelium. The epidermis shows flattening of the rete ridges (Beck). Occasionally however they may show acanthosis as well as hyperkeratosis and even inclusion of dilated capillaries into the epidermis. In that case histologic distinction from angiokeratoma may be impossible (Traub and Tolmach).

## GLOMUS TUMOR

This tumor usually occurs as a single small, deep seated pink or purplish nodule which is tender and gives rise to severe paroxysmal pains. In rare instances there are numerous lesions in which case tenderness is absent in most of them (Weidman and Wise, Eyster and Montgomery). The most common sites of the solitary lesions are




FIG. 921. Glomus.

a. a venous vessel ( $\times 200$ )

the nail bed and the fingertips; however, the lesions may occur elsewhere.

**Histopathology.** The glomus tumor represents a benign tumor of the cutaneous glomus, a structure composed of an arterial segment, the Sucquet Hoyer canal and a venous segment. The normal Sucquet Hoyer canal possesses a narrow lumen lined with a single layer of endothelial cells and a thick mantle of glomus cells (see page 24). Glomus cells have a faintly eosinophilic cytoplasm and

pa-

ci

fit

A glomus tumor is composed of vascular lumina and numerous

largement of the affected region for instance microcheilia and macroglossia

**Histopathology** In lymphangioma *circumscriptum* one observes in the uppermost portion of the dermis cystically dilated lymph vessels lined by a single layer of endothelium (Fig 230) They contain



FIG 230 Lymphangioma circumscriptum Cystically dilated lymph vessels lined by a single layer of endothelial cells are present in the upper dermis The epidermis shows downward growth and more or less surrounds some of the lymph vessels There is moderate hyperkeratosis ( $\times 50$ )

coagulated lymph and lymphocytes The stratum *malpighii* varies greatly in thickness Over some of the lymph cysts it is thinned else where it may show considerable acanthosis and irregular downward growth Some of the dilated lymph vessels may be completely surrounded by epidermis Hyperkeratosis is common The histologic picture may be similar to that of angiokeratoma except that the dilated areas contain lymph fluid instead of blood

In lymphangioma *externosum* large lymph filled cystic spaces lined by a single layer of endothelium are present in the dermis and the subcutaneous tissue There is concomitant hypertrophy of the connective tissue

## GLOMUS TUMOR

small deep-seated pink or  
paroxysmal  
which case  
tenderness is absent in most of them (Weidman and Wise, Eyster  
and Montgomery) The most common sites of the solitary lesions are



FIG 231 Glomus tumor There are numerous vascular lumina of varying size lined by a single layer of endothelial cells and surrounded by proliferating rows of glomus cells. However some of the masses of glomus cells do not show a central vessel (X200)

the nail bed and the fingertips however the lesions may occur elsewhere

**Histopathology** The glomus tumor represents a benign tumor of the cutaneous glomus a structure composed of an arterial segment the Sucquet Hoyer canal and a venous segment. The normal Sucquet Hoyer canal possesses a narrow lumen lined with a single layer of endothelial cells and a thick mantle of glomus cells (see page 24). Glomus cells have a faintly eosinophilic cytoplasm and large, oval pale nuclei with a distinct chromatin structure. Thus they resemble epithelioid cells. They are richly supplied with nonmyelinated nerve fibrils.

A glomus tumor is composed of vascular lumina and numerous

glomus cells. As a rule the lumina are small but sometimes especially in those cases with multiple lesions, the lumina are of considerable size (Weidman and Wise, Eyster and Montgomery). The vascular lumina are lined by a single layer of flattened endothelial cells and usually, by several layers of glomus cells (Fig 231). Some of the vessels closely resemble the Sucquet Hoyer canals of the normal glomus. In many areas the glomus cells proliferate irregularly from the vascular walls into the connective tissue stroma of the tumor. In addition masses of glomus cells without central lumina and scattered glomus cells are present in the stroma.

The connective tissue stroma of the tumor is loose, edematous and contains scattered fibroblasts and glomus cells. It may show marked mucoid degeneration. Special staining for nerve fibers (Bodian stain) will reveal numerous nerve fibers, most of them nonmyelinated. They show considerable branching and terminate as fine fibrils around the glomus cells.

### HEMANGIOPERICYTOMA

This rare tumor may arise wherever there are capillaries. Its most common sites are the skin and the subcutaneous tissue. Hemangio-




FIG. 232 Hemangiopericytoma. The capillary lumina are lined with a single layer of endothelial cells and are surrounded by irregularly proliferating closely packed pericytes. Most of the pericytes are spindle shaped ( $\times 400$ ).

pericytomas have no diagnostic clinical appearance. They are of varying size, firm and nodular. Some are benign, some are gradually growing and some even metastasize.

✓ **Histopathology** The tumor is characterized by the presence of endothelial tubes and sprouts surrounded by irregularly proliferating cells (Fisher and Kaufman, 1930).

In some cases the pericytes show frequent mitotic figures and invasion into vascular lumina so that the tumor must be regarded as malignant or at least potentially malignant (Fisher, Kaufman and Mason). In one instance widespread metastases resulting in death have been reported (Forrester and Houston).

In the glomus tumor show a more orderly arrangement and are more uniform in appearance, being round to oval and never spindle shaped.

### KAPOSI'S SARCOMA (MULTIPLE IDIOPATHIC HEMORRHAGIC SARCOMA)

Kaposi's sarcoma consists of an eruption of multiple bluish red or dark brown nodules and plaques. The lesions not infrequently show a verrucous surface. They may undergo ulceration. Spontaneous involution of some of the lesions occurs occasionally. The sites of predilection are the distal portions of the extremities but other areas of the skin may also become involved.

Visceral lesions occur in about 10 per cent of the cases (Tedeschi, Folsom and Carnicelli). The most frequent sites are, in order of frequency, the gastro-intestinal tract, the liver, the lungs and the retroperitoneal and the mesenteric lymph nodes (Dorffel). In rare instances there may be visceral lesions of Kaposi's sarcoma without cutaneous involvement (Tedeschi, Folsom and Carnicelli).

**Histogenesis** The histogenesis of Kaposi's sarcoma is not fully ascertained. A widely accepted view (with which the author agrees) is that Kaposi's sarcoma is a benign angioma arising from embryonal vascular cells and that the lesions are autochthonous in origin rather than metastatic (Gilchrist and Keison, Lang and Hasl).



hofer, Becker and Thrasher, Tedeschi, Folsom and Carnicelli). Dissenting views are held by Pautrier and Diss who regard Kaposi's sarcoma as a neuro Dörffel, who regards it as a disease of the and related to the lymphoma group of disease is a sarcoma proposed originally by Kaposi has been abandoned by most writers



FIG. 233 Kaposi's sarcoma, early granulomatous stage. The capillaries are increased in size and number. Their endothelial cells are large. A diffuse chronic inflammatory infiltrate is present. In the lower central section of the illustration one can see groups of endothelial cells, attempting to form new blood vessels. ( $\times 200$ ).

Nevertheless some authors (Aegerter and Peale, Sachs, Azuly and Convit) still maintain that it represents a sarcoma.

Evidence against the concept of Kaposi's sarcoma as sarcoma and in favor of the autochthonous rather than the metastatic origin of the lesions are (1) the absence of a primary focus that progressively enlarges, (2) the appearance of widely separated lesions in crops, (3) the spontaneous regression of some lesions and (4) the fact that histologic examination may reveal very early stages of development in late appearing lesions. Occasionally, however, a lesion may undergo malignant degeneration and then grow as a true sarcoma and cause metastases.



FIG 234 Kaposi's sarcoma neoplastic stage On the left the neoplasia is fibroblastic on the right it is angiomatous ( $\times 100$ )




FIG 235 Kaposi's sarcoma angiomatous lesion There are numerous vascular lumina Most lumina are lined by only a single layer of endothelial cells but some are surrounded also by perithelial cells The stroma is edematous and contains extravasated erythrocytes ( $\times 200$ )

hofer, Becker and Thatcher, Tedeschi, Tolson and Carnicelli). Dissenting views are held by Pautrier and Diss who regard Kaposi's sarcoma as a neurovascular dysgenesis and by Dorfman who regards it as a disease of the reticulo-endothelial system and related to the lymphoma group of diseases. The idea that the disease is a sarcoma proposed originally by Kaposi has been abandoned by most writers.



FIG. 233 Kaposi's sarcoma early granulomatous stage. The capillaries are increased in size and number. Their endothelial cells are large. A diffuse chronic inflammatory infiltrate is present. In the lower central section of the illustration one can see groups of endothelial cells attempting to form new blood vessels ( $\times 900$ ).

Nevertheless some authors (Aegerter and Peale, Sachs, Azulay and Condit) still maintain that it represents a sarcoma.

Evidence against the concept of Kaposi's sarcoma as sarcoma and in favor of the autochthonous rather than the metastatic origin of the lesions are (1) the absence of a primary focus that progressively enlarges, (2) the appearance of widely separated lesions in crops, (3) the spontaneous regression of some lesions and (4) the fact that histologic examination may reveal very early stages of development in late appearing lesions. Occasionally, however, a lesion may undergo malignant degeneration and then grow as a true sarcoma and cause metastases.

20 reticulum represents young collagen) In old regressing lesions one may find considerable collagenization due to maturation of the young fibroblasts This may lead eventually to fibrosis cicatrization and disappearance of the lesion

If malignant degeneration occurs in a lesion of Kaposi's sarcoma the resulting sarcoma is indistinguishable from fibrosarcoma

Seven instances are recorded in the literature in which lymphoma occurred in patients with Kaposi's sarcoma In four of these patients lymphocytic lymphoma was present (Cole and Crump Hufnagel and Dupont Sachs and Gray Fischer and Cohen) in two mycosis fungoides (Lane and Greenwood Winer) and in one Hodgkin's disease (Greenstein and Consten) There is not sufficient proof that Kaposi's sarcoma and lymphoma are related It is therefore, best to regard their coexistence as accidental

### HEMANGIO ENDOTHELIOMA (HEMANGIOSARCOMA)

Malignant tumors arising from blood vessels are rare According to the two types of cells of which capillaries are composed perithelial cells and endothelial cells two types occur hemangio fibrosarcoma and hemangio endothelioma

Hemangio fibrosarcomas have the histologic appearance of fibrosarcomas with conspicuous capillary proliferation and may therefore be regarded as fibrosarcomas The sarcomas which occasionally arise in lesions of Kaposi's sarcoma are of that type This leaves hemangio endothelioma as the only specific malignant tumor arising from blood vessels

Hemangio endothelioma usually occurs as a diffusely infiltrating mass which is soft dark red and raised above the surface of the skin It grows slowly but progressively and may attain large size Metastases occur

Hist. is characterized (1) by the pr greater numbers than required to line the vessels with a simple endothelial membrane and (2) by the new formation of vascular tubes with a marked tendency for their lumina to anastomose (Stout)

The number of vascular lumina varies in different tumors but as a rule vascular lumina are numerous They are irregular in size and shape Large tortuous sinuses may be present (Fig 236) The vascular channels are lined by large, atypical endothelial cells In many are is endothelial cells lie in several layers and proliferate into the lumina to such a degree that the vascular tubes are completely obscured and cannot be made out as such when routine stains are used When however a reticulum stain is used the outline of the vessels

**Histopathology.** The two types of cells of which capillaries are composed participate in the formation of the lesions: endothelial and perithelial cells. The perithelial cell is a pericapillary histiocyte which may develop into a fibroblast (see page 33). In addition, an inflammatory reaction is observed in early lesions. One may divide the lesions of Kaposi's sarcoma into early, granulomatous lesions and late neoplastic lesions. The latter may be either 'angiomatous', if endothelial cell proliferation predominates, or 'fibroblastic' if perithelial cell proliferation predominates.

In early granulomatous lesions, the blood vessels of the dermis are dilated and increased in number (Fig 233). Their endothelial cells are swollen. There is perivascular as well as diffuse cellular infiltration, varying in severity. The infiltrate is composed of lymphocytes, plasma cells and some histiocytes and fibroblasts. One may see groups of endothelial cells attempting to form new blood vessels. Frequently, one sees small groups of extravasated erythrocytes and deposits of hemosiderin. The histologic picture in the early stage is not always diagnostic, however, the presence of extravasated erythrocytes and of hemosiderin in a granulomatous lesion as described above should always make one think of early Kaposi's sarcoma.

In late lesions, the histologic picture may be either angiomatous or fibroblastic, frequently both phases are found intermingled in the same lesion (Fig 234). In angiomatous lesions one finds numerous vascular lumina. They vary greatly in size and occasionally are saccular. Most lumina show only a single layer of swollen endothelial cells, but some are surrounded, in addition, by perithelial cells (Fig 235). The stroma in which the vessels are embedded often is edematous and usually contains hemorrhages and hemosiderin deposits.

In fibroblastic lesions, one observes marked proliferation of spindle shaped cells which represent young fibroblasts (Symmers) and are derived from perithelial cells. They lie in strands which extend irregularly in all directions. The nuclei vary in size and staining qualities and some of them are atypical. Mitotic figures are present though usually in small number. The histologic picture is thus very much like that of a fibrosarcoma. One feature, however, distinguishes the fibroblastic lesions of Kaposi's sarcoma from fibrosarcoma and that is the presence of small or large groups of extravasated erythrocytes and of granules of hemosiderin between the fibroblasts. This feature sometimes is not conspicuous but a thorough search or staining for iron almost invariably will reveal such areas (see Plate 3). The fibroblasts, being immature, form only little collagen as a rule but staining with Foot's stain will reveal a rather dense network of reticulum fibers produced by them (Symmers). (As discussed on page

can be seen. In some tumors, one may see endothelial cells extend as invading cords between the fibers of the connective tissue and, occasionally, they may occur in large, solid sheets.

The endothelial cells seen in this tumor are large and polyhedral and have a well-defined cellular membrane and relatively clear cytoplasm (Fig. 237). The nuclei tend to be round to oval, pale and vesicular. However, many nuclei are atypical, being irregular in shape and hyperchromatic. In addition, numerous mitotic figures are usually present. Some of the endothelial cells are multinucleated.

## POST MASTECTOMY LYMPHANGIO SARCOMA

Lymphangio sarcoma may occur in post mastectomy lymphedema (Stewart and Treves). Several years after radical mastectomy for carcinoma, subcutaneous and cutaneous nodules appear in the edematous tissue of the arm on the side of the operation. The cutaneous nodules have a bluish color. The clinical resemblance to Kaposi's sarcoma may be great. The nodules are not radiosensitive. Metastases, especially to the lungs, occur.

**Histopathology.** The nodules are composed of large, atypical cells which, in some places form solid proliferations and, in others, produce capillary vessels and lacunar structures. The lumens are usually empty, but, in some cases, contain occasional accumulations of red blood cells. In such cases, the tumor represents a mixed lymphangio and hemangio-endothelioma (Jesner Zilk and Rein). Outside the tumor formations, one observes numerous dilated, proliferating lymphatics in the dermis in the subcutaneous tissue and even deeper, in the intermuscular fascia.

**Differential Diagnosis.** The resemblance to hemangio endothelioma is so great that differentiation is impossible except perhaps on the basis of the amount of red blood cells in the lumina. The tumor differs from Kaposi's sarcoma by the intralymphatic proliferation and greater atypicality of the endothelial cells and by the absence of atypical fibroblastic formations.

## 6 TUMORS OF MUSCULAR TISSUE

### LEIOMYOMA

There are three types of leiomyoma of the skin: (1) multiple cutaneous leiomyomas, (2) solitary angioleiomyomas and (3) solitary genital leiomyomas (myomes dartoïques).

Multiple cutaneous leiomyomas are from pinhead- to pea sized, brown or bluish, firm elevated nodules which tend to occur either




FIG. 236 Hemangioendothelioma. Low magnification. There are numerous vascular lumina; on the right side is a large vascular sinus. The vascular channels are lined by large atypical endothelial cells which in some areas proliferate irregularly into the lumen ( $\times 100$ ).




FIG. 237 Hemangioendothelioma. High magnification of Figure 236. A large sinus is shown into which atypical endothelial cells proliferate. Some of the endothelial cells are multinucleated ( $\times 200$ ).

muscle of veins and genital leiomyomas from either the muscularis sexualis or the muscularis mamillae (Stout)

All leiomyomas are composed of interlacing bundles of smooth muscle fibers (Fig 239) Proliferating collagen bundles are often intermingled with the smooth muscle bundles The muscle fibers composing the bundles are straight or slightly wavy and contain centrally located thin very long blunt edged nuclei The muscle bun



collagen is present within a leiomyoma The muscle bundles of the vein merge with those of the tumor ( $\times 200$ )

... in which blue stain may be used With the ...

... new blood vessels

Solitary angioleiomyomas lie largely in the subcutaneous tissue, are encapsulated and contain large blood vessels, probably veins, with thick muscular walls (Fig 239) Some of these vessels have a stellate lumen because of contraction of the muscular tissue In some areas, one sees the muscle bundles of the vessels merge with those of the tumor (Jansen)

Solitary

encaps

rather la



on the back the face or the extensor surfaces of the extremities and usually are arranged in groups They often are painful and sensitive to pressure



FIG 238 Leiomyoma nonvascular type The tumor is composed of interlacing bundles of smooth muscle fibers The nuclei of the smooth muscle fibers are thin long and blunt edged Collagen bundles are intermingled with the smooth muscle bundles Both stain alike with hematoxylin eosin To differentiate them an aniline blue stain may be used (See Plate 4) ( $\times 100$ )

Solitary angioleiomyomas usually are subcutaneous in location but are adherent to the overlying skin They rarely grow larger than 1 cm in diameter and usually are nontender

Solitary genital leiomyomas are located either on the scrotum the labia majora or rarely on the nipples They may attain considerable size several centimeters in diameter and are nontender

**Histopathology** Multiple cutaneous leiomyomas arise from the erectores pilorum muscles solitary angioleiomyomas from the smooth

uently, shows active downward proliferation, even with horn pearl formation (Fig. 241) (Bloom and Ginzler). This pseudo epitheliomatous hyperplasia has been mistaken for squamous cell carcinoma in several cases reported in the literature (Eickhoff).



FIG. 241. Pseudo-epitheliomatous hyperplasia. The dermis contains numerous large pale cells as they are typical of this tumor (X200).

In lesions of the tongue one not infrequently sees areas suggesting transitions between the tumor cells and mature muscle fibers (Crane and Tremblay). This observation has been the reason that granular cell myoblastomas were at first generally regarded as tumors of immature striated muscle cells (myoblasts). However, in recent years, several authors have expressed the view that these transitions were only apparent and that granular-cell myoblastomas were composed of either schwannian cells or endoneurial fibroblasts and thus were of neural origin (Fust and Custer, Ashburn and Rodger, Bangle)

## GRANULAR-CELL MYOBLASTOMA

Granular-cell myoblastomas are solitary tumors occurring most commonly in the tongue, the skin and the subcutaneous tissue (Crane and Tremblay). They usually are benign, but malignant degeneration occurs occasionally (Ross, Miller and Foote). Cutaneous myoblastomas usually consist of a firm, round, well-circumscribed, non-




FIG. 240. Granular-cell myoblastoma. The tumor is composed of large cells having a pale cytoplasm filled with coarse granules. ( $\times 400$ )

tender nodule, from 0.5 to 2 cm. in diameter, which may lie within the thickness of the skin or may be pedunculated. In some instances, the surface of the tumor is hyperkeratotic. Subcutaneous myoblastomas consist of a firm nodule which may or may not be attached to the overlying skin.

**Histopathology.** On histologic examination, the cells of the tumor appear large and polyhedral. Most cells measure from 20 to 60 microns in diameter, but some are even larger. They have a pale cytoplasm filled with coarse acidophilic granules (Fig. 240). The nuclei are small, round or oval and somewhat vesicular (Cipollaro and Einhorn). Some cells contain more than one nucleus. Delicate strands of collagen surround the cells of the tumor.

The overlying epidermis usually is hyperplastic and, not infre-



FIGURE 21<sup>o</sup> Osteoma cutis. Low magnification. The bone appears laminated about several Haversian canals ( $\times 100$ )



FIGURE 22<sup>o</sup> Osteoma cutis. High magnification of Figure 21<sup>o</sup>. The bone lined in many areas by osteoblasts. In the center of the field several osteoclasts lie within a niche called Howship's lacuna ( $\times 400$ )

The reasons these authors have given are first, granular cells occasionally are arranged concentrically around a core consisting of bundles of axis cylinders (Fust and Custer), second groups of granular cells have been found within nerve sheaths both within and in the vicinity of the tumor (Fust and Custer, Ashburn and Rodger) and, third, the granules stain slightly positive for lipids, suggesting that they form as a result of the disintegration of axis cylinders and myelin sheaths (Bangle).

In the rare cases of malignant granular cell myoblastomas, one can observe all stages of transition from typical granular cells to malignant spindle and giant cells devoid of granules. Widespread hematogenous metastases may occur (Ross, Miller and Foote).

**Differential Diagnosis.** On cursory examination, the large, pale cells of the tumor resemble the foam cells of xanthoma. However, the cells of granular cell myoblastoma contain a granular and not a foamy cytoplasm and take fat stains only very faintly. Furthermore, in granular cell myoblastoma, the overlying epidermis tends to be hyperplastic rather than atrophic as in xanthoma.

## 7. TUMORS OF OSSEOUS TISSUE

### OSTEOMA CUTIS

Cutaneous bone formation may be primary (heterotopic) or secondary (metaplastic). Only lesions with primary bone formation should be called osteomas.

In primary bone formation bone develops in areas which were not the site of previous lesions. The bone probably develops from embryonal rests. Such lesions therefore represent nevus tumors or hamartomas (Hopkins, Dietrich, Vero, Machacek and Bartlett).

In secondary bone formation, bone develops in areas of tissue degeneration. Bone may develop in tumors particularly in the calcifying epithelioma of Malherbe (see page 368), in scar tissue (Lisga and Burns) in lesions of scleroderma in various granulomas and in areas of fat necrosis or hemorrhage. Also multiple small foci of ossification may occur in the skin of the face in prolonged, severe acne with scarring (Leider). In all these instances the bone develops by metaplasia and frequently, though not always calcification precedes the ossification.

Osteomas may be single (Dietrich) or multiple (Hopkins, Vero, Machacek and Bartlett, Tjstens and Rueter). Their clinical appearance is not uniform. Usually they occur as small hard plaques or nodules within the dermis or subcutis.



**Histopathology** On histologic examination, one or several pieces of bone are observed within a lesion. The bone appears lamellated in concentric rings about several Haversian canals. These canals contain blood vessels and connective tissue (Fig 212). Most of the bone lamellae contain numerous small lacunae each filled with a cell. These cells called osteocytes possess various shapes, some of them are stellate. Along the margin of the bone one sees many osteoblasts which build bone and a few osteoclasts which absorb bone. The osteoblasts have small oval or elongated nuclei. As they lay down bone substance they become enclosed in the bone as osteocytes. The osteoclasts have multiple large nuclei and resemble multinucleated foreign body giant cells. Frequently they lie within deep grooves called Howship's lacunae which extend into the bone substance (Fig 213).

The tissue surrounding the piece or pieces of bone often is highly vascular and cellular and may contain fat cells so that it resembles bone marrow.

Occasionally one may find in osteomas not only bone but also fibrocartilage (Vero Marcheck and Bartlett).

#### BIBLIOGRAPHY

##### Dermatofibroma Histiocytoma Nevo xantho endothelioma and Reticulo histiocytoma

- Arnold H L Jr and Tilden I L. Histiocytoma cutis a variant of xanthoma. *Arch Dermat & Syph* 47:198 1943
- Caro M R and Seneff F E. Reticulohistiocytoma of the skin. *Arch Dermat & Syph* 65:701 1952
- Gross R E and Wolbach S B. Sclerosing hemangiomas. *Am J Path* 19:533 1943
- Michelson H E. Nodular subepidermal fibrosis. *Arch Dermat & Syph* 27:81 1933
- Montgomery H and Osterberg A E. Xanthomatosis. *Arch Dermat & Syph* 37:373 1938
- Rentiers P L and Montgomery H. Nodular subepidermal fibrosis (dermatofibroma versus histiocytoma). *Arch Dermat & Syph* 59:568 1949 (Good review)
- Seneff F E and Caro M R. Histiocytoma cutis. *Arch Dermat & Syph* 33:209 1936
- Nevo xantho endothelioma or juvenile xanthoma. *Arch Dermat & Syph* 34:195 1936
- Truab E F and Monash S. Dermatofibroma. *Arch Dermat & Syph* 26:250 1932
- Woringer F and Kwiatkowski S I. L'histiocytome de la peau. *Ann de dermat et syph* 3:998 1932
- Zak F G. Reticulohistiocytoma (ganglioneuroma) of the skin. *Brit J Dermat* 62:351 1950

## Keloid

- Garb J and Stone M J Keloids *Am J Surg* 58 315 1942

## Fibrosarcoma

- Binkley G W Dermatofibrosarcoma protuberans *Arch Dermat & Syph* 40 578 1939  
 — — — — — 1950  
 of soft  
 Costa O G Progressive recurrent dermatofibroma (Darier Ferrand) *Arch. Dermat & Syph* 54 432 1946  
 Foot N C Pathology in Surgery p 60 Philadelphia Lippincott 1945  
 Cencile H Malignant fibroblastic tumors of the skin *Acta dermat venereol* 31 suppl 27 1951 (Good review)

## Myxoma and Myxosarcoma

- Foot N C Pathology in Surgery p 77 Philadelphia Lippincott 1945

## Synovial Cyst of the Skin

- Mackee G M and Andrews G C The pathologic histology of synovial lesions of the skin *Arch Dermat & Syph* 5 361 1922  
 Savatard L Periarticular fibroma of the skin (synovial lesions of the skin) *Arch Dermat & Syph* 9 441 1924  
 Woodburne A R Myxomatous degeneration cysts of skin and subcutaneous tissues *Arch Dermat & Syph* 56 407 1947

## Lipoma Hibernoma and Liposarcoma

- Brines O A and Johnson M H Hibernoma a special fatty tumor *Am J Path* 25 467 1919  
 Sternberg S S Liposarcoma arising within a subcutaneous lipoma *Cancer* 5 975 1952  
 Stout A P Liposarcoma—the malignant tumor of lipoblasts, *Ann Surg* 119 86 1944  
 Sutherland J C Callahan W P and Campbell C L Hibernoma a tumor of brown fat *Cancer* 5 364 1952  
 Fedeschi C G Systemic multicentric lipoblastosis *Arch Path* 42 320 1946

## Neuroma

- Duening W M Cutaneous neuroma *Arch Dermat & Syph* 19 226 1929  
 Ludy J B Cutaneous neuroma *Arch Dermat & Syph* 21 419 1930

## Neurofibromatosis

- Charache H Multiple neurofibroma with sarcomatous transformation and skeletal involvement *Arch Dermat & Syph* 40 185 1959  
 Foot N C Histology of tumors of the peripheral nerves *Arch Path* 30 772 1940  
 Crill J and Kurma J F Recklinghausen's disease with unusual symptoms from inter- and intra- — — — — —  
 Howell K — — — — —  
 crence — — — — —



**Histopathology** On histologic examination one or several pieces of bone are observed within a lesion. The bone appears lamellated in concentric rings about several Haversian canals. These canals contain blood vessels and connective tissue (Fig 242). Most of the bone lamellae contain numerous small lacunae each filled with a cell. These cells called osteocytes possess various shapes, some of them are stellate. Along the margin of the bone one sees many osteoblasts which build bone and a few osteoclasts which absorb bone. The osteoblasts have small oval or elongated nuclei. As they lay down bone substance they become enclosed in the bone as osteocytes. The osteoclasts have multiple huge nuclei and resemble multinucleated foreign body giant cells. Frequently they lie within deep grooves called Howship's lacunae which extend into the bone substance (Fig 243).

The tissue surrounding the piece or pieces of bone often is highly vascular and cellular and may contain fat cells so that it resembles bone marrow.



Occasionally one may find in osteomas not only bone but also fibrocartilage (Vero Machacek and Bartlett).

#### BIBLIOGRAPHY

##### Dermatofibroma Histiocytoma Nevo xantho endothelioma and Reticulohistiocytoma

- Arnold H L Jr and Tilden J L. Histiocytoma cutis a variant of xanthoma. *Arch Dermat & Syph* 47:498 1943
- Cirio M R and Senear F E. Reticulohistiocytoma of the skin. *Arch Dermat & Syph* 65:701 1952
- Gross R E and Wolbach S B. Sclerosing hemangiomas. *Am J Path* 19:533 1913
- Michelson H E. Nodular subepidermal fibrosis. *Arch Dermat & Syph* 27:81<sup>o</sup> 1933
- Montgomery H and Osterberg A E. Xanthomatosis. *Arch Dermat & Syph* 37:373 1938
- Rentiers P L and Montgomery H. Nodular subepidermal fibrosis (dermatofibroma versus histiocytoma). *Arch Dermat & Syph* 59:668 1949 (Good review)
- Senear F E and Cirio M R. Histiocytoma cutis. *Arch Dermat & Syph* 33:209 1936
- Nevo xantho endothelioma et juvenile xanthoma. *Arch Dermat & Syph* 34:190 1936
- Traub E F and Monrad S. Dermatofibroma. *Arch Dermat & Syph* 26:200 1932
- Woringer F and Kwiatkowski S L. L'histiocytome de la peau. *Ann de dermat et syph* 3:998 1932
- Zak F G. Reticulohistiocytoma (ganglioneuroma) of the skin. *Brit J Dermat* 62:351 1950

Leiomyoma Aniline blue (Mallory) stain  
This stain serves to differentiate collagen from smooth muscle. With hematoxylin-eosin both stain red but with aniline blue collagen stains blue and muscle red ( $\times 170$ )



Blue nevus Numerous large, spindle shaped deeply pigmented cells are located in the lower dermis. They are melanoblasts ( $\times 175$ )

- McNairy D J and Montgomery H Cutaneous tumors of von Recklinghausen's disease (neurofibromatosis) Arch Dermat & Syph 51 381 1915 (Good review)
- Masson P Experimental and spontaneous schwannomas (peripheral gliomas) Am J Path 8 367 389 1932
- Murray M R and Stout A P Schwann cell versus fibroblast as the origin of the specific nerve sheath tumor Am J Path 16 41 1910
- Penfield W Tumors of the sheaths of the nervous system Arch Neurol & Psychiat 27 1298 1932
- Stout A P The malignant tumors of the peripheral nerves Am J Cancer 25 1 1935
- Tarlov I M The origin of perineural fibrosarcoma Am J Path 16 33 1910
- Verocay J Zur Kenntnis der Neurofibrome Beitr z path Anat u z allg Path 48 1, 1910
- Wachstein M and Wolf E General neurofibromatosis (von Recklinghausen's disease) with local sarcomatous change and metastasis to regional lymph nodes Arch Path 37 331 1911
- Westcott R J and Ackerman I V Elephantiasis neurofibrosa Arch Dermat & Syph 55 233 1947

#### Neurolemoma

- Stout A P The peripheral manifestations of the specific nerve sheath tumor (neurolemoma) Am J Cancer 21 751 1935
- Neurolemoma in Tumors of the Peripheral Nervous System p 15 Washington D C Armed Forces Institute of Pathology 1949

#### Hemangioma

- Miescher G Ueber plane Angiome (Naevi hyperaemici) Dermatolog 106 176 1953

#### Granuloma Pyogenicum

- Ireund H Zur Kenntnis des telangiectatischen Granuloms Arch f Dermat u Syph 166 669 1932
- Michelson H L Granuloma pyogenicum Arch Dermat & Syph 12 192 1925

#### Angiokeratoma

- Ruster M Angiokeratoma corporis diffusum Arch Dermat & Syph 68 21 1953
- Früh I F and Isenbach J A Angiokeratoma Arch Dermat & Syph 21 39 1931
- Wile U J and Belote G H Angiokeratoma Arch Dermat & Syph 18 501 1928

#### Senile Hemangioma

- Beck C H Zur Histologie der disseminierten papulösen Telangiectasen (Angioma senilis) und der Telangiectasia traneæ Arch f Dermat u Syph 175 481 1937

#### Nevus Araneus

- Patek A J Post J and Victor J C The vascular spider associated with cirrhosis of the liver Am J Med Sc 200 341 1910
- Walsh E N and Becker S W Erythema palmare and nevus araneus like telangiectases Arch Dermat & Syph 41 616 1941







## 448 Mesodermal Tumors

- Mackee G M and Cipollaro A C Multiple idiopathic hemorrhagic sarcoma (Kaposi) *Am J Cancer* 26 1 1936 (Good review)
- Pautrier L M and Diss A Kaposi's idiopathic sarcoma is not a genuine sarcoma but a neurovascular dysgenesis *Brit J Dermat* 41 93 1929
- Sachs W Azulay R D, and Convit J Multiple idiopathic hemorrhagic sarcoma of Kaposi *J Invest Dermat* 8 317 1947
- and Gray M Kaposi's sarcoma and lymphatic leukemia *Arch Dermat & Syph* 51 325 1945
- Symmers D Kaposi's disease *Arch Path* 32 764 1941
- Tedeschi C G Folsom H F and Carnicelli T J Visceral Kaposi's disease *Arch Path* 43 335 1947
- Winer L H Mycosis fungoides *Arch Dermat & Syph* 56 480 1947

### Hemangioendothelioma

- Caro M R and Stubenrauch C H Jr Hemangioendothelioma of the skin *Arch Dermat & Syph* 51 295 1945
- Stout A P Hemangioendothelioma a tumor of blood vessels featuring vascular endothelial cells *Ann Surg* 118 445 1943
- Sweitzer S E and Winer L H Hemangioendothelioma *Arch Dermat & Syph* 34 997 1936

### Post mastectomy Lymphangio sarcoma

- Jessner M Zak F G and Rein C R Angiosarcoma in postmastectomy lymph edema (Stewart Treves syndrome) *Arch Dermat & Syph* 65 123 1952
- Stewart F W and Treves A Lymphangiosarcoma in postmastectomy lymphedema *Cancer* 1 64 1948

### Leiomyoma

- Jansen L H Leiomyomata cutis *Acta dermat venereol* 32 40 1952
- Stout A P Solitary cutaneous and subcutaneous leiomyoma *Am J Cancer* 29 435 1937

### Granular Cell Myoblastoma

- Ashburn L L and Rodger R C Myoblastoma neural origin *Am J Clin Path* 22 440 1952
- Bangle R Jr A morphological and histochemical study of the granular cell myoblastoma *Cancer* 5 950 1952
- Bloom D and Ginzler A M Myoblastoma *Arch Dermat & Syph* 56 618 1947
- Cipollaro A C and Einhorn M B Granular cell myoblastoma Report of two cases *Arch Dermat & Syph* 56 812 1947
- Crane A R and Tremblay R G Myoblastoma (granular cell myoblastoma or myoblastic myoma) *Am J Path* 21 357 1945
- Eickhoff H Myoblastenmyom und Carcinom Virchows *Arch* 304 432 1939
- Fust J A and Custer R P The neurogenesis of so called granular cell myoblastoma *Am J Clin Path* 19 522 1949
- Ross R C Miller T R and Foote F W Malignant granular cell myoblastoma *Cancer* 5 112 1952

### Osteoma Cutis

- Dietrich C Osteomatosis cutis *Arch Dermat & Syph* 41 562 1940
- Hopkins J G Multiple miliary osteomas of the skin *Arch Dermat & Syph* 18 706 1928 (Good review)

- Leider M Osteoma cutis as a result of severe acne vulgaris of long duration  
Arch Dermat & Syph 62:403 1950
- Lalga H V and Burns D C Osteomatosis cutis Arch Dermat & Syph  
46:872 1942
- Tijdens E F and Ruiter M Ueber Osteosis Kulis Acta dermat venercol  
29:140 1949
- Vero F Machacek G F and Bartlett F H Disseminated congenital osteomas  
of the skin J A M A 129:728 1943



## Pigmented Nevi and Malignant Melanoma

Pigmented nevi and malignant melanomas are composed of nevus cells. Lentigo, the Mongolian spot and the blue nevus represent special types of pigmented nevi which will be discussed at the end of this chapter.

### PIGMENTED NEVUS

From a histologic point of view, it is practical to divide pigmented nevi into intradermal (resting) nevi and junction (active) nevi. However, intermediary forms so called compound nevi, are common. It is thus understandable that one cannot always predict from the clinical appearance whether a nevus is of the intradermal type or the junction type.

The intradermal nevus, as a rule, is a circumscribed, smooth or slightly verrucous elevation which may or may not contain a number of hairs and varies in color from normal skin to dark brown. Intradermal nevi are very rare on the palms, the soles and the genitalia. Nevi located in these locations almost always are junction nevi.

The junction nevus is a relatively flat or slightly raised, smooth, pigmented mark that is devoid of hairs. Its color is light brown or brownish black, rarely slate blue or bluish black. The majority of the marks that are slate blue or bluish black are not junction nevi but blue nevi.

**Histogenesis of the Nevus Cell** In former years the nevus cell had been thought to be of epidermal origin. It was believed that nevus cells were modified basal cells that had migrated (dropped off) from the epidermis into the dermis (Abtropfung theory). Today there are but few adherents of this theory (Allen). The great majority of investigators have accepted the theory first proposed by Masson in 1926, that both the melanocytes in the epidermis, the so called clear cells, and the nevus cells are of neural origin (see page 5). Masson stated that nevus cells may develop from two sources: from melanocytes in the epidermis and from schwannian cells of cutaneous nerves. He believed that junction nevi develop exclusively from

melanocytes whereas compound and intradermal nevi develop from both melanocytes and schwannian cells. So long as there was migration of melanocytes from the epidermis into the dermis the nevus was a compound nevus but when this migration ceased it became an intradermal nevus. Thus compound and intradermal nevi according to Masson have a dual origin from two primordia which fused together. This dualistic view of the origin of compound and intra



FIG. 244 Junction nevus. Well circumscribed nevus cell nests are present in the lower epidermis. Otherwise the epidermis appears normal. This type of junction nevus is not apt to become malignant. (X200)

dermal nevus has been questioned by some authors (Lund and Stobbe) who believe that all pigmented nevi develop solely by dropping off of melanocytes.

As evidence in favor of his theory of the neural origin of

trands of schwan  
in the deep por  
roid tubes and  
Additional evidence in favor of Mas  
son's theory is the observation by Berkhiser and Rappoport that  
nevi cells may be found proliferating within the perineural sheath  
of superficial cutaneous nerves

## Pigmented Nevi and Malignant Melanoma

Pigmented nevi and malignant melanomas are composed of nevus cells. Vitigo, the Mongolian spot, and the blue nevus represent special types of pigmented nevi which will be discussed at the end of this chapter.

### PIGMENTED NEVUS

From a histologic point of view, it is practical to divide pigmented nevi into intradermal (resting) nevi and junction (active) nevi. However, intermediary forms so-called compound nevi, are common. It is thus understandable that one cannot always predict from the clinical appearance whether a nevus is of the intradermal type or the junction type.

The intradermal nevus, as a rule, is a circumscribed, smooth or slightly verrucous elevation which may or may not contain a number of hairs and varies in color from normal skin to dark brown. Intradermal nevi are very rare on the palms, the soles and the genitalia. Nevi located in these locations almost always are junction nevi.

The junction nevus is a relatively flat or slightly raised, smooth pigmented mark that is devoid of hairs. Its color is light brown or brownish black, rarely slate blue or bluish black. The majority of the marks that are slate blue or bluish black are not junction nevi but blue nevi.

**Histogenesis of the Nevus Cell** In former years the nevus cell had been thought to be of epidermal origin. It was believed that nevus cells were modified basal cells that had migrated (dropped off)

Today  
: majority  
of investigators have accepted the theory proposed by Masson in 1926, that both the melanocytes in the epidermis, the so-called clear cells, and the nevus cells are of neural origin (see page 5). Masson stated that nevus cells may develop from two sources: from melanocytes in the epidermis and from schwannian cells of cutaneous nerves. He believed that junction nevi develop exclusively from

at inclined homogeneous cytoplasm. The nucleus is large,  
 or show varia  
 the epi  
 they may




FIG. 216 Junction nevus of the premalignant type. Nevus cells lie diffusely scattered in the lower epidermis, which appears disorganized. No actual invasion of the nevus cells into the dermis is present, however. The upper dermis contains a ~~handlike inflammatory infiltrate~~ intermingled with melanophores (It is possible that this type of junction nevus actually represents the earliest phase of a malignant melanoma) ( $\times 200$ ).

**JUNCTION NEVUS** This form of pigmented nevus, well described by Fbert by Traub and Keil and by Allen and Spitz, is characterized by the active formation of nevus cells in the lower epidermis—i.e. at the epidermal-dermal junction. There are two types of junction nevus between which, however, there are transitions. In one type, the newly formed nevus cells are present largely as well-circumscribed

**Histopathology.** In a junction (active) nevus, there is active formation of nevus cells in the basal layer of the epidermis (i.e., at the epidermal dermal junction) and no nevus cell nests are found in the dermis. In a compound nevus, there also is junction activity, but, in addition, well formed nevus cell nests are present in the dermis. In an intradermal nevus junction, activity is no longer present and the




FIG 245 Nevus pigmentosus compound type. The cells composing the intra epidermal and the subepidermal nevus cell nests are spindle shaped and resemble schwannian cells. The nevus cell nests suggest Messner tactile bodies (X100)

nevus cells are all located in the dermis. However, completely intradermal (resting) nevi are rare. If serial sections are made it will be seen that nearly all intradermal nevi show at least a few areas of junction activity.

That the development from junction nevus into a compound nevus and, further, into an intradermal nevus is a matter of aging of a nevus is suggested by the observation of Lund and Stobbe that a much higher percentage of nevi show junctional activity in children than in adults. Only nevi located at the distal portions of the extremities were found consistently to be junction nevi, even in adults.

The typical nevus cell is oval or cuboidal in shape.

suggest Meissner tactile bodies (Fig 245). Although all junction nevi have a certain potentiality to become malignant, because they are active nevi, the danger of malignant degeneration is slight in this type of junction nevus with its regularly formed nests of nevus cells.



FIG. 245. Nevus pigmentosus, intradermal type. In the upper dermis the nevus cells lie in nests and cords. In the lower dermis the nevus cells are arranged more loosely and embedded in fibrous tissue ( $\times 100$ ).

In the second type of junction nevus, the premalignant junction nevus of Allen and Spitz, in which the nevus cells lie diffusely scattered in the lower epidermis, the lower epidermis appears disorganized by the presence of numerous vacuolated nevus cells with irregularly shaped nuclei (Fig 246). The nevus cells usually contain a considerable amount of melanin. The border between the epidermis and the dermis is somewhat irregular, but no actual invasion of

nevus cell nests within the lower epidermis while in the other type the nevus cells are scattered diffusely through the lower epidermis. This latter type of junction nevus has been called premalignant junction nevus by Allen and Spitz.

In junction nevus with well circumscribed nevus cell nests in the lower epidermis the residual epidermis appears essentially normal.




FIG 247 Nevus pigmentosus compound type. In the basal layer of the epidermis are seen clear cells (melanocytes) lying singly and in nests. Those lying in nests have already the appearance of nevus cells. One nest (N) is in the stage of dropping off. In addition typical nevus cell nests (N.C.N.) lie free in the dermis. Considerable amounts of melanin are present in the clear cells and in some of the nevus cell nests ( $\times 900$ ).

(Fig 244) Frequently there are nests which lie beneath the epidermis but still are in contact with it and thus are in the stage of dropping off. Occasionally in other parts of the same nevus one observes nevus cell nests located free in the dermis representing a development into a compound nevus. The nevus cells contain varying amounts of melanin. Melanophores may be present in the dermis but no inflammatory infiltrate is present. The nevus cells comprising the intraepidermal nests have a regular cuboidal appearance. In rare instances however they are spindle shaped and endowed with stroma so that they resemble schwannian cells and the cell nests

suggest Meissner tactile bodies (Fig 245). Although all junction nevi have a certain potentiality to become malignant, because they are active nevi the danger of malignant degeneration is slight in this type of junction nevus with its regularly formed nests of nevus cells



FIG 246 Nevus pigmentosus intradermal type. In the upper dermis the nevus cells lie in nests and cords. In the lower dermis the nevus cells are arranged more loosely and embedded in fibrous tissue ( $\times 100$ )

In the second type of junction nevus the premalignant junction nevus of Allen and Spitz in which the nevus cells lie diffusely scattered in the lower epidermis the lower epidermis appears disorganized by the presence of numerous vacuolated nevus cells with irregularly shaped nuclei (Fig 246). The nevus cells usually contain a considerable amount of melanin. The border between the epidermis and the dermis is somewhat irregular, but no actual invasion of



nevus cell nests within the lower epidermis while in the other type the nevus cells are scattered diffusely through the lower epidermis. This latter type of junction nevus has been called premalignant junction nevus by Allen and Spitz.

In junction nevi with well circumscribed nevus cell nests in the lower epidermis, the residual epidermis appears essentially normal.




FIG 247 Nevus pigmentosus compound type. In the basal layer one sees clear cells (melanocytes) lying singly and in nests. Those lying in nests have already the appearance of nevus cells. One nest (N) is in the stage of dropping off. In addition typical nevus cell nests (N-C-N) lie free in the dermis. Considerable amounts of melanin are present in the clear cells and in some of the nevus cell nests (X200).

(Fig 244) Frequently there are nests which lie beneath the epidermis but still are in contact with it and thus are in the stage of dropping off. Occasionally in other parts of the same nevus one observes nevus cell nests located free in the dermis representing a development into a compound nevus. The nevus cells contain varying amounts of melanin. Melanophores may be present in the dermis but no inflammatory infiltrate is present. The nevus cells comprising the intraepidermal nests have a regular cuboidal appearance. In rare instances however they are spindle shaped and endowed with stroma so that they resemble schwannian cells and the cell nests

fore can be taken as evidence of the benign nature of the nevus in which they occur. They should not be confused with the giant cells occurring in so-called juvenile melanoma and occasionally, in malignant melanoma (see page 462). The nevus cells in the upper portion of the dermis frequently though not always contain melanin. In the presence of much melanin melanophores are often present.




FIG. 250 Verrucous nevus pigmentosus. There is fingerlike downward growth of the epidermis around nests of nevus cells. There are numerous multinucleated nevus cells. (This is not a sign of malignancy.) ( $\times 100$ )

No inflammatory reaction is present except in the case of mechanical irritation or of secondary infection. In that case the inflammatory infiltrate however is focal rather than bandlike or diffuse as it is in malignant melanoma.

In the lower dermis the nevus cells lie farther apart, tend to be spindle shaped and are embedded in fibrous tissue. The fibrous tissue may have the same loose, waxy, pale appearance as in neurofibroma (often). Within this fibrous tissue the nevus cells may be arranged in narrow columns so that they suggest neural sheaths (neural tubes of Masson). In other areas the fibrous tissue may be in concentric arrangement (lamellae foliaceae of Masson) resulting in structures resembling Meissner tactile bodies (Fig. 249). Montgomery and

nevus cells in the dermis is present. The upper dermis contains numerous melanophores and, in many instances, also a bandlike inflammatory infiltrate. This type of junction nevus is often difficult to differentiate from an early malignant melanoma (see page 459) and it is possible that it actually represents the earliest phase of a malignant melanoma. In any event, it possesses a definite potentiality to progress into a malignant melanoma.




FIG. 249 Nevus pigmentosus, intradermal type. The lower dermis is shown. The nevus cells are largely spindle shaped. They are embedded in fibrous tissue which has the same loose, wavy, pale appearance as in neurofibroma. In the center one sees a neuroid structure (N S) resembling a Meissner tactile body. ( $\times 200$ )

**COMPOUND NEVUS** In a compound nevus one observes features of both junction nevus and intradermal nevus. Nevus cell nests are seen within the epidermis, dropping off from the epidermis, as well as in the dermis free from contact with the epidermis (Fig. 247).

**INTRADERMAL NEVUS** Intradermal nevi show, in the upper dermis, nests and cords of nevus cells surrounded by bundles of collagen (Fig. 248). In some instances the nevus cells lie in dense masses. Not infrequently, one encounters multinucleated nevus cells showing either clumping or rosette like arrangement of small, darkly staining nuclei. These giant cells occur only in well matured nevi and, there-

melanomas whereas intradermal nevi, which show little or no junction activity, give rise to a malignant melanoma very rarely. Especially the so-called premalignant junction nevus which is characterized by a diffuse out to the upper



Fig. 251 Margin of an early malignant melanoma. On the left the epidermis is normal. In the center because of the presence of many

... a ... extends on the left beyond the margin of the tumor under the normal epidermis. (X100)

In an older lesion but only dermis ... of numerous atypical nevus cells, appears vacuolated, torn apart and even disintegrated (Fig. 251). This type of early malignant melanoma greatly resembles the premalignant type of junction nevus and differs from it only by the greater atypicality of the cells and the beginning invasion into the dermis (see also Differential Diagnosis). As in the premalignant

Kernohan found such structures in 11 per cent of their material of pigmented nevus. With special nerve stains such as the Bodian stain one frequently can see nerve fibers in the fibrous tissue. They may occasionally be seen in close relation to the limes follicles.

An occasional intradermal nevus is devoid of nevus cell nests in the upper dermis and shows only spindle shaped nevus cells embedded in abundant fibrous tissue. Differentiation from a neurofibroma may then be difficult on a histologic basis (Cohen Becker). Such nevi are referred to as neural nevus.

The epidermis over intradermal nevus may be normal but often is flattened because of pressure from below. In some nevus the epidermis shows hyperkeratosis, papillomatosis and icelike downward growth (verrucous nevus pigmentosus) (Fig. 250). In others large hair follicles are present (nevus pigmentosus et pilosus).

## MALIGNANT MELANOMA

Malignant melanoma may arise as such or may develop from a pigmented nevus. If a malignant melanoma arises from a pigmented nevus such nevus is almost invariably of the junction type. Clinical evidence that a malignant change is occurring in a pigmented nevus is presented by an increase in the size of the lesion, an increase in the depth of pigmentation and not infrequently the development of an inflammatory border with or without spilling of pigment from the lesion into the surrounding skin.

Early malignant melanoma is characterized by a gradually enlarging deeply pigmented nodule usually surrounded by erythema. Later the lesion becomes fungating and satellite lesions may appear. Ulceration is a late symptom. In occasional instances hyperpigmentation is slight or absent.

Metastasis takes place at first through the lymphatics resulting in involvement of the regional lymph nodes. Blood spread is a late event and may be absent until nearly the end. When it occurs metastases are usually widespread. The liver, the lungs, the brain and the skin are the most common sites of hematogenous metastases.

Malignant melanoma is rare before puberty and when it occurs it is usually clinically benign. For instance in a series of 13 histologically malignant melanomas in children published by Spitz only one death occurred.

**Histopathology** The malignant changes invariably begin at the dermal epidermal junction irrespective of whether a malignant melanoma starts as such or develops from a pigmented nevus (Miescher). For this reason pigmented nevus with considerable junction activity like the junction nevus predispose to the development of malignant

melanomas, whereas intradermal nevi, which show little or no junction activity, give rise to a malignant melanoma very rarely. Especially with in the



FIG 251 Margin of an early malignant melanoma. On the left, the epidermis is normal. In the center, because of the presence of many

left beyond the margin of the tumor under the normal epidermis ( $\times 100$ )

In an early malignant melanoma and at the advancing border of older lesions one may find considerable changes within the epidermis but only slight invasion of the dermis by tumor cells. The lower epidermis, because of the presence of numerous atypical nevus cells, appears vacuolated, torn apart and even disintegrated (Fig 251). This type of early malignant melanoma greatly resembles the pre-malignant type of junction nevus and differs from it only by the greater atypicality of the cells and the beginning invasion into the dermis (see also "Differential Diagnosis"). As in the pre-malignant



irregularly shaped, often deeply pigmented nests of atypical nevus cells which may show mitotic figures (Fig. 253). The upper epidermis may be invaded and become so permeated with tumor cells that it




Fig. 253

(x 100)

disintegrates and ulceration results (Fig. 254). In addition, there is deep penetration of the dermis by atypical nevus cells. Simultaneously with the downward penetration of the nevus cells, the epidermis undergoes ulceration of the rete ridges by the downward

The size and the shape of the tumor cells in the dermis show great



type of junction nevus, one finds in the upper dermis, close to the epidermis, a dense, bandlike inflammatory infiltrate intermingled with many melanophores. This inflammatory infiltrate often extends for a distance away from the malignant melanoma under the normal




FIG. 252 Malignant melanoma. Low magnification. There is considerable junction activity. One finds not only dropping off of nevus cells and nevus cell nests downward into the dermis but also migration upward into the stratum malpighii. The tumor cells are largely of the cuboidal type and lie in alveolar formations ( $\times 100$ ).

epidermis (Fig. 251). It is of interest that once deep invasion of the dermis has taken place, the inflammatory infiltrate is absent except at the margins if there is extension of the tumor and in well progressed malignant melanoma.

In an advanced malignant melanoma there is also considerable irregular junction activity (Fig. 252). The epidermis contains large,

in malignant melanoma usually is strongly positive in the cells near the dermo-epidermal junction. The tumor cells deep in the dermis, however, react weakly or not at all (Miescher)

In children malignant melanoma is uncommon and even lesions which histologically have the appearance of malignant melanoma



110-111 Malignant melanoma. The tumor cells are fusiform and lie in irregular strands. The tumor thus resembles a melanophore. (x100)

cause metastases only in the rarest instances. These malignant melanomas generally referred to as juvenile melanomas differ histologically from the adult type.

In about 5% of the cases, the tumor contains giant cells.

These giant cells occur in adult malignant melanoma (Spitz, Haber). These giant cells have a round, oval or stellate

variation. Nevertheless two types of cells can be recognized clearly a cuboidal and a fusiform type. Although most tumors show both cell types almost invariably one type greatly predominates. Predominance of cuboidal cells is much more common than predominance of fusiform cells. The cuboidal cells tend to lie in alveolar formations (Figs 252 and 253). The fusiform cells tend to lie in irregular branching strands (Fig 255). Tumors in which fusiform cells predominate





FIG 254 Malignant melanoma. The epidermis has disintegrated due to permeation with tumor cells (X200)

resemble fibrosarcoma but differ from it by the presence of junction activity. Mitotic figures are usually present in malignant melanoma but often only in small numbers. Their presence is of great value as evidence that the tumor is malignant inasmuch as in pigmented nevi mitotic figures are very rare or absent (Miescher). Bizarre multinucleated giant cells occur occasionally.

The amount of melanin varies considerably in malignant melanomas. In some tumors considerable amounts of melanin are present in both tumor cells and melanophores. In others, particularly in rapidly growing tumors there may be no evidence of melanin in hematoxylin-eosin stains. Such tumors have been referred to as amelanotic malignant melanoma. However staining of sections with silver will reveal in most instances a few cells containing melanin. Irrespective of the amount of melanin present the dopa reaction

in malignant melanoma usually is strongly positive in the cells near the dermo-epidermal junction. The tumor cells deep in the dermis, however, react weakly or not at all (Vriescher)

In children malignant melanoma is uncommon and even lesions which histologically have the appearance of malignant melanoma



irregular strands. The tumor thus resembles a fibrosarcoma but differs from it by the presence of junction activity and of considerable amounts of melanin which are located in tumor cells as well as in melanophores (X100)

cause metastases only in the rarest instances. These malignant melanomas generally referred to as juvenile melanomas differ in their histologic aspects from malignant melanomas of adults.

variation. Nevertheless, two types of cells can be recognized clearly, a cuboidal and a fusiform type. Although most tumors show both cell types, almost invariably one type greatly predominates. Predominance of cuboidal cells is much more common than predominance of fusiform cells. The cuboidal cells tend to lie in alveolar formations (Figs 252 and 253). The fusiform cells tend to lie in irregular branching strands (Fig 255). Tumors in which fusiform cells predominate




Fig 254 Malignant melanoma. The epidermis has disintegrated due to permeation with tumor cells ( $\times 200$ )

resemble fibrosarcoma but differ from it by the presence of junction activity. Mitotic figures are usually present in malignant melanoma but often only in small numbers. They should be searched for because their presence is of great value as evidence that the tumor is malignant inasmuch as in pigmented nevi mitotic figures are very rare or absent (Miescher). Bizarre multinucleated giant cells occur occasionally.

The amount of melanin varies considerably in malignant melanomas. In some tumors considerable amounts of melanin are present in both tumor cells and melanophores. In others, particularly in rapidly growing tumors, there may be no evidence of melanin in hematoxylin-eosin stains. Such tumors have been referred to as amelanotic malignant melanoma. However, staining of sections with silver will reveal in most instances a few cells containing melanin. Irrespective of the amount of melanin present the dopa reaction

the basal layer of the projecting rete ridges shows considerable hyperpigmentation and a great increase in the number of clear cells. No junction activity i.e. dropping-off of clear cells is observed. The upper dermis often contains melano



FIG. 256. Lentigo. The rete ridges are elongated. Numerous clear cells (melanocytes) are present in the basal layer ( $\times 900$ ).

phores and sometimes a mild perivascular lymphocytic infiltrate. In the case of senile lentigo the upper dermis also shows basophilic degeneration of the collagen.

Whether or not lentigo can progress by the onset of dropping-off activity into a junction nevus is not yet fully decided, but this appears likely since junction nevi not infrequently show at the periphery.

(fr hyperpigmentation of the basal layer but no elongation of the rete ridges.

### MONGOLIAN SPOT

The typical Mongolian spot occurs as a bluish gray discoloration in the lumbosacral region. It is found with great regularity in mem

shape and an acidophilic cytoplasm with one or more large, vacuolated, irregularly outlined nuclei. They differ from the giant cells of intradermal nevi, which have small, deeply basophilic nuclei, and from those found occasionally in malignant melanoma, which are larger and more bizarre.

The histologic appearance of metastatic lesions of the skin differs from that of primary lesions by the absence of junction activity. Metastatic lesions never show any inflammatory infiltrate (Dixon).

**Differential Diagnosis.** Differentiation of early malignant melanoma from junction nevus may cause considerable difficulties. That is easily explainable: a junction nevus may develop into a malignant melanoma. If this development proceeds slowly, as it often does, tumors in an intermediate stage of development result. The type of junction nevus most likely to undergo malignant degeneration is the one with diffusely scattered nevus cells in the lower epidermis, the so called premalignant junction nevus (page 455, fig. 246). This type of junction nevus, like early malignant melanoma, has numerous vacuolated nevus cells with atypical nuclei in the lower epidermis which thus appears disorganized. It may show, in addition, a band like inflammatory infiltrate in the upper dermis. It differs from malignant melanoma by the absence of invasion of the nevus cells into the dermis and by the absence of mitotic figures. However, these are not absolutely reliable criteria, and in some instances a decision as to whether malignancy exists already is impossible. There are some authors, like Miescher, who are inclined to regard as malignant any junction nevus showing a bandlike inflammatory infiltrate. According to Miescher, this is evidence that aggressive invasion is being fought off. It matters little, however, whether such junction nevi are called premalignant or "early malignant", it is important that they require wide excision.

A highly malignant amelanotic melanoma may be difficult to differentiate from a highly malignant fibrosarcoma or a Grade IV spindle cell squamous cell carcinoma. For their differentiation see under Fibrosarcoma page 110.

## LENTIGO

Lentigo is a smooth, not infiltrated dark brown mark usually measuring only a few millimeters in diameter. A juvenile and a senile type occur. Juvenile lentigines begin to appear in childhood and occur on all parts of the body. Senile lentigines ("liver spots") occur in old age on the dorsal of the hands, on the forearms and on the face. Both types are benign. Clinical differentiation from junction nevus may be impossible.

**Histopathology** The histologic findings are the same in juvenile and senile lentigo. The rete ridges appear elongated and club shaped (Fig 256). Eccentric thumb like buds may project from them (Zeisler and Becker, Cauley and Curtis). The basal layer of the projecting rete ridges shows considerable hyperpigmentation and a great increase in the number of clear cells. No junction activity, i.e. X dropping-off of clear cells is observed. The upper dermis often contains melano-




FIG. 256. Lentigo. The rete ridges are elongated. Numerous clear cells (melanocytes) are present in the basal layer ( $\times 200$ ).

phores and sometimes a mild perivascular lymphocytic infiltrate. In the case of senile lentigo the upper dermis also shows basophilic degeneration of the collagen.

Whether or not lentigo can progress by the onset of dropping-off activity into a junction nevus is not yet fully decided but this appears likely since junction nevus not infrequently show at their periphery the histologic picture of lentigo.

**Differential Diagnosis** Lentigo must be differentiated from ephelis (freckle). Ephelides show hyperpigmentation of the basal layer but no elongation of the rete ridges.

### MONGOLIAN SPOT

The typical Mongolian spot occurs as a bluish gray discoloration in the lumbosacral region. It is found with great regularity in mem-



bers of the Mongolian race but occurs occasionally also in other races. It is present at birth and gradually fades.

Occasionally, Mongolian spots occur outside the usual location as aberrant Mongolian spots (Cole, Hubler and Lund). They may even be multiple (Carleton and Biggs) or appear during adult life and gradually increase in size (Pariser and Beerman).

**Histopathology.** The dermis shows, especially in its midportion fusiform and stellate cells filled with brown pigment granules. These cells lie widely scattered as well as in small groups. They give a positive dopa reaction (see page 12), indicating that they are melanocytes (Cole, Hubler and Lund).

Formerly, the cells composing the Mongolian spot were regarded as mesodermal melanocytes, as opposed to the epidermal melanocytes forming pigmented nevi. However, the theory of the neural origin of melanocytes has done away with this artificial division into epidermal and mesodermal melanocytes. The cells of the Mongolian spot (as well as those of the blue nevus—see below) are now regarded as melanocytes which in their embryonal migration from the neural crest to the epidermis failed to reach the epidermis but became arrested in the dermis (Montgomery).

## BLUE NEVUS

The blue nevus is a sharply circumscribed, round or oval, soft nodule of slate blue or bluish black color which measures, as a rule, only a few millimeters in diameter. There usually is only one lesion, although multiple blue nevi do occur. In rare instances, large, infiltrated plaques have been encountered (Uppshaw, Ghormley and Montgomery). Malignant degeneration of blue nevus is very rare, but it does occur (Montgomery and Kähler, Allen and Spitz). Metastases may develop and cause death.

**Histopathology.** The cells composing blue nevus are of the same type as those composing the Mongolian spot. They are dopa positive melanocytes which became arrested in the dermis during their embryonal migration from the neural crest to the epidermis (Montgomery). The number of these cells is however much greater in blue nevus than in the Mongolian spot.

In the blue nevus greatly elongated melanocytes lie grouped in irregular masses and bundles in the middle and the lower thirds of the dermis (Fig. 257). Occasionally these cell masses extend into the subcutaneous layer or close to the epidermis. The epidermis, however, is normal. The melanocytes are spindle shaped and have long bipolar, occasionally branching processes (see Plate 4). They lie predominantly with their long axis parallel to the epidermis. Most of

them are filled with numerous fine granules of melanin, often so completely that their nucleus cannot be visualized. In addition, melanophores (chromatophores) frequently occur within and beside the masses of melanocytes. The melanophores differ from the melanocytes by being shorter and thicker and by containing coarser granules (Fig. 258). In addition, the melanocytes are dopa positive while the melanophores are dopa negative.



FIG. 258. Blue nevus. Low magnification. Numerous spindle-shaped melanocytes filled with melanin, lie grouped in irregular bundles in the middle and the lower thirds of the dermis and in the subcutaneous fat (X50).

Blue nevus may undergo fibrosis. In that case, thick bundles of collagen separate the melanocytes. In occasional instances, the presence of a pigmented nevus and of a blue nevus in the same lesion has been reported (Montgomery and Kallier).

The blue color which the blue nevus presents is due to the presence of melanin in the dermis and subcutaneous fat.

Blue nevus may undergo malignant degeneration, irregularly hyperchromatic nuclei, blue nevus in which malignant degeneration has occurred.

be . . . . . occasionally also in other races

It . . . . .

outside the usual location as aberrant Mongolian spots (Cole, Hubler and Lund) They may even be multiple (Carleton and Biggs) or appear during adult life and gradually increase in size (Pariser and Beerman)

**Histopathology.** The dermis shows, especially in its midportion fusiform and stellate cells filled with brown pigment granules These cells lie widely scattered as well as in small groups They give a positive dopa reaction (see page 12), indicating that they are melanocytes (Cole, Hubler and Lund)

Formerly, the cells composing the Mongolian spot were regarded as mesodermal melanocytes, as opposed to the epidermal melanocytes forming pigmented nevi However, the theory of the neural origin of melanocytes has done away with this artificial division into epidermal and mesodermal melanocytes The cells of the Mongolian spot (as well as those of the blue nevus—see below) are now regarded as melanocytes which in their embryonal migration from the neural crest to the epidermis failed to reach the epidermis but became arrested in the dermis (Montgomery)

### BLUE NEVUS

The blue nevus is a sharply circumscribed, round or oval, soft nodule of slate blue or bluish black color which measures, as a rule, only a few millimeters in diameter There usually is only one lesion, although multiple blue nevi do occur In rare instances, large, infiltrated plaques have been encountered (Upshaw, Ghormley and Montgomery) Malignant degeneration of blue nevus is very rare, but it does occur (Montgomery and Kahler, Allen and Spitz) Metastases may develop and cause death

**Histopathology** The cells composing blue nevus are of the same type as those composing the Mongolian spot They are dopa positive melanocytes which became arrested in the dermis during their embryonal migration from the neural crest to the epidermis (Montgomery) The number of these cells is, however, much greater in blue nevus than in the Mongolian spot

In the blue nevus greatly elongated melanocytes lie grouped in irregular masses and bundles in the middle and the lower thirds of the dermis (Fig 257) Occasionally these cell masses extend into the subcutaneous layer or close to the epidermis The epidermis, however, is normal The melanocytes are spindle shaped and have long bipolar, occasionally branching processes (see Plate 4) They lie predominantly with their long axis parallel to the epidermis Most of

them are filled with numerous fine granules of melanin often so completely that their nucleus cannot be visualized. In addition, melanophores (chromatophores) frequently occur within and beside the masses of melanocytes. The melanophores differ from the melano-



FIG. 257 Blue nevus. Low magnification. Numerous spindle shaped melanocytes filled with melanin are grouped in irregular bundles in the middle and the lower thirds of the dermis and in the subcutaneous fat ( $\times 50$ )

Blue nevus may undergo fibrosis. In that case thick bundles of collagen separate the melanocytes. In occasional instances the presence of a pigmented nevus and of a blue nevus in the same lesion has been reported (Montgomery and Kahler).

The blue color which the blue nevus presents clinically results from the deep location of the pigment.

Blue nevus in which malignant degeneration has occurred (large irregularly hyperchromatic nuclei)

curred show areas of necrosis as a diagnostically important feature (Allen) The malignant cells may appear swollen and vacuolated and may lose their spindle character so that recognition of the lesion is




FIG. 258 Blue nevus High magnification of Figure 257 There are numerous spindle shaped melanocytes filled with fine melanin granules In addition melanophores filled with large irregularly shaped melanin granules are present in the middle third of the field shown ( $\times 400$ )

A malignant blue nevus must be based on the presence of residual portions of the original benign blue nevus

## BIBLIOGRAPHY

### Pigmented Nevus

- Allen A C A reorientation on the histogenesis and clinical significance of cutaneous nevus and melanoma *Cancer* 2:28 1949  
 — and Spitz S Malignant melanoma *Cancer* 6:1 1953 (Informative)  
 Becker S W Diagnosis and treatment of pigmented nevus *Arch Dermat & Syph* 60:44 1949 (Good review)  
 Berkhiser S W and Rappoport A E The comparative morphogenesis of the dermo epidermal nevus and malignant melanoma *Am J Path* 28:477 1952  
 Cohen E L Solitary neurofibroma of scalp *Brit J Dermat* 57:172 1915  
 Ebert M H Multiple pigmented nevus *Arch Dermat & Syph* 37:1 1938  
 Lund H Z and Stobbe G D The natural history of the pigmented nevus factors of age and anatomic location *Am J Path* 25:1117 1949

- Masson P. Les naevi pigmentaires tumeurs nerveuses. *Ann d'anat path* 3 417 607 1926
- My conception of cellular nevi. *Cancer* 49 1951 (Informative)
- Montgomery H. and Kethohan J. W. Pigmented nevi with special studies regarding a possible neuro-epithelial origin of the nevus cell. *J Invest Dermat* 3 465 1940
- Traub E. F. and Keil H. The common mole. *Arch Dermat & Syph* 41 214 1940
- Unna P. G. Die Histopathologie der Hautkrankheiten p 1147 Berlin V Hirschfeld 1934

### Malignant Melanoma

- Ackerman I. V. Malignant melanoma of the skin. *Am J Clin Path* 18 602 1949
- Allen A. C. and Spitz S. Malignant melanoma. *Cancer* 6 1 1953
- Dixon H. A. Melanotic sarcoma with extreme melanosis. *Arch Dermat & Syph* 38 571 1938
- Haber H. Cellular naevi in children (juvenile melanoma). *Tr St John's Hosp Dermat Soc* 33 44 1952
- Miescher G. Die Entstehung der bösartigen Melanome der Haut. *Virchows Arch* 261 86 1927 (Good review)
- Spitz S. Melanomas of childhood. *Am J Path* 24 591 1918

### Lentigo

- Cawley E. P. and Curtis A. C. Lentigo senilis. *Arch Dermat & Syph* 62 635 1950
- Zenker E. P. and Becker S. W. Generalized lentigo. *Arch Dermat & Syph* 33 109 1936

### Mongolian Spot

- Carleton A. and Biggs R. Diffuse mesodermal pigmentation with congenital cranial abnormality. *Brit J Dermat* 60 10 1949
- Cole H. N. Jr. Hubler W. R. and Lund H. Z. Persistent aberrant Mongolian spots. *Arch Dermat & Syph* 61 244 1950
- Montgomery H. Benign and malignant dermal neoplasms. *J A M A* 150 1182 1952
- Pariser H. and Beerman H. Extensive blue patchlike pigmentation. *Arch Dermat & Syph* 59 396 1949

### Blue Nevus

- Allen A. C. and Spitz S. Malignant melanoma. *Cancer* 6 1 1953
- Montgomery H. Benign and malignant dermal neoplasms. *J A M A* 150 1182 1952
- and Kahler J. E. The blue nevus (Jadassohn-Tièche) its distinction from ordinary moles and malignant melanomas. *Am J Cancer* 36 527 1939 (Good review)
- Upshaw B. Y. Ghormley R. L. and Montgomery H. Extensive blue nevus of Jadassohn-Tièche. *Surgery* 22 761 1947

## Lymphoma and Myelosis

Lymphoma designates a group of malignant tumors arising, usually in multiple foci, from the lymphoid reticular system. Myelosis is a designation for malignant tumors arising, invariably in multiple foci, from the myeloid system.

### LYMPHOMA

Lymphomas are composed primarily of immature and mature cells of the lymphoid reticular system. The mother cell of the lymphoid reticular system is the lymphoid reticular stem cell. This cell may differentiate into a lymphoid cell (lymphoblast to lymphocyte) or into a reticular cell (reticulum cell to histiocyte to fibroblast) (see Chart 2 page 33). The type of lymphoma which results in a given case depends on the degree of immaturity of the cells and on the direction of their differentiation.

The lymphomas may start as a solitary lesion and may remain as such for a long period of time. As a rule, however, the lesions are multiple from the beginning. The multiplicity of the lesions is due to the systemic nature of the disease and usually not to metastasis. Occasionally, however, dissemination by metastasis occurs in addition. Leukemia is frequent with some types, rare with other types of lymphoma. Leukemia is merely the result of a release of immature tumor cells into the blood stream and does not represent a separate form of disease. Invariably one finds in patients with leukemia involvement of the bone marrow. However, marrow involvement may occasionally be found in patients with normal blood pictures (Gall and Mallory).

Just as leukemia is a part of the disease which may be present or absent, so involvement of the skin is a variable manifestation of lymphoma, found more frequently in some forms of lymphoma than in others, but a possible occurrence in all. In a review of 618 cases of lymphoma, Gall and Mallory found leukemic blood changes in 17 per

groups, the monomorphous and the polymorphous. In the

former the tumors are composed entirely of lymphoid reticular cells while in the latter group the tumors show an admixture of inflammatory cells. The following classification which is based on that by Gall and Mallory is suggested

Monomorphous group

- 1 Stem cell lymphoma
- 2 Reticulum cell lymphoma
- 3 Lymphoblastic lymphoma
- 4 Lymphocytic lymphoma
- 5 Follicular lymphoma (Brill Symmers)

Polymorphous group

- 6 Hodgkin's disease
- 7 Mycosis fungoides

It is impossible to assign every case of lymphoma to one of these seven types because some tumors are in an intermediary stage of differentiation or show features of more than one type. Furthermore different types of lymphoma may be encountered in different areas of the same patient and not infrequently as the disease progresses the lesions may become less differentiated and require reassignment to another type (Heim Gall and Mallory Herbut Miller and Erl).

The terms round-cell sarcoma lymphosarcoma and reticulum-cell sarcoma formerly were and occasionally still are used as designation for single tumors of lymphoma. The use of the term sarcoma is misleading however because even in the case of a single lesion the potentially systemic nature of the disease may become apparent at any time.

Non-specific lesions which when found in association with leukemia are often referred to as leukemids may consist of macules papules purpuric lesions vesicles bullae eczematous lesions and exfoliative dermatitis. Specific lesions may consist of plaques nodules and tumors, in addition however any of the lesions mentioned as occurring as a non-specific reaction may show a specific lymphomatous infiltrate. This holds true especially of eczematous lesions and exfoliative dermatitis. Thus it is impossible to predict always from the clinical appearance whether a lesion is non-specific or specific.

**Histopathology** Histologically specific lesions of lymphoma of the skin show either large masses of lymphoma cells patchy accumulations of lymphoma cells or an inflammatory infiltrate intermixed with lymphoma cells. The lymphoma cells vary with the type of lymphoma and may be immature lymphoid reticular cells such as



# Lymphoma and Myelosis

✓ Lymphoma designates a group of malignant tumors arising usually in multiple foci from the lymphoid reticular system. Myelosis is a designation for malignant tumors arising invariably in multiple foci from the myeloid system.

## LYMPHOMA

Lymphomas are composed primarily of immature and mature cells of the lymphoid reticular system. The mother cell of the lymphoid reticular system is the lymphoid reticular stem cell. This cell may differentiate into a lymphoid cell (lymphoblast to lymphocyte) or into a reticular cell (reticulum cell to histiocyte to fibroblast) (see Chart 2 page 33). The type of lymphoma which results in a given case depends on the degree of immaturity of the cells and on the direction of their differentiation.

The lymphomas may start as a solitary lesion and may remain as such for a long period of time. As a rule however the lesions are multiple from the beginning. The multiplicity of the lesions is due to the systemic nature of the disease and usually not to metastasis. Occasionally however dissemination by metastasis occurs in addition. ✓ Leukemia is frequent with some types rare with other types of lymphoma. Leukemia is merely the result of a release of immature tumor cells into the blood stream and does not represent a separate form of disease. Invariably one finds in patients with leukemia involvement of the bone marrow. However marrow involvement may occasionally be found in patients with normal blood pictures (Gall and Mallory).

Just as leukemia is a part of the disease which may be present or absent so involvement of the skin is a variable manifestation of lymphoma found more frequently in some forms of lymphoma than in others but a possible occurrence in all. In a review of 618 cases of lymphoma Gall and Mallory found leukemic blood changes in 17 per

groups the monomorphous and the polymorphous 60.1%

forms of lymphoma. They occur most commonly in stem-cell lymphoma, reticulum-cell lymphoma, lymphoblastic lymphoma and in the tumor stages of Hodgkin's disease and mycosis fungoides. The tumor masses may be sharply demarcated, but usually there are small, outlying islands of tumor cells and, in addition, single rows of lymphoma




FIG. 240 Stem-cell lymphoma. High magnification of Figure 239. The tumor cells possess abundant, pale staining cytoplasm and large round nuclei. The nuclei contain delicate dustlike chromatin and one or several prominent nucleoli. Mitotic figures are numerous. A large atypical mitotic figure (M) is present in the upper right corner ( $\times 400$ ).

phoma cells which extend from the main tumor mass into the

are in evidence at the breast—see pages 469 &

ly:

th

as may occur even in the subcutaneous tissue. They usually show a blood vessel in their center around which the infiltrate lies as a thick sheath.

An inflammatory infiltrate intermixed with tumor cells is the usual picture in Hodgkin's granuloma and in the erythematous and the

stem cells, reticulum cells and lymphoblasts, or mature lymphoid reticular cells, such as histiocytes and lymphocytes. Usually atypical cells and a varying number of mitotic figures are present. Nonspecific lesions consist of an inflammatory infiltrate without immature cells.




FIG 259 Stem cell lymphoma. Low magnification. A large mass of loosely packed tumor cells is present in the dermis ( $\times 100$ ).

It is probable that most if not all nonspecific lesions result from lymphoma cells (Gates). It may be assumed that the lymphoma cells either have been overwhelmed and removed by the inflammatory infiltrate which they themselves have provoked or do not differ sufficiently in their appearance from the inflammatory cells to be recognized as lymphoma cells.

Large masses of lymphoma cells in the dermis and the subcutaneous layer, which clinically appear as cutaneous tumors may occur in all

ture reticular cell (histiocyte) [The author prefers to call the immature reticular cell a reticulum cell, and the mature reticular cell a histiocyte. However, it should be kept in mind that many authors use the term reticulum cell also for the mature reticular cell (see page 35)] The reticulum cells in reticulum-cell lymphoma have an eosinophilic cytoplasm the border of which tends to be irregular in




FIG. 261 Reticulum-cell lymphoma. The tumor cells possess abundant pale-staining cytoplasm and variously shaped nuclei. Some are round but most are oval or kidney shaped. The nuclei are pale staining and possess a distinct nuclear membrane. ( $\times 400$ )

outline suggesting ameboid propensities. The nuclei differ in configuration: a few are round, more are oval, and still others are kidney shaped. They are pale staining and appear vesicular because of the presence of a well-defined, strongly basophilic nuclear membrane (Fig. 261). They show a moderately heavy chromatin network, nucleoli are rarely evident. A moderate number of mitotic figures is present. The reticulum cells, being immature, form only a scanty reticulum network. In most tumors a fair number of mature reticular cells (histiocytes) are present. Tumors containing many histiocytes may have a well-developed reticulum network. Lymphocytes are sometimes also found, presumably evidence of exudative reaction (Wayson and Weidman, Gall and Mallory, Director and Kern).

plaque stages of mycosis fungoides. This type of infiltrate, too, tends to have a patchy arrangement.

So called nonspecific lesions, which show a nonspecific chronic inflammatory infiltrate, occur in the skin, particularly in early mycosis fungoides, in Hodgkin's disease and in lymphocytic lymphoma.

### 1 STEM CELL LYMPHOMA

Stem cell lymphoma produces, as a rule, tumorlike lesions. In about one third of the cases, it arises as a single lesion, either in the skin or elsewhere, and may remain localized for some time. Leukemia is very rare.

**Histopathology.** The lesions are composed of large masses of cells (Fig. 259). The vast majority of cells are stem cells, but most tumors show an admixture of reticulum cells. Stem cells are large in size and possess abundant, pale staining cytoplasm. They may lie separate from one another, but frequently the cytoplasm of neighboring cells is fused. Their nuclei are large, from two to four times the size of a normal lymphocyte, round and filled with delicate, dustlike chromatin. They usually possess one but occasionally several large, deeply staining nucleoli (Fig. 260). Typical and atypical mitotic figures are numerous. Stem cells are too immature to form reticulum fibers (Gill and Mallory).

### 2 RETICULUM CELL LYMPHOMA

Tumorlike lesions are the most common type. In about one third of the cases, reticulum cell lymphoma, like stem cell lymphoma, starts as a solitary lesion. Occasionally, the first lesion or group of lesions occurs in the skin (Director and Keim).

If leukemia develops in reticulum cell lymphoma, it is of the monocytic variety, since the blood monocyte is derived from the reticular group of cells (Herbut and Miller). True monocytic leukemia is sometimes referred to as the Schilling type of monocytic leukemia, in contrast to the Megrel type of monocytic leukemia in which monocyte like cells arise from myeloblasts and which therefore is a form of myeloid leukemia (Montgomery and Watkins) (see page 195). In the presence of monocytic leukemia the skin frequently shows purpuric, vesicular and papular lesions (Herbut and Miller). In addition, plaques and tumorlike lesions may be present (Wayson and Weidman). Exfoliative dermatitis may occur (Montgomery and Watkins). Swelling and ulceration of the gums are frequent.

**Histopathology.** In tumorlike lesions, large compact masses of cells are present. The type cell is an immature reticular cell (reticulum cell). This cell is smaller than a stem cell but larger than a ma-

ture reticular cell (histiocyte) [The author prefers to call the immature reticular cell a reticulum cell, and the mature reticular cell a histiocyte. However, it should be kept in mind that many authors use the term reticulum cell also for the mature reticular cell (see page 35)] The reticulum cells in reticulum cell lymphoma have an eosinophilic cytoplasm the border of which tends to be irregular in




FIG 261 Reticulum-cell lymphoma. The tumor cells possess abundant round eosinophilic cytoplasm.

outline suggesting ameboid propensities. The nuclei differ in configuration: a few are round, more are oval, and still others are kidney shaped. They are pale-staining and appear vesicular because of the presence of a well-defined, strongly basophilic nuclear membrane (Fig 261). They show a moderately heavy chromatin network, nucleoli are rarely evident. A moderate number of mitotic figures is present. The reticulum cells, being immature, form only a scanty reticulum network. In most tumors a fair number of mature reticular cells (histiocytes) are present. Tumors containing many histiocytes may have a well-developed reticulum network. Lymphocytes are sometimes also found, presumably evidence of exudative reaction (Wayson and Weidman, Gall and Mallory, Director and Kern).

In small cutaneous lesions, as they occur particularly in association with monocytic leukemia, the infiltrate is less extensive and the cells are of smaller size. Nevertheless, the cells have the same appearance as in the large lesions having indented or kidney shaped, pale nuclei with a distinct nuclear membrane (Hubler and Nether




FIG. 262 Lymphoblastic lymphoma. Low magnification. Masses of densely packed tumor cells are present in the lower dermis ( $\times 50$ )

ton). In the presence of monocytic leukemia, extravasation of erythrocytes is frequently found

**Differential Diagnosis.** It is important not to confuse reticulum-cell lymphoma with reticulo endotheliosis (histiocytosis) of the Letterer-Siwe and Hand Schuller Christian types (see page 262) Reticulum-cell lymphoma and reticulo endotheliosis are occasionally so difficult to differentiate histologically that in the past they have been confused (Sweetzer, Winer and Cumming) Reticulo endotheliosis, like reticulum cell lymphoma, shows large, irregularly shaped reticular cells (histiocytes) but shows no atypical reticular cells, par-

icularly no mitotic figures. The lesions contain, as a rule, eosinophils and one may find cholesterol deposits in the reticular cells.

### 3. LYMPHOBLASTIC LYMPHOMA

The cutaneous lesions consist predominantly of nodules, plaques and tumors. Purpura is not uncommon.




FIG. 263 Lymphoblastic lymphoma. High magnification of Figure 262. The cells have only little cytoplasm. The nuclei lie therefore more densely packed than in stem cell lymphoma or " "

4) *pericities*. A few lymphocytes are present ( $\times 100$ )

Leukemia develops in a large percentage of cases—according to Gall and Mallory, in 40 per cent. The type is lymphatic leukemia.

**Histopathology.** The cutaneous lesions show usually large masses of tumor cells and occasionally a patchy infiltrate. Lymphoblasts predominate, but in most lesions stem cells and lymphocytes are present in moderate number. Lymphoblasts possess a narrow basophilic rim of cytoplasm and large round or slightly indented nuclei. Because of the relatively small amount of cytoplasm, the nuclei lie close together much closer than in stem-cell or reticulum-cell lymphoma (Fig. 262). The nuclei are larger than those of lymphocytes and more uniform in appearance than those of stem cells or reticulum cells. The chromatin in the nuclei of lymphoblasts is distributed rather evenly and much less clumped than in lymphocytes, giving the nuclei



a vesicular appearance (Fig 263) Nucleoli are observed infrequently Mitotic figures are usually numerous A fine, evenly distributed reticulum framework is occasionally present (Gall and Mallory)

#### 4 LYMPHOCYTIC LYMPHOMA

Cutaneous nodules, plaques and tumors occur Exfoliative dermatitis is relatively common In addition, nonspecific lesions, especially




FIG 264 Lymphocytic lymphoma In the upper third of the photograph one sees the periphery of a large mass of tumor cells Rows of tumor cells extend from there between the collagen bundles The cells are indistinguishable from normal lymphocytes ( $\times 200$ )

purpules and petechiae, may be observed Of interest is the occasional occurrence of generalized herpes zoster in association with lymphocytic lymphoma (see page 491)

Lymphatic leukemia occurs in about half the cases of lymphocytic lymphoma (Gall and Mallory)

Histopathology. The specific cutaneous lesions may show either large masses of cells (Fig 264), scattered patches of cells (Fig 265) or, in the case of exfoliative dermatitis, a diffuse infiltrate in the upper dermis

In lesions showing large masses or scattered patches of cells, the predominating cell is indistinguishable from a normal lymphocyte. Lymphoblasts may be present in small numbers. Mitotic figures are sparse. Thus, the infiltrate appears homogeneous. At the periphery of the large masses one frequently sees rows of tumor cells extending between and even around intact collagen bundles in a similar way as it may be observed in scirrhous metastatic carcinoma.




FIG. 265 Lymphocytic lymphoma. Sharply demarcated, large patches of lymphocytes are distributed through the dermis. Most patches show a blood vessel in their center ( $\times 100$ ).

If the infiltrate consists of scattered patches, some difficulty may arise in differentiating lymphocytic lymphoma from chronic discoid lupus erythematosus in which the infiltrate is also patchy. In lymphocytic lymphoma, however, the patches are distributed indiscriminately throughout the dermis without predilection to the upper dermis and the vicinity of the cutaneous appendages. In addition the patches are uniformly composed of lymphocytes without an admixture of plasma cells and histiocytes as in lupus erythematosus. Furthermore the epidermal changes associated with lupus erythematosus are absent.

In instances of exfoliative dermatitis one finds a diffuse infiltrate in the upper dermis. It is composed, in addition to lymphocytes, of

a vesicular appearance (Fig 263) Nucleoli are observed infrequently Mitotic figures are usually numerous A fine, evenly distributed reticulum framework is occasionally present (Gall and Mallory)

#### 4 LYMPHOCYTIC LYMPHOMA

Cutaneous nodules, plaques and tumors occur Exfoliative dermatitis is relatively common In addition, nonspecific lesions especially




FIG 264 Lymphocytic lymphoma. In the periphery of the tumor one sees the periphery of the tumor cells extend from there and are indistinguishable from normal lymphocytes ( $\times 200$ )

papules and petechiae, may be observed Of interest is the occasional occurrence of generalized herpes zoster in association with lymphocytic lymphoma (see page 491)

Lymphatic leukemia occurs in about half the cases of lymphocytic lymphoma (Gall and Mallory)

**Histopathology.** The specific cutaneous lesions may show either large masses of cells (Fig 264) scattered patches of cells (Fig 265) or, in the case of exfoliative dermatitis a diffuse infiltrate in the upper dermis

## 6 HODGKIN'S DISEASE

In most instances Hodgkin's disease affects primarily and predominantly the lymph nodes. In rare instances the first lesions are noted in the skin (Reimann, Havens and Herbut).

Cutaneous lesions are observed quite frequently. Nonspecific lesions, however, are more common than specific lesions. Most commonly found are papular lesions which are intensely pruritic. Other lesions are eczematous patches, generalized exfoliative dermatitis, nodules or tumors. The latter frequently undergo ulceration (Senear and Caro). Generalized herpes zoster occurs occasionally (see page 491).

**Histopathology.** In the specific lesions of Hodgkin's disease two stages can be recognized: Hodgkin's granuloma and Hodgkin's sarcoma. Hodgkin's sarcoma may follow Hodgkin's granuloma, but the disease may show the morphology of Hodgkin's sarcoma at the onset (Jackson and Parker).

Hodgkin's granuloma is composed of a polymorphous infiltrate. The majority of the constituent cells are the usual components of a chronic inflammatory infiltrate, namely, eosinophils, neutrophils, lymphocytes, plasma cells, histiocytes, and fibroblasts. In addition

from stem cells. They occur as mononucleated and multinucleated cells. The mononucleated Sternberg-Reed giant cell possesses a very large, irregularly shaped nucleus. The multinucleated Sternberg-Reed giant cell has either a double nucleus (mirror image nucleus) or several nuclei of dissimilar size and shape clustered in the center of the cell. The nuclei of both mononucleated and multinucleated Sternberg-Reed cells contain a prominent nucleolus and heavy clumps of chromatin. Mitotic figures occur with moderate frequency in the stem cells, reticulum cells, and Sternberg-Reed cells. Hodgkin's granuloma has a tendency to focal necrosis and in lesions showing this phenomenon phagocytes are quite numerous. Eosinophils are usually but not invariably present. In some cases they constitute the predominant cell. Collagen production is the result of the natural evolution of maturing histiocytes into fibroblasts. Collagen production is scanty in the early phases and progresses steadily until broad fibrous septa separate the foci of cellularity into islands (Jackson and Parker).

Hodgkin's sarcoma differs from Hodgkin's granuloma by the preponderance of stem cells, reticulum cells, and Sternberg-Reed cells.

neutrophils, eosinophils and plasma cells, resulting in an inflammatory infiltrate of banal appearance. However, immature lymphocytes and mitotic figures here and there usually reveal the nature of the process. Sometimes one also finds in the lower cutis variously sized patches composed solely of lymphocytes, further facilitating the diagnosis (Keim).

### 5 FOLLICULAR LYMPHOMA (BRILL-SYMMERS)

Follicular lymphoma, also called Brill Symmers disease, occurs predominantly in the lymph nodes, causing considerable enlargement of the affected lymph nodes. The skin is involved in only rare instances, showing discrete, firm, brownish to reddish nodules (Gall Morrison and Scott).

Follicular lymphoma of the lymph nodes was described originally as a hyperplasia but is now generally regarded as lymphoma, because it may progress into other forms of lymphoma and is ultimately fatal.

**Histopathology.** The enlarged lymph nodes show replacement of the normal architecture by numerous round, follicle like nodules of varying size. These follicles are composed of densely packed lymphoblasts and are surrounded by a thin rim of normal small lymphocytes. Mitotic figures are present within them in small number. Very frequently, the follicles are partially separated from the surrounding stroma by fissures. Although these fissures are an artefact caused by shrinkage of tissue during fixation they are of considerable aid in the diagnosis of follicular lymphoma (Gall, Morrison and Scott).

The cutaneous lesions, in rare instances, show the same follicular pattern as the lymph nodes. Usually, however, the infiltrate has the appearance of either lymphocytic or lymphoblastic lymphoma.

**Differential Diagnosis.** Follicular lymphoma of the lymph nodes must be differentiated from reactive hyperplasia such as occurs in dermatopathic lymphadenitis (see page 75). It differs from the latter by the presence of mitotic figures, obliteration of the sinuses, absence of inflammatory cells and absence of phagocytosis.

Confusion of follicular lymphoma with dermatopathic lymphadenitis has caused several authors to describe erroneously the occurrence of generalized exfoliative dermatitis in Brill Symmers disease (Combes and Bluefarb, Rost). These cases actually were instances of idiopathic generalized exfoliative dermatitis with secondary dermatopathic lymphadenitis.

For differentiation of follicular lymphoma from lymphocytoma cutis, see page 491.

## 6 HODGKIN'S DISEASE

In most instances Hodgkin's disease affects primarily and predominantly the lymph nodes. In rare instances the first lesions are noted in the skin (Reimann, Havens and Herbut).

Cutaneous lesions are observed quite frequently. Nonspecific lesions however are more common than specific lesions. Most commonly found are papular lesions which are intensely pruritic. Other lesions are eczematous patches, generalized exfoliative dermatitis, nodules or tumors. The latter frequently undergo ulceration (Senear and Caro). Generalized herpes zoster occurs occasionally (see page 491).

**Histopathology.** In the specific lesions of Hodgkin's disease two stages can be recognized: Hodgkin's granuloma and Hodgkin's sarcoma. Hodgkin's sarcoma may follow Hodgkin's granuloma, but the disease may show the morphology of Hodgkin's sarcoma at the onset (Jackson and Parker).

Hodgkin's granuloma is composed of a polymorphous infiltrate. The majority of the constituent cells are the usual components of a chronic inflammatory infiltrate, namely, eosinophils, neutrophils, lymphocytes, plasma cells, histiocytes and fibroblasts. In addition

from stem cells. They occur as mononucleated and multinucleated cells. The mononucleated Sternberg-Reed giant cell possesses a very large, irregularly shaped nucleus. The multinucleated Sternberg-Reed giant cell has either a double nucleus (mirror image nucleus) or several nuclei of dissimilar size and shape clustered in the center of the cell. The nuclei of both mononucleated and multinucleated Sternberg-Reed cells contain a prominent nucleolus and heavy clumps of chromatin. Mitotic figures occur with moderate frequency in the stem cells, reticulum cells and Sternberg-Reed cells. Hodgkin's granuloma has a tendency to focal necrosis and in lesions showing this phenomenon phagocytes are quite numerous. Eosinophils are usually but not invariably present. In some cases they constitute the predominant cell. Collagen production is the result of the natural evolution of mature histiocytes into fibroblasts. Collagen production is scanty in the early phases and progresses steadily until broad fibrous septa separate the foci of cellularity into islands (Jackson and Parker).

Hodgkin's sarcoma differs from Hodgkin's granuloma by the preponderance of stem cells, reticulum cells and Sternberg-Reed cells.

over all other elements comprising the tumor. Multinucleated cells and mitotic figures are numerous. Fibrosis is absent or minimal. Except for the presence of Sternberg-Reed cells, the histologic picture may greatly resemble that of stem-cell or reticulum cell lymphoma (Gall and Mallory)



FIG. 266 Hodgkin's disease. Low magnification. There are two large masses of tumor cells. In addition, numerous small collections of tumor cells are present throughout the dermis ( $\times 50$ ).

The nodules and the tumors of the skin show large masses of cells in the dermis and not infrequently also in the subcutaneous fat (Fig. 266). The histologic picture is more apt to be that of Hodgkin's granuloma than that of Hodgkin's sarcoma and is rarely as typical as in the lymph nodes. The number of Sternberg-Reed cells, on the presence of which the diagnosis depends, is often small and fibrosis

is less pronounced. The presence of atypical reticulum cells with mitotic figures and of chronic inflammation may suggest mycosis fungoides. Even then, however, a thorough search of serial sections will usually show a few Sternberg Reed giant cells (Fig. 267). It is important not to confuse clumped reticulum or endothelial cells with multinucleated Sternberg Reed cells. Reticulum and endothelial




FIG. 267 Hodgkin's disease. High magnification of Figure 266. There is a dense polymorphous infiltrate. It contains many atypical reticulum cells and several mitotic figures (MF). A mononucleated Sternberg Reed giant cell (MO) is present on the left; a multinucleated Sternberg Reed giant cell (MW) on the right ( $\times 400$ ).

cells have more regularly shaped, paler staining nuclei than Sternberg Reed cells (Miller, Wrong).

In the presence of a  
matit

case c... because Sternberg Reed cells frequently are absent. In such instances, the histologic picture is either the same as in mycosis fungoides or shows merely a nonspecific chronic inflammatory infiltrate.

The papular lesions show, as a rule, merely a nonspecific infiltrate of lymphocytes and histiocytes. Occasionally, a few atypical reticulum cells with mitotic figures are present.



over all other elements comprising the tumor. Multinucleated cells and mitotic figures are numerous. Fibrosis is absent or minimal. Except for the presence of Sternberg Reed cells, the histologic picture may greatly resemble that of stem cell or reticulum cell lymphoma (Gall and Mallory).



FIG. 266 Hodgkin's disease. Low magnification. There are two large masses of tumor cells. In addition numerous small collections of tumor cells are present throughout the dermis ( $\times 50$ ).

The nodules and the tumors of the skin show large masses of cells in the dermis and not infrequently also in the subcutaneous fat (Fig. 266). The histologic picture is more apt to be that of Hodgkin's granuloma than that of Hodgkin's sarcoma and is rarely as typical as in the lymph nodes. The number of Sternberg Reed cells, on the presence of which the diagnosis depends, is often small and fibrosis

atrophicans vascularis (see page 305) develop either as a precursor or as a residuum of plaque like lesions

**Histogenesis** The histogenesis of mycosis fungoides is not solved fully. Some authors explain the multiplicity of cell types as due to combined lymphoid and myeloid stimulation (Reimann, Havens and Herbut). A most logical explanation is that first suggested by Fraser (1925, 1936). He regarded mycosis fungoides as a form of reticulum cell lymphoma and believed that all other cells represented merely a defense reaction of the tissue against the tumor cells. He pointed out that in the early stage of the disease the number of tumor cells is small and the inflammatory defense reaction pronounced. As the disease advances the number of tumor cells increases while the defense reaction slackens until finally in the tumor stage the tumor cells proliferate uninhibitedly. Fraser's theory finds support in the fact that also in other malignant diseases such as squamous-cell carcinoma and malignant melanoma an inflammatory tissue reaction tends to be present as long as the tumor cells are only moderately malignant but disappears as the malignancy of the tumor cells increases (see Squamous Cell Carcinoma, page 330 and Malignant Melanoma, page 460).

**Histopathology** The division of mycosis fungoides into three stages as described for the clinical picture pertains also to the histologic picture.

In the first or erythematous stage the histologic picture not infrequently shows merely a banal inflammatory reaction so that a diagnosis of mycosis fungoides may be impossible. In some instances however particularly if several specimens are taken for biopsy and serial sections are made areas may be found in which specific changes (described in the next paragraph) are present. A finding which should always arouse one's suspicion of mycosis fungoides is the presence of patches of cellular infiltrate deeper in the dermis than one would expect to find them in a banal inflammatory infiltrate. It must be emphasized that the decision whether or not early mycosis fungoides exists often is a very difficult one.

In the second or plaque stage the histologic picture usually is diagnostic. The following six changes may be present: (1) great multiplicity of cell types, (2) polymorphism of the histiocytes, (3)

of

in

de

and abscesses

The variety of cells present includes histiocytes (= mature reticular cells), reticulum cells (= immature reticular cells), lymphocytes, neutrophils, eosinophils, plasma cells and fibroblasts. The number of

## 7. MYCOSIS FUNGOIDES

Mycosis fungoides affects primarily and predominantly the skin. In the late stage, however, the internal organs may show involvement (Berman; Post and Lincoln)

Several authors (Symmers; Gall and Mallory) have questioned the advisability of regarding mycosis fungoides as an entity. Symmers has called mycosis fungoides a "clinical and pathologic nonexistent," and states that cases reported as mycosis fungoides represent histologically "either Hodgkin's disease, reticulum cell sarcoma or lymphosarcoma." It is true that some cases which clinically appear as mycosis fungoides prove, on histologic examination, to be one of the other types of lymphoma, and that cases which at the onset show the histologic picture of mycosis fungoides later assume the histologic characteristics of some other form of lymphoma. Nevertheless mycosis fungoides has a distinct histologic picture which in the majority of cases remains as such throughout the disease and is found also in the internal lesions. Transformation of one form of lymphoma into another occurs not only in mycosis fungoides but also in other forms of lymphoma. This is evidence that all forms of lymphoma are related to each other but it does not obviate any classification.

Clinically, as well as histologically, mycosis fungoides can be divided into three stages, the erythematous ("premycotic" or 'pre fungoid') stage, the plaque stage and the tumor stage. All three stages may be present simultaneously and, occasionally, tumors develop without the previous presence of erythematous or plaque lesions (mycosis fungoides d'emblee) (Eller and Rein).

In the erythematous stage, the eruption may resemble eczema, psoriasis or parapsoriasis or have the appearance of generalized exfoliative dermatitis. Most commonly one observes scattered erythematous, scaling patches of irregular outline. They may be poorly demarcated or show a fairly sharp border. Since such lesions often resemble eczema closely it is always advisable to consider the possibility of early mycosis fungoides when dealing with an atypical chronic eczematoid eruption. Similarly, in cases of generalized exfoliative dermatitis of unknown genesis the possibility of mycosis fungoides must always be borne in mind. Itching is a prominent symptom in most instances of early mycosis fungoides.

In the plaque stage circinate well defined, elevated plaques are seen. They may show central clearing resulting in ringlike lesions.

In the tumor stage, one observes round or lobulated, raised tumors of bluish to brownish red color. They often undergo ulceration.

In rare instances of mycosis fungoides, lesions of poikiloderma

vessel in their center. If the patches are fairly large in size, they are strong evidence for lymphoma. An almost pathognomonic finding occasionally encountered is the presence of so-called Pautrier 'micro-abscesses' in the malpighian layer. They consist of small accumula-



FIG 269 Mycosis fungoides plaque stage. High magnification of Figure 268. The infiltrate in the upper dermis shows marked multiplicity of cell types and polymorphism of the histiocytes. Atypical reticulum cells with large hyperchromatic nuclei (mycosis cells) are numerous. There are several Pautrier micro-abscesses in the epidermis ( $\times 400$ ).

tions of cells mainly lymphocytes and histiocytes (Fig 269). (It should be recalled that the Munro micro-abscesses of psoriasis are located in the horny layer and are composed of neutrophils.) The epidermis in the plaque stage usually shows acanthosis with elongation of the rete ridges and may have an appearance similar to that found in psoriasis.

In the third or tumor stage the infiltrate consists of large masses of cells and may occupy large areas of the dermis and even penetrate

eosinophils varies but often is considerable. Histiocytes are numerous. Staining for reticulum with Foot's stain will show, commensurate with the rather large number of histiocytes, a well-developed network of reticulum fibers. The nuclei of the histiocytes vary greatly in size and shape. In addition, their nuclei may show pyknosis (shrinking), karyorrhexis (breaking up into particles, "nuclear dust")



FIG. 268 *Mycosis fungoides*, plaque stage. Low magnification. The infiltrate in the upper dermis is diffuse, while in the lower dermis it consists of variously sized sharply demarcated patches. The epidermis contains several Pautrier microabscesses (P.A.) ( $\times 100$ )

and clumping. Immature and atypical reticulum cells, called mycosis cells, are present. Mycosis cells differ from histiocytes by having larger, more irregularly shaped and more deeply staining nuclei. Mitotic figures, though not numerous, can, as a rule, be found. (An occasional mitotic figure may occur in a banal inflammatory infiltrate, so that the finding of one, two or even three mitotic figures in a section does not necessarily mean mycosis fungoides.) The infiltrate in the plaque stage of mycosis fungoides is located largely in the upper dermis just as in a nonspecific chronic inflammation. Frequently, however, one finds, in addition, patches of cellular infiltrate in the lower dermis (Fig. 268). These patches usually have a blood

vessel in their center. If the patches are fairly large in size, they are strong evidence for lymphoma. An almost pathognomonic finding, occasionally encountered, is the presence of so-called Pautrier "micro abscesses" in the malpighian layer. They consist of small accumula-



of the histiocytes. Atypical reticulum cells with large, hyperchromatic nuclei ( mycosis cells ) are numerous. There are several Pautrier "micro-abscesses" in the epidermis ( $\times 400$ )

tions of cells, mainly lymphocytes and histiocytes (Fig. 269). (It should be recalled that the Munro micro-abscesses of psoriasis are located in the horny layer and are composed of neutrophils.) The epidermis in the plaque stage usually shows acanthosis with elongation of the rete ridges and may have an appearance similar to that found in psoriasis.

In the third or tumor stage, the infiltrate consists of large masses of cells and may occupy large areas of the dermis and even penetrate

into the subcutaneous layer. The pressure of the infiltrate may destroy the epidermis so that ulceration results. The infiltrate still shows as a rule the same characteristics as in the plaque stage except that the number of mycosis cells and of mitotic figures is larger. In some cases, however, the mycosis cells attain considerable size and may possess more than one nucleus so that they resemble mononucleated or multinucleated Sternberg Reed cells.




FIG. 270. Mycosis fungoides tumor stage. The infiltrate is polymorphous. There are numerous mycosis cells. Some of them are of considerable size and one located in the center has two nuclei. These large mycosis cells resemble mononucleated or multinucleated Sternberg Reed cells ( $\times 400$ ).

cleaved or multinucleated Sternberg Reed giant cells (Fig. 270). If such cells are conspicuous the infiltrate may be identical with that of Hodgkin's disease (Fraser and Schwartz, Wile and Stiles). In other cases large immature reticulum cells like those seen in reticulum cell lymphoma may be present within the multicellular infiltrate (Fig. 271). Occasionally immature reticulum cells are so numerous and inflammatory cells so few that the infiltrate is identical with that of reticulum cell lymphoma (Fraser and Schwartz).

The histologic appearance of generalized exfoliative dermatitis in mycosis fungoides may be that of either the erythematous or the plaque stage. If it is that of the erythematous stage a diagnosis of mycosis fungoides may be impossible (Montgomery).

In poikiloderma atrophicans vasculare due to mycosis fungoides, the epidermis is atrophic and shows vacuolization of the cells of the basal layer. A dense infiltrate lies in the upper dermis in close approximation to the epidermis and invades the epidermis in some areas. The infiltrate may be that of either the erythematous or the plaque stage. As a rule, sufficient atypical cells are present to permit the diagnosis of mycosis fungoides (Oliver, Hazel, Dostrovsky, and




FIG 271 Mycosis fungoides tumor stage. The polymorphous infiltrate contains many large immature reticulum cells like those seen in reticulum-cell lymphoma ( $\times 400$ )

Sagher) (For a discussion of poikiloderma atrophicans vasculare see page 305.)

**Involvement of Internal Organs** The incidence of specific visceral lesions in mycosis fungoides is difficult to estimate from the literature. Many reviews set the figure too high, largely because the tendency exists in the dermatologic literature to regard a case of lymphoma as mycosis fungoides just because the primary or predominant lesions of the lymphoma are located on the skin. It should be stressed that for a diagnosis of mycosis fungoides it is necessary that the majority of the cutaneous lesions either show

as many cases of lymphoma that



were not mycosis fungoides, in a review of autopsies recorded in the literature found visceral involvement in 51 out of 54 cases. Crowley, Curtis and Leach, in their own material, found visceral lesions in 8 out of 10 cases of supposed mycosis fungoides, but only 1 case showed the multiplicity of cell types necessary for a diagnosis of mycosis fungoides and that case had no visceral lesions. It may be estimated that visceral lesions are found on autopsy in only about 20 per cent of the cases of mycosis fungoides, and usually they are not prominent or widespread.

When involvement of the internal organs is present in mycosis

cutaneous lesions (Ormsby and Tinnerud). On the other hand, the visceral lesions, just as some of the cutaneous lesions, may be composed largely of immature reticulum cells and be indistinguishable from reticulum cell lymphoma (Fraser and Schwartz) or may present, due to the presence of Sternberg Reed giant cells the picture of Hodgkin's disease (Fraser, Wile and Stiles, Poulsen).

Of the various internal organs, the subcutaneous lymph nodes are affected most frequently. However, the infiltrate in these lymph nodes is usually not that of mycosis fungoides but is nonspecific inflammatory, showing the histologic characteristics of dermatopathic lymphadenitis (see page 75). Nevertheless, the subcutaneous lymph nodes as well as the internal lymph nodes, the spleen, the lungs, the liver, the kidneys, the gastro intestinal tract and many other organs may be involved by mycosis fungoides (Berman, Gates, Post and Lincoln). Involvement of the bone marrow, however, has never been reported in true cases of mycosis fungoides (Poulsen). For this reason, mycosis fungoides shows no significant hemocytologic changes except that occasionally immature monocytes or lymphocytes are found in the blood. In rare instances true lymphatic leukemia develops in the terminal phase of the disease (Lane and Greenwood).

#### GENERALIZED EXFOLIATIVE DERMATITIS IN LYMPHOMA

Generalized exfoliative dermatitis is a fairly common occurrence in lymphoma. It occurs most commonly in lymphocytic lymphoma, Hodgkin's disease and mycosis fungoides, but also in reticulum cell lymphoma, particularly when accompanied by monocytic leukemia. Therefore, every case of persistent generalized exfoliative dermatitis requires investigation into the possibility of lymphoma. According to Montgomery, 25 per cent of all cases of generalized exfoliative dermatitis prove to be associated with lymphoma. The histologic picture of generalized exfoliative dermatitis in lymphoma may be,

in the early stage that of a nonspecific chronic inflammation. In cases in which it is doubtful whether or not lymphoma exists thorough hemocytologic studies, lymph node biopsy and sternal marrow biopsy are indicated and additional skin biopsies should be performed at intervals.

### HERPES ZOSTER IN LYMPHOMA

The relatively frequent association of herpes zoster particularly of herpes zoster generalisatus with lymphoma is of interest. Herpes zoster is most likely to occur in Hodgkin's disease and in lymphocytic lymphoma especially when lymphatic leukemia is present.

**Histopathology.** Histologic examination of the herpes zoster lesions may show the presence of a lymphomatous infiltrate (Barney Barton and O'Leary). In addition autopsy may reveal lymphomatous cells in the intercostal nerve in the spinal ganglion in the nerve roots or in the cord segment corresponding to the site of the herpes zoster (Bluefarb). It may be assumed that the lymphomatous infiltrate produces a locus minoris resistentiae where the zoster virus localizes and multiplies.

### LYMPHOCYTOMA CUTIS (LYMPHADENOSIS BENIGNA CUTIS SPIEGLER-FENDT SARCOID)

This condition occurs in two types: a localized type (Loveman and Fliegelman, Mopper and Rogin) and a disseminated type (Bäckerstedt). In the localized type there is either a solitary nodule or a group of nodules; in the disseminated type the lesions are scattered widely. The face and the ear lobes are the sites of predilection. The nodules are soft, asymptomatic and radiosensitive. Development into a lymphoma does not seem to occur.

**Histopathology.** A heavy infiltrate is present in the dermis usually separated from the epidermis by a narrow zone of normal collagen. The infiltrate occasionally consists only of mature lymphocytes but in most cases there are also reticulum cells. The two types of cells lie either intermingled with one another or in a follicular arrangement. In the latter type of arrangement lymphocytes surround islands of reticulum cells resulting in structures resembling the follicles of lymph nodes (Fig. 272). The lymphocytes and the reticulum cells are differentiated easily. The lymphocytes have small round deeply staining nuclei lying closely packed because lymphocytes possess only little cytoplasm. The reticulum cells have large irregularly shaped pale nuclei lying in loose arrangement here and there.

The nature of lymphocytoma cutis is not known. Although it suggests a lymphoma in its histologic architecture and is highly radio sensitive, it apparently is not a form of lymphoma. European authors (Hallam and Vickers; Hellier, Balverstedt) have suggested that it represents a hyperplasia of pre-existing rudimentary lymphoid tissue.



FIG. 272 Lymphocytoma cutis (lymphadenosis benigna cutis, Spiegler Fendt sarcoïd). Low magnification. The infiltrate is composed of two types of cells: lymphocytes, which lie in the dark staining areas, and reticulum cells, which lie in the light staining areas. At the right (L.F.), the arrangement of these cells resembles that encountered in a lymph follicle ( $\times 50$ ).

**Differential Diagnosis.** The diagnosis is easily established in those cases of lymphocytoma cutis in which the infiltrate shows lymphocytes and reticulum cells in follicular arrangement. Even in cases in which the two cell types lie intermingled, the diagnosis usually can be made because the cells composing lymphocytoma, in contrast with those of reticulum cell lymphoma, are mature. Nevertheless, mistakes in diagnosis occur (Director and Kern). The greatest difficulties in diagnosis occur in those cases with a purely lymphocytic infiltrate because it may be impossible to rule out lymphocytic lymphoma. It is thus advisable always to make a diagnosis of lymphocytoma cutis with reservations, except in cases with obvious follicular formations (Loveman and Fliegelman).

Differentiation from follicular lymphoma (Brill Symmers disease) with which lymphocytoma cutis is not related is not difficult since in follicular lymphoma the follicular centers are larger more numerous and outlined more distinctly. Also they are partially separated from the surrounding stroma by fissures which probably are the result of shrinkage of tissue during fixation. Such fissures are not apt to occur in lymphocytoma cutis.




FIG. 273 Lymphocytoma cutis (lymphadenosis benigna cutis Spiegler Fendt sarcoid). High magnification of Figure 272. Lymphocytes lie in the left upper half; reticulum cells in the right lower half of the illustration ( $\times 200$ ).

### MYELOSIS

Myelosis is nearly always associated with leukemia in contrast with lymphoma in which leukemia is often absent. The type of leukemia is myeloid leukemia. On the other hand, cutaneous lesions are less common in myelosis than in lymphoma. Specific and nonspecific cutaneous lesions may occur, however. Just as in lymphoma, there is no clearcut distinction between the two. The two conditions merge imperceptibly and it is probable that most if not all of the nonspecific cutaneous manifestations result from tumor cells (Gates) (see page 472).

Clinically, the specific cutaneous lesions consist of from pinhead to walnut sized, firm nodules, which may coalesce into plaques and occasionally ulcerate. The nonspecific cutaneous lesions (leukemids) may consist of macules, papules, pustules and purpuric lesions.

**Histopathology.** Histologic examination of specific lesions shows the presence of dense masses of cells in the dermis and, occasionally, in the subcutaneous layer. Frequently, rows of cells extend from these masses into the surrounding tissue. In addition, small groups of cells may be found scattered in the tissue spaces. The majority of cells belong to the myeloid series, but lymphocytes and phagocytic histiocytes not infrequently are observed (Ketron and Gay). Some of the cells of the myeloid series are mature, but most are immature, being either myeloblasts or myelocytes. They may be so immature that it is impossible to decide whether they belong to the myeloid or to the lymphoid reticular series of cells.

Myeloblasts are about double the size of a mature polymorphonuclear leukocyte. They possess a large, oval or round, pale staining nucleus and relatively little, nongranular cytoplasm. They resemble lymphoblasts, although, as a rule, lymphoblasts have a coarser chromatin structure and less cytoplasm (Nekam). Myelocytes are about the same size as myeloblasts or slightly smaller. They have, however, a smaller indented or lobate nucleus and abundant cytoplasm containing granules which may be either fine neutrophilic or coarse eosinophilic. Myeloblasts may or may not give a positive peroxidase reaction (also called oxidase reaction), while myelocytes always give a positive peroxidase reaction.

The peroxidase reaction indicates the presence of the enzyme peroxidase, which occurs in mature and partly matured cells of the myeloid series but is absent in very immature cells of the myeloid series and in all cells of the lymphoid reticular series. The peroxidase reaction is of great diagnostic value since it is positive in the majority of cases with myelosis and always negative in lymphoma. Even very immature tumors of myelosis usually show at least a few areas in which the reaction is positive. For the peroxidase reaction the specimen should be fixed in a 10 per cent formalin solution. Frozen sections are used for staining. On staining peroxidase positive cells show numerous black granules (Ketron and Gay).

Occasionally, the myeloid cells found in the cutaneous lesions are more immature than those in the bone marrow and the circulating blood. This may be regarded as evidence that the cutaneous lesions in myelosis are autochthonous (Paul and Imrizzi). It is probable that the myeloid cells form by myeloid transmutation of local reticular cells (Heller, Lewisohn and Palin).



- Eller J J and Rein C R *Mycosis fungoides à tumeur d'emblée solitaire* Arch Dermat & Syph 27 807 1933
- Fraser J T *Mycosis fungoides* Arch Dermat & Syph 12 814 1925
- and Schwartz H J *Neoplastic disease of the reticuloendothelial system* Arch Dermat & Syph 33 1 1936 (Good discussion)
- Gall E A and Mallory T B *Malignant lymphoma* Am J Path 18 381 1949 (Classification)
- Morrison H R and Scott A T *The follicular type of malignant lymphoma* Ann Int Med 14 2073 1941
- Gates O *Cutaneous tumors in leukemia and lymphoma* Arch Dermat & Syph 37 1015 1938
- Hazel O G *Poikiloderma atrophicum vasculare* Arch Dermat & Syph 40 776 1939
- Hente H J and Socha P *Häufigkeitsanalytische Untersuchungen zur Symptomatologie der Mycosis fungoides* Arch f Dermat u Syph 193 118 1951
- Herbut P A and Miller F R *Histopathology of monocytic leukemia* Am J Path 23 93 1947
- — and Erf L A *The relation of Hodgkin's disease lymphosarcoma and reticulum cell sarcoma* Am J Path 21 233 1945
- Hubler W R and Aetherton E W *Cutaneous manifestations of monocytic leukemia* Arch Dermat & Syph 56 70 1947
- Jackson H and Parker F *Hodgkin's disease—II Pathology* New England J Med 231 35 1944 (Good review)
- Keim H L *The lymphoblastomas their interrelationship* Arch Dermat & Syph 19 533 1929
- Lane C G and Greenwood A M *Lymphoblastoma (mycosis fungoides) and hemorrhagic sarcoma of Kaposi in the same person* Arch Dermat & Syph 27 643 1933
- Miller H *Lymphogranulomatosis cutis* Arch Dermat & Syph 17 156 1928
- Montgomery H *Exfoliative dermatitis and malignant erythrodermia* Arch Dermat & Syph 27 253 1933
- and Watkins C H *Monocytic leukemia cutaneous manifestations of the Naegeli and Schilling types hemocytologic differentiation* Arch Int Med 60 51 1937
- Oliver E A *Mycosis fungoides with poikiloderma like symptoms* Arch Dermat & Syph 33 267 1936
- Qewashy O S and Finnerud C W *Mycosis fungoides* Arch Dermat & Syph 27 631 1933
- Post C F and Lincoln C S *Mycosis fungoides* J Invest Dermat 10 135 1948
- Poulsen A *On mycosis fungoides* Acta dermat venercol 21 365 1940
- Reimann H A Havens W P and Herbut P A *Hodgkin's disease with specific lesions appearing first in the skin* Arch Int Med 70 431 1942
- Rost G A *Die Symmersche Erkrankung* Arch f Dermat u Syph 187 351 1949
- Senear F E and Caro M R *Ulcerative Hodgkin's disease of the skin* Arch Dermat & Syph 35 114 1937
- Sweitzer S E Winer L H and Cumming H A *Reticuloendotheliosis* Arch Dermat & Syph 40 192 1939
- Symmers D *Mycosis fungoides is a clinical and pathologic nonexistent* Arch Dermat & Syph 23 1 1932
- Wayson J T and Weidman F D *Aleukemic reticulosis* Arch Dermat & Syph 34 755 1936

Wile U J and Sules F Clinical mutations in lymphoblastomas JAMA 104 532 1935

Winer L H Mycosis fungoides Arch Dermat & Syph 56 480 1947

Wrong N M Hodgkin's disease of the scalp Arch Dermat & Syph 33 259 1936

### Lymphocytoma Cutis

Balverstedt B Ueber Lymphadenosis benigna cutis Acta dermat venereol 24 suppl XI 1945

Director W and Kern A B Reticulum cell sarcoma Arch Dermat & Syph 62 69 1950

Hallam R and Vickers H R Two cases of miliary lymphocytoma or benign lymphadenoid granuloma of the skin Brit J Dermat 51 251 1959

Hellier F F Lymphocytoma of the face Brit J Dermat 51 260 1959

Loveman A B and Fliegelman M T Lymphocytoma cutis Arch Dermat & Syph 63 169 1951

Mopper C and Rogin J R Benign solitary lymphocytoma Arch Dermat & Syph 63 184 1951

### Myelosis

Carmel W J Minno A M and Cook W L Eosinophilic leukemia with report of a case Arch Int Med 87 280 1951

Deme I Eosinophile Leukämie mit Hautsymptomen Dermatolog 98 150 1949

Gates O Cutaneous tumors in leukemia and lymphoma Arch Dermat & Syph 37 1015 1938

Goodman E G and Iverson L Chloroma a clinico-pathologic study of two cases Am J M Sc 211 205 1946

Heller E L Lewisohn M G and Palin W E Aleukemic myelosis Am J Path 23 327 1947

Hollender L Kaslin G J Permar H H and Schmitt C L Myeloid leukemia with cutaneous manifestations Arch Dermat & Syph 29 821 1934

Heuton L W and Gay L N Myeloid leukemia of the skin Arch Dermat & Syph 7 176 1923

Montgomery H and Watkins C H Monocytic leukemia cutaneous manifestations of the Naegeli and Schilling types hemocytologic differentiation Arch Int Med 60 51 1937

Nekam L Ein Fall von myeloischer Leukämie mit Beteiligung der Haut Dermat Ztschr 62 158 1931

Paul J T and Lanza L R Specific cutaneous lesions in chronic myeloid leukemia Arch Dermat & Syph 45 897 1942

Watkins C H and Hall B E Monocytic leukemia of Naegeli and Schilling types Am J Clin Path 10 387 1940 1941



- Eller J J and Rein C R Mycosis fungoides à tumeur d'emblée solitaire Arch Dermat & Syph 27 807 1933
- Fraser J F Mycosis fungoides Arch Dermat & Syph 12 814 1925
- and Schwartz H J Neoplastic disease of the reticuloendothelial system Arch Dermat & Syph 33 1 1936 (Good discussion)
- Gall E A and Mallory T B Malignant lymphoma Am J Path 18 381 1912 (Classification)
- Morrison H R and Scott A T The follicular type of malignant lymphoma Ann Int Med 14 2073 1941
- Gates O Cutaneous tumors in leukemia and lymphoma Arch Dermat & Syph 37 1015 1938
- Hazel O G Poikiloderma atrophicum vasculare Arch Dermat & Syph 40 716 1939
- Heite H J and Sochr P Häufigkeitsanalytische Untersuchungen zur Symptomatologie der Mycosis fungoides Arch f Dermat u Syph 191 118 1951
- Herbut P A and Miller F R Histopathology of monocytic leukemia Am J Path 23 93 1917
- and Erf L A The relation of Hodgkin's disease lymphosarcoma and reticulum cell sarcoma Am J Path 21 233 1913
- Hubler W R and Netherton E W Cutaneous manifestations of monocytic leukemia Arch Dermat & Syph 56 70 1917
- Jackson H and Parker F Hodgkin's disease—II Pathology New England J Med 231 35 1944 (Good review)
- Keim H F The lymphoblastomas their interrelationship Arch Dermat & Syph 19 533 1929
- Lane C G and Greenood A M Lymphoblastoma (mycosis fungoides) and hemorrhagic sarcoma of Kaposi in the same person Arch Dermat & Syph 27 643 1933
- Miller H Lymphogranulomatosis cutis Arch Dermat & Syph 17 156 1928
- Montgomery H Exfoliative dermatitis and malignant erythroderma Arch Dermat & Syph 27 253 1933
- and Watkins C H Monocytic leukemia cutaneous manifestations of the Naegeli and Schilling types hemocytologic differentiation Arch Int Med 60 51 1937
- Oliver E A Mycosis fungoides with poikiloderma like symptoms Arch Dermat & Syph 33 267 1936
- Ormsby O S and Finnerud C W Mycosis fungoides Arch Dermat & Syph 27 631 1933
- Post C F and Lincoln C S Mycosis fungoides J Invest Dermat 10 135 1948
- Poulsen A On mycosis fungoides Acta dermat venerol 21 365 1910
- Reimann H A Havens W P and Herbut P A Hodgkin's disease with specific lesions appearing first in the skin Arch Int Med 70 434 1942
- Rost G A Die Symmerssche Erkrankung Arch f Dermat u Syph 187 331 1919
- Senechal F E and Caro M R Ulcerative Hodgkin's disease of the skin Arch Dermat & Syph 35 114 1937
- Sweitzer S E Winer L H and Cumming H A Reticuloendotheliosis Arch Dermat & Syph 40 192 1939
- Symmers D Mycosis fungoides as a clinical and pathologic nonexistent Arch Dermat & Syph 25 1 1932
- Wayson J T and Weidman F D Leukemic reticulosis Arch Dermat & Syph 34 755 1936

# Glossary

④ **Acantholysis** Loss of coherence between epidermal cells due to degeneration of the intercellular bridges. It leads to the formation of clefts, vesicles and bullae within the epidermis. Occurs in pemphigus vulgaris, pemphigus foliaceus, pemphigus erythematosus, pemphigus neonatorum, pemphigus herpetiformis, pemphigus vegetans, pemphigus coccidioides, pemphigus paraneoplastic, pemphigus parapsoriasis, pemphigus paraneoplastic, pemphigus parapsoriasis, pemphigus paraneoplastic, pemphigus parapsoriasis.

④ **Acant**

④ **Alteration cavitaire** See Intracellular Edema

**Anaplasia** Atypical differentiation of cells occurring in malignant diseases. Anaplastic cells have large hyperchromatic irregularly shaped nuclei and frequently show atypical mitotic figures.

④ **Bull** - - - - -

tion occurs in virus vesicles and is diagnostic of them. See also Reticular Degeneration of Epidermal Cells

④ **Bulla** A cavity forming either within or beneath the epidermis and filled with lymph fluid. A small bulla generally is called a vesicle and a slitlike bulla as seen in Darier's disease and senile keratosis a lacuna. About the different modes of formation of bullae see the introduction to Chapter 7.

**Caseation necrosis** A type of tissue death in which the affected area has lost its structural outline and consists of pale eosinophilic amorphous finely granular material. Unless the necrosis is far advanced some pyknotic nuclei are still present. No invasion of neutrophils occurs. Caseation necrosis occurs especially in tuberculosis, syphilis, granuloma annulare and beryllium granuloma.

④ **Dyskeratosis** Faulty keratinization of individual epidermal cells. There are two types, one occurring in benign diseases and the other in malignant diseases. Benign dyskeratosis occurs in Darier's disease and occasionally in familial benign chronic pemphigus and consists of the formation of corps ronds and grains (see page 49). Malignant dyskeratosis occurs especially in Bowen's disease but also in squamous cell carcinoma and senile keratosis and consists of premature and atypical keratinization of individual cells.



# Glossary

- ⑥ **Acantholysis** Loss of coherence between epidermal cells due to degeneration of the intercellular bridges. It leads to the formation of clefts, vesicles and bullae within the epidermis. Occurs in pemphigus, Darier's disease, familial benign chronic pemphigus, virus bullae and senile keratosis.
- ⑥ **Acanthosis** Increase in thickness of the stratum malpighii.
- ⑥ **Alération cavitaire** See Intracellular Edema.
- Anaplasia** Atypical de-differentiation of cells occurring in malignant diseases. Anaplastic cells have large hyperchromatic irregularly shaped nuclei and frequently show atypical mitotic figures.
- ⑥ **Ballooning degeneration** A type of degeneration of epidermal cells causing marked swelling with loss of the intercellular bridges. Acantholysis results and a bulla forms. Ballooning degeneration occurs in virus vesicles and is diagnostic of them. See also Reticular Degeneration of Epidermal Cells.
- ⑥ **Bulla** A cavity forming either within or beneath the epidermis and filled with lymph fluid. A small bulla generally is called a vesicle and a fistlike bulla as seen in Darier's disease and senile keratosis a lacuna. About the different modes of formation of bullae see the introduction to Chapter 7.
- Caseation necrosis** A type of tissue death in which the affected area has lost its structural outline and consists of pale eosinophilic amorphous finely granular material. Unless the necrosis is far advanced some pyknotic nuclei are still present. No invasion of neutrophils occurs. Caseation necrosis occurs especially in tuberculosis, syphilis, granuloma annulare and beryllium granuloma.
- ⑥ **Dyskeratosis** Faulty keratinization of individual epidermal cells. There are two types: one occurring in benign diseases and the other in malignant diseases. Benign dyskeratosis occurs in Darier's disease and occasionally in familial benign chronic pemphigus and consists of the formation of corps ronds and grains (see page 49). Malignant dyskeratosis occurs especially in Bowen's disease but also in squamous-cell carcinoma and senile keratosis and consists of premature and atypical keratinization of individual cells.

**Hydropic degeneration of basal cells:** See "Liquefaction Degeneration of Basal Cells"

- ⑤ **Hyperkeratosis:** Excessive thickness of the horny layer. If caused by excessive formation of keratin, the granular layer is also thickened, as in lichen planus and lupus erythematosus. If caused by retention of the horny layer, as in ichthyosis vulgaris, the granular layer is even smaller than normal.

**Incontinence of pigment:** Loss of melanin from the cells of the basal layer due to damage to these cells with accumulation of the melanin in the upper dermis, inside as well as outside of melanophores. It occurs in an idiopathic form in the disease called incontinentia pigmenti, and in a symptomatic form in lichen planus, lupus erythematosus, poikiloderma atrophicum vasculare, Riehl's melanosis and melanodermatitis toxica.

- ⑥ **Intercellular edema (spongiosis):** Edema between squamous cells causing an increase in the width of the spaces separating them. It occurs frequently in inflammatory processes of the skin especially in dermatitis eczema. It does not cause formation of bullae or vesicles but contributes to their increase in size. See also Intracellular Edema.

- ⑦ **Intracellular edema (altération cavitaire):** Edema within squamous cells. If severe, it results in reticular degeneration (see below) of the affected cells and in the formation of multilocular bullae.

**Karyorrhexis:** Fragmentation of nuclei.

**Liquefaction degeneration of basal cells:** A type of degeneration causing vacuolization and disintegration of basal cells. It occurs in incontinentia pigmenti, lichen planus, lupus erythematosus, poikiloderma atrophicum vasculare, Riehl's melanosis, melanodermatitis toxica and lichen sclerosus et atrophicus with its variants kraurosis vulvae and balanitis xerotica obliterans. In several of these diseases the liquefaction degeneration may cause incontinentia pigmenti (see above). In lichen planus, lupus erythematosus and lichen sclerosus et atrophicus it may cause the formation of subepidermal bullae (see under Formation of Bullae introduction to Chapter 7).

**Metachromasia:** The phenomenon of reacting with a different color than that of the dye used for the staining. Important examples of metachromasia occur in the staining of the granules of mast cells (see "Urticaria Pigmentosa") and in the staining of mucin (see "Myxedema") of amyloid (see "Amyloidosis") and of the fibrinoid material in fibrinoid degeneration (see "Acute Systemic Lupus Erythematosus"). All four materials

stain purple with toluidine blue methylene blue thionine and cresyl violet Incidentally all four materials because of the presence of polysaccharides stain deeply red with the periodic acid Schiff reaction

**Metaplasia** Change of one type of tissue into another as it occurs for instance in the formation of bone in scars and in calcifying epithelioma (see Osteoma)

⑤ **Micro-abscesses** Small accumulations of cells in the epidermis Two types of micro abscesses occur the micro abscess of Munro in psoriasis and the micro abscess of Pautrier in mycosis fungoides For their description see those two diseases

**Papilloma** A tumor or tumor like proliferation of the skin characterized by papillomatosis (see below) and hyperkeratosis Five diseases show this type of proliferation nevus verrucosus (Jadassohn) keratosis senilis basal cell papilloma verruca vulgaris and acanthosis nigricans In typical instances histologic differentiation of these five diseases is easy but occasionally no more specific diagnosis than papilloma can be made

**Papillomatosis** Upward proliferation of papillae causing the surface of the epidermis to show irregular undulation

⑥ **Parakeratosis Imperfecta** nuclei in the horn is absent in areas

**Pyknosis** Shrinking of nuclei

⑦ **Reticular degeneration of epidermal cells** A process in which severe intracellular edema causes bursting of epidermal cells and formation of a multilocular bulla The septa inside the bulla are formed by resisting cell walls Reticular degeneration plays a role in the formation of virus vesicles and those of contact dermatitis

⑧ **Spongiform pustule of Kogoj** A multilocular pustule located in the upper stratum malpighii and characterized by the presence of neutrophils inside of edematous epidermal cells The cellular walls of these epidermal cells traverse the pustule like the network of a sponge This type of pustule has a diagnostic appearance and occurs in eczematoid dermatitis of Hallopeau impetigo herpetiformis keratosis blennorrhagica and Reiter's disease

⑨ **Spongiosis** See Intercellular Edema

⑩ **Vesicle** A small bulla (see Bulla) No sharp borderline can be drawn between a vesicle and a bulla Generally the term bulla is preferred in histopathology except for very small lesions that clinically are hardly visible

**Hydropic degeneration of basal cells:** See "Liquefaction Degeneration of Basal Cells"

- ④ **Hyperkeratosis:** Excessive thickness of the horny layer. If caused by excessive formation of keratin, the granular layer is also thickened, as in lichen planus and lupus erythematosus. If caused by retention of the horny layer, as in ichthyosis vulgaris, the granular layer is even smaller than normal.

**Incontinence of pigment:** Loss of melanin from the cells of the basal layer due to damage to these cells with accumulation of the melanin in the upper dermis, inside as well as outside of melanophores. It occurs in an idiopathic form in the disease called incontinentia pigmenti, and in a symptomatic form in lichen planus, lupus erythematosus, poikiloderma atrophicum vasculare, Riehl's melanosis and melanodermatitis toxica.

- ④ **Intercellular edema (spongiosis):** Edema between squamous cells causing an increase in the width of the spaces separating them. It occurs frequently in inflammatory processes of the skin especially in dermatitis eczema. It does not cause formation of bullae or vesicles but contributes to their increase in size. See also 'Intracellular Edema'.

- ④ **Intracellular edema (altération cavitaire):** Edema within squamous cells. If severe it results in reticular degeneration (see below) of the affected cells and in the formation of multilocular bullae.

**Karyorrhexis:** Fragmentation of nuclei.

**Liquefaction degeneration of basal cells:** A type of degeneration causing vacuolization and disintegration of basal cells. It occurs in incontinentia pigmenti, lichen planus, lupus erythematosus, poikiloderma atrophicum vasculare, Riehl's melanosis, melanodermatitis toxica and lichen sclerosus et atrophicus with its variants kraurosis vulvae and balanitis xerotica obliterans. In several of these diseases the liquefaction degeneration may cause incontinentia pigmenti (see above). In lichen planus, lupus erythematosus and lichen sclerosus et atrophicus it may cause the formation of subepidermal bullae (see under 'Formation of Bullae', introduction to Chapter 7).

**Metachromasia:** The phenomenon of reacting with a different color than that of the dye used for the staining. Important examples of metachromasia occur in the staining of the granules of mast cells (see Urticaria Pigmentosa) and in the staining of mucin (see Myxedema) of amyloid (see 'Amyloidosis') and of the fibrinoid material in fibrinoid degeneration (see 'Acute Systemic Lupus Erythematosus'). All four materials

# Index

Numerals in boldface type indicate the main reference to the subject

- Abscesses intra epidermal 81 82 151  
     170 222  
 Abtropfung theory of Unna 450  
 Acantholytic, 50 52 64 65 66 79 82  
     83 85 86 87 88 89 240 244  
     328 499  
 Acanthosis 499  
     nigricans 282 302 324  
 Acne necrotica miliaris 121  
     rosacea 124 183  
     varioliformis 124  
     vulgaris 123  
         ossification in 442  
 Acrodermatitis chronica atrophicans  
     118 163  
     continua 100 102 104 172  
     pustulosa 102  
 Acrokeratosis verruciformis 53  
 Actrosclerosis 307  
 Actinomycosis 221 228  
 Addison's disease 281 284 285  
 Adeno-acanthoma of sweat glands 387  
 Adenocarcinoma of apocrine glands  
     318 342 383  
     eccrine sweat glands 318 386  
     sebaceous glands 318 385  
 Adenoma 318 340 347  
     apocrine 317 349 352  
     definition of 347  
     sebaceous 319 347 386  
     sebaceum (Pringle) 319 344 346  
 Adiponecrosis subcutanea neonatorum  
     133  
 Alergic granulomatosis 312  
 Avascular 32 89 127 171 311  
 Alimenta strata 355  
     cicatrizata 120  
     splillica 208 209  
 Altered in cavitate 64 68 499 500  
 Amyloid 161  
     fast on and staining for 28 29 30  
 Amyloidosis 165 272  
     Congo-red test, in 272  
 Amyloidosis—(Continued)  
     lichenoid 272, 274  
     primary systemic 272  
     secondary systemic 272 275  
 Anaphylactoid purpura 32 33 89 127  
     130 311 313  
 Anaplasia 317 499  
 Anetoderma 165  
 Angiokeratoma corporis diffusum 426  
     Mibelli 426  
 Angiolipoma 437 438 439  
 Angioma 422 *See also* Hemangioma  
     capillary 423 425  
     cavernous 422 423  
     senile 426  
     serpiginosum 129  
 Anhidrosis thermogenic 76  
 Anisotropy of fat 31 *See also* Polariz-  
     scopic examination  
 Anthrax 198  
 Apocrine adenomas 317 349 352  
     epitheliomas 351  
     gland nevus 319 346  
     glands 4 16 125 318 342 343 349  
         352 354 356 380  
     carcinoma of 318 342 389  
     embryology 4  
 Argyria 153 286  
 Arrectores pilorum 25  
 Arsenical carcinoma 155  
     keratosis 155 328 338  
 Asteroid inclusion bodies in sarcoid  
     190  
     spores in sporotrichosis 231  
 Aubrine dermatitis 150  
 Atopic dermatitis 33 38 67 68 71 72  
     244  
 Atrophoderma reticulatum 163  
 Atrophy macular 163  
     senile 157 296  
     of vulva 161 162, 328  
 Autonomic nerves of skin 21  
 Autotechnicon 30



**Villi** Elongated and often tortuous papillae which are covered, as a rule, with but one or two layers of epidermal cells and extend into a vesicle a bulla or a cystic cavity. Formation of villi is observed in Darier's disease, familial benign chronic pemphigus, pemphigus vulgaris, pemphigus vegetans, syringo-cystadenoma papilliferum and hidradenoma papilliferum.

# Index

Numerals in boldface type indicate the main reference to the subject

- Abscesses intra-epidermal 81 82 151  
     170 222  
 Atropfung theory of Unna 450  
 Acantholysis 50 52 61 63 66 79 82  
     83 85 86 87 88 89 240 244  
     398 493  
 Acanthosis 499  
     nigricans 282 322 394  
 Acne necrotica miliaris 124  
     rosacea 124 183  
     varioliformis 124  
     vulgaris 123  
         ossification in 442  
 Acrodermatitis chronica atrophicans  
     118 163  
     continua 100 102 104 172  
     pusulosa 192  
 Acrokeratosis verruciformis 53  
 Acroclerosi 307  
 Actinomycosis 221 228  
 Addison's disease 281 284 285  
 Adeno-canthoma of sweat glands 387  
 Adenocarcinoma of apocrine glands  
     318 349 389  
     eccrine sweat glands 318 386  
     sebaceous glands 318 385  
 Adenoma 318 390 347  
     apocrine 317 349 352  
     definition of 347  
     sebaceous 319 347 386  
     sebaceum (Pringle) 319 344 346  
 Adiponecrosis subcutanea neonatorum  
     153  
 Allergic granulomatosis 312  
     vasculitis 39 89 127 171 311  
 Alopecia areata 125  
     cicatricata 126  
     syphilitica 208 209  
 Alération cavitaire 64 68 499 500  
 Amyloid 161  
     fixation and staining for 28 29 30  
 Amyloidosis 165 272  
     Congo red test in 272  
 Amyloidosis—(Continued)  
     lichenoid 272 274  
     primary systemic 272  
     secondary systemic 272 275  
 Anaphylactoid purpura 32 33 89 127,  
     130 311 313  
 Anaplasia 317 499  
 Anetoderma 163  
 Angiokeratoma corporis diffusum 426  
     Mibelli 496  
 Angioleiomyoma 437 438 439  
 Angioma 422 *See also* Hemangioma  
     capillary 423 425  
     cavernous 422 425  
     senile 426  
     serpiginosum 129  
 Anhidrosis thermogenic 76  
 Anisotropy of fat 31 *See also* Polar  
     scope examination  
 Anthrax 198  
 Apocrine adenomas 317 349 352  
     epitheliomas 354  
     gland nevus 319 346  
     glands 4 16 125 318 342 345 349  
         352 354 356 389  
     carcinoma of 318 342 389  
     embryology 4  
 Argyria 153 286  
 Arrectores pilorum 25  
 Arsenical carcinoma 155  
     keratosis 155 328 338  
 Asteroid inclusion bodies in sarcoid  
     190  
     spores in sporotrichosis 231  
 Atabrine dermatitis 150  
 Atopic dermatitis 33 38 67 68 71 72  
     244  
 Atrophoderma reticulatum 163  
 Atrophy macular 163  
     senile 157 296  
     of vulva 161 162 328  
 Autonomic nerves of skin 21  
 Autotechnicon 30

- Bacterid pustular 101, 102  
 Brilantia xerotica obliterans 158 162  
 Balanoposthitis chronica circumscripta 339  
 Ballooning degeneration 65 66 239  
     213 244 499  
 Basal cell epithelioma 44 53 122 155  
     318 319 326 331 344 351  
     354 356 368 369 385 386  
     adenoid 319 372 376 377  
     cystic 319 353 372 376  
     differentiated 319 371 372 376  
         from squamous cell carcinoma 380  
     fibrosing 369 372 371  
     histogenesis 318 370  
     intra epidermal 376  
     keratotic 319 372 376 377 379 381  
     metastasis in 370  
     morpher like 369 372 374  
     nodulo ulcerative 369 372  
     occurrence in nevus sebaceous (Ja  
         drassohn) 344  
     syngonocystadenoma papilliferum  
         351  
     pigmented 369 372 373  
     primordial type 372  
     relationship to tricho epithelioma  
         361 379  
         basal cell papilloma 380  
         solid 372 379  
         superficial 155 369 372 374 400  
         undifferentiated 319 371 372  
 Basal cell papilloma 322 323 328 382  
     385  
 Basal cells 4 7 12  
 Basal layer of epidermis 7  
 Basal squamous cell epithelioma 379  
     380  
 Basement membrane 8 9  
 Basophilic degeneration of collagen  
     44 157 161 296  
     leukocytes 37  
 Berylliosis systemic 141 190  
 Beryllium granuloma 140 141  
 Biliary xanthomatosis 257 260  
 Biopsy technique for 1  
 Blastomycosis European (torulosis)  
     221 224 231  
     North American 36 152 221 222  
         225 226 334  
 Blisters classification 65  
     Blisters—(Continued)  
         definition 499 501  
         mode of formation 61  
     Blood vessels of the skin 24  
     Blue nevus 450 466  
     Boeck's sarcoid 188 See also Sarcoid  
         dosis  
     Bone formation in tumor etc etc  
     Brill Symmers disease 75 471 480 493  
     Bromoderma 151 152 170 334  
     Bronze diabetes 280  
     Bulla definition 499  
     Bullae classification 60  
         mode of formation 64  
     Burger Grütz type of xanthomatosis  
         261  
     Burn 60 50 334 380  
     Calcification 275 303 310 324 325  
         368 442  
     Calcifying epithelioma (Mafferbé) 36  
         159 326 366 442  
     Carcinosis cutis 275  
         metabolic 276  
         metastatic 275  
     Calcium fixation and staining for 28  
         29 30  
     Cancer en cuirasse 390 391 392 473  
     Carcinoma of apocrine glands 318  
         342 389  
         arsenical 155  
         basal cell See Basal-cell epithelioma  
         of breast metastases to skin 390  
         definition of 317  
         due to arsenic 155  
         of eccrine sweat glands 318 386  
         inflammatory of breast 390 391  
         of meibomian glands 385  
         metastatic of the skin 389  
         mixed 379  
         of sebaceous glands 318 385  
         squamous cell 329 See also Squa  
             mous-cell carcinoma  
         telangiectatic 390 392  
     Caseation necrosis 175 176 499  
     Cavernous angioma 422 423  
     Cellule chure 9 See also Melanocytes  
     Cervicous glands 16

- Chancre syphilitic 208 209  
   tuberculous 176  
 Chancroid 199  
 Chloroma 495  
 (cholesterol) 31  
 Cholesteatoma extracellular 31 257,  
   268  
 Chondrodermatitis nodularis chronica  
   helicis 135  
 Chromatophores 12 35  
 Chromoblastomycosis 221 223 225  
 Chronic undermining burrowing ulcer  
   170  
 Chrysiasis 154  
 Civatte's poikiloderma reticulare 283  
 Clasmatocyte 34 *See also* Histocyte  
 Clear cell 9 *See also* Melanocytes  
 Clear cell myo-epithelioma 361  
 Coccidioidomycosis 36 221 226  
 Collagen 5 19 34 35 36  
   basophilic degeneration of 44 157  
   164 296  
   development of 20 404  
   fibrinoid degeneration of 30 118  
   297 302 309 310  
 Collastin 157  
 Colloid degeneration of the skin 163  
   milium 163 275  
 Comedo 123  
 Condyloma acuminatum 208 246 251  
   latum 208 209  
 Congenital diseases 40  
   ectodermal defect 44  
   ichthyosiform erythroderma 41 65  
   lymphedema (Milroy's disease) 135  
   syphilis 209 213  
 Congo-red test in amyloidosis 272  
 Contact dermatitis 38 67 68 69 72  
   Corium *See* Dermis  
 Cornification of epidermis, 11  
   hair cells 19  
 Cornoid lamella 43  
 Cornu cutaneum 327 328  
 Corps ronds 43 48 49 338  
 Cryptococcosis 224 *See also* Torulosis  
 Cushing's disease 162  
 Cutaneous hyperelasticity 56  
   rhomboidal & rufae 157  
 Cylindroma 319 351 357 359 366  
 Cyst dermoid 324  
   epidermal 47 139 317 324 368  
   Cyst—(Continued)  
   myxomatous degeneration of the  
     skin 413  
   sebaceous 139 324 368 369  
   synovial of the skin 413  
 Darier's disease 45 48 52 65 77 81  
   263 328 338 350  
   bullous 52  
 Darier Roussy's subcutaneous sarcoid  
   188 303  
 Degeneration ballooning of the epi-  
   dermis 63 66 239 243 244  
   499  
   basophilic of collagen 44 157 164  
   296  
   fibrinoid of collagen 30 118 297  
   302 309 310  
   reticular of the epidermis 65 66 69  
   239 240 243 244  
   senile of the skin 157  
 Dehydrating of specimens 28  
 Delhi boil (leishmaniasis) 235  
 Dendritic cell 9 *See also* Melanocytes  
 Dermatitis 65 66 150 471 484  
   acute 32 34 68 218  
   atopic 33 38 67 68 71 72 244  
   chronic 34 70 73 106 123 125 211  
   218 220 287 303  
   contact 38 67 68 69 72  
   due to Atabrine (quinacrine hydro-  
     chloride) 120  
   fungus 67 218  
   lymphoma 67 471 483  
   generalized exfoliative 67 74 75  
   150 151 471 474 478 479  
   480 481 483 484 488 490  
   due to drugs 150 151  
   in lymphoma 68 74 75 471 474  
   478 479 481 483 484 488  
   490  
   herpetiformis 33 63 69 87 88 89  
   infectious eczematoid 67 72  
   papularis capillitis 169  
   pigmented purpuric lichenoid (Gou-  
     gerot and Blum) 129  
   seborrheic 67 68 73 100 101 263  
   solar 122  
   stasis 67 68 73 130 334  
   subacute 34 70 218 219 220  
   venenata *See* Dermatitis contact

- Dermatofibroma* 403, 408 410 411, 424
- Dermatofibrosarcoma protuberans* 406 408 410
- Dermatomyositis* 276 302 303 305 306  
 relationship to acute systemic lupus erythematosus 302  
 visceral lesions in 305
- Dermatophilic lymphadenitis* 75 480 490
- Dermatophytosis* 217  
 of hands and feet 217 218
- Dermatosis exudative discoid and lichenoid chronic* 67 73 211  
*papulosa nigra* 383  
 progressive pigmentary of Schimberg 129
- Dermis* 5 19  
 embryology 5
- Dermoid cyst* 324 326
- Dihydroxyphenylalanine (dopa)* 12
- Diphtheria of the skin* 197
- Diseases due to bacteria* 167  
*tungi* 217  
 protozoa 235  
 spirochetes 203  
 viruses 238
- Donovan bodies of granuloma inguinale* 36 200 201
- Dopa stain* 9 10 12 28 29 30 285 342 462 466  
 in blue nevus 466  
 in malignant melanoma 462  
 in Mongolian spot 466  
 in vitiligo 285
- Drug eruptions* 33 67 68 150  
 fixed 150
- Dyshidrotic eruption* 76
- Dyskeratosis* 499  
 congenita 42
- Dyskeratosis in arsenical keratosis* 153  
*Bowen's disease* 338  
*Darier's disease* 48  
*familial benign chronic pemphigus* 52  
*pemphigus foliaceus* 84  
*radiodermatitis* 121  
*senile keratosis* 328  
*squamous-cell carcinoma* 331
- Eccrine sweat glands* 13 16 76 77, 123  
 carcinoma of 318 386  
 embryology 4
- Ectodermal defect congenital* 41
- Eczema* 66 *See also* *Dermatitis nummular* 67 72  
*vaccinatum* 244
- Eczematous drug eruptions* 33 67 68 150
- Edema intercellular of epidermis* 68 500  
 intracellular of epidermis 68 500  
 of the legs hereditary (*Milroy's disease*) 133
- Ehlers Danlos syndrome* 56
- Elastic fibers* 5 19 20  
 fixation and staining of 20 29
- Elastosis senilis* 56 157
- Eleidin* 12
- Embedding* 28
- Embryology of apocrine sweat glands* 4  
 dermis 5  
 eccrine sweat glands 4  
 elastic fibers 5  
 epidermal appendages 4  
 epidermis 3  
 hair 4  
 melanocytes 5  
 sebaceous glands 4
- Embryonal cells definition of* 317
- Endocarditis in acute systemic lupus erythematosus* 300  
 subacute bacterial 127 170
- Endoneurium* 22 418 419 441
- Endothelial cells* 21 24 431 435 437
- Eosinophilic granuloma* 33 113 257 262 265  
 of the face 111 *See also* *Granuloma faciale*  
*leukemia* 495
- Eosinophils* 33
- Ephelis* 465
- Epidermal cyst* 47 139 317 324 368
- Epidermal nevus* 321 *See also* *Nevus verrucosus*
- Epidermal tumors histogenesis and classification* 317
- Epidermis basal cell layer* 4 7  
 herence with dermis 8 20 21  
 embryology 5

- Epidermis—(Continued)**  
 epitrichial layer 3  
 granular cell layer 7 11  
 histology 7  
 horny layer 7 11  
 nerves 13  
 pigment 5 9 12  
 squamous-cell layer 4 7 10  
 stratum lucidum 12
- Epidermodysplasia verruciformis** 41  
 52 250
- Epidermoid carcinoma** 329 *See also*  
 Squamous cell carcinoma
- Epidermolysis bullosa** 47 65 87 278  
 279
- Epineurium** 22
- Epithelioid cells** 34 36 188
- Epithelioma** 390  
 adenoides cysticum 363 *See also*  
 Tricho-epithelioma  
 apocrine 354  
 basal-cell 369 *See also* Basal cell ep-  
 ithelioma  
 benign 318 320 366  
 calcifying (Mallherbe) 36 139 319  
 326 366 442  
 definition of 390  
 multiple benign cystic 363 *See also*  
 Tricho-epithelioma  
 sebaceous 319 333 386  
 squamous-cell 329 *See also* Squa-  
 mous-cell carcinoma  
 superficial basal-cell 155 369 372  
 374 403
- Epitrichial layer** 3
- Eruptions due to drugs** 33 67 68 150
- Frysupelas** 32 168
- Erythema elevatum diutinum** 32 114  
 induratum 96 176 181 184 183  
 215 231  
 multiforme 47 65 69 87 89 150  
 313  
 nodosum 33 93 98 150 186 226
- Erythrasma** 218
- Erythroderma** congenital ichthyosi-  
 form 41 63  
 generalized *See* Exfoliative derma-  
 titis generalized
- Erythroplasia of Queyrat** 338
- European blastomycosis (torulosis)**  
 221 224 231
- Exfoliative dermatitis** generalized 67  
 74 75 150 151 471 474 478  
 479 480 481 483 481 488  
 490  
 due to drugs 150 151  
 in lymphoma 68 74 75 471 474  
 478 479 481 483 181 488  
 490
- Extracellular cholesterol** 31 257 268
- Exudative discoid and lichenoid**  
 chronic dermatosis 67 73 211
- Familial benign chronic pemphigus**  
 50 51 63 77 80  
 hemorrhagic telangiectasia 427
- Fat anisotropy** 31 *See also* Fat re-  
 fractility  
 necrosis subcutaneous of the new-  
 born 133 154  
 refractility 17 31 133 193 258 268  
 271 406
- Fatty tissue** mucoid degeneration of  
 299 303
- Favus** 218 220
- Feulgen reaction** 30 241 246 251 302
- Fibrinoid degeneration of collagen** 30  
 118 297 302 309 310
- Fibroblast** 5 20 34 35 36 470  
 differentiation from histiocyte 35
- Fibroma durum** 403 *See also* Dermato-  
 fibroma
- Fibromyxoma** 412
- Fibrosarcoma** 122 403 406 408 410  
 415 421 434 435 462 464
- Fibrosis** nodular subepidermal 403  
 404
- Fixation** 28
- Fixed drug eruption** 150
- Foam cells** 33 258 260 406 407
- Follicular lymphoma** 75 471 480 493
- Folliculitis** 32 150 168 219  
 barbae 168 220  
 decalvans 126  
 keloidalis 169  
 ulerythematosa reticulata 163
- Fordyce's disease** 319 346
- Foreign body giant cells** 36 139 189  
 213 270 278 368  
 granuloma 31 133 139 190
- Formalin fixation** 28
- Fox Fordyce disease** 125

- Freckle 465  
 Frisch bacilli of rhinoscleroma 35 202  
 Fungus dermatitis 67 218  
   diseases 175 217  
   staining for 31  
 Furuncle 168  
  
 Gaucher's disease 257 265  
 Generalized exfoliative dermatitis 67  
   74 75 150 151 471 474 478  
   479 480 481 483 484 488  
   490  
   due to drugs 150 151  
   in lymphoma 68 74 75 471 474  
   478 479 481 483 484 488  
   490  
 Generative cells of sebaceous glands  
   17 448  
 Genital nerve corpuscles 22 24  
 Genodermatoses 40  
 Ghon complex of tuberculosis 174  
   176  
 Giant cells foreign body 36 139 189  
   213 270 278 368  
   Langhans 36 175 189 213 222  
   Sternberg Reed 75 481 488 490  
   Touton 259 269 406 407  
 Gilchrist's disease 222 *See also* Blastomycosis North American  
 Gitterfasern 20 *See also* Reticulum fibers  
 Glands apocrine 16 *See also* Sweat glands  
   apocrine  
   ceruminous 16  
   eccrine 13 *See also* Sweat glands eccrine  
   mammary 16 353  
   meibomian 17 385  
   Moll's 16  
   sebaceous 17 *See also* Sebaceous glands  
 Glomerulitis wire loop 300 310  
 Glomus 24 423 429 430  
   tumor 429 431  
 Glycogen 30  
   staining for 30  
   storage disease 261  
 Gonorrheal keratosis 100 103 101  
   172  
 Gout 36 139 277  
 Grains (Darier's disease) 49 50 81  
 Granular cell myoblastoma 334 440  
 Granular layer of epidermis 7 11  
 Granuloma annulare 114 118 271  
   beryllium 140 141  
   eosinophilic 33 113 257 262 265  
   of the face 111 *See also* Granuloma faciale  
   faciale 33 111 114 265  
   foreign body 31 133 139 190  
   inguinal 37 200 204 232 236 237  
   334  
   lipid 140  
   lipophagic 133  
   pyogenicum 424  
   silicon 140 112  
   swimming pool 142  
   tattoo 140  
 Granulomatosis allergic 312  
   disiformis chronica progressiva 271  
 Guarnieri bodies in variola 238 243  
   246  
 Gumma 181 188 208 213 216 334  
   380  
  
 Hair 1 18  
   embryology of 4  
   keratinization of 19  
   nevus 319 345 346  
 Hairy pigmented nevus 458  
 Hamaroma 317 318 320 371 413  
 Hand Schuller-Christian disease 31  
   113 257 262 265 474  
 Helly's fixative 26  
 Hemangioendothelioma 435 437  
 Hemangioma 422 425  
   capillary 423 425  
   cavernosum 422 423  
   sclerosing 403 404 424  
   senile 126  
 Hemangiopericytoma 430  
 Hemangiosarcoma 435  
 Hemochromatosis 285  
 Hemosiderin 32 35 151 285  
   differentiation from melanin 285  
 Henle layer of hair sheath 1 19  
 Herpes 37 38  
 Hereditary edema of the legs 135  
 Herpes gestationis 88  
   simplex 65 238 239 243 244  
   zoster 65 238 239 243 478 481  
   491

- Herpes—(Continued)  
   zoster—(Continued)  
     in lymphoma 478 481 491  
 Fibroma 414  
   malignant 414 415  
 Hidradenitis suppurativa 169  
 Hidradenoma papilliferum 319 352  
 Hidrocytoma 123  
 Histiocyte 21 21 31 31 35 36 401 470  
   472 475 485  
   differentiation from fibroblast, 35  
 Histiocytoma 31 260 403 408  
 Histiocytosis 262 *See also* Reticulo-  
   endotheliosis  
 Histochemical staining methods 30  
 Histoplasmosis 36 202 204 221 221  
   231 236 237  
 Hodgkin's disease 33 75 137 191  
   224 231 433 471 473 474  
   481 484 488 490 491  
   sarcoma 481  
 Horn cyst 363 364 379 381  
   pearls 329 334 363 379 381  
 Horny layer of the epidermis 7 11  
 Hotchkiss McManus stain 8 50 73 99  
   218 271 225 230 231 272  
   279 297 299 501  
 Huxley layer of hair sheath 4 19  
 Hyaluronic acid 30 37 38 279 281  
   staining for 30  
 Hyaluronidase 38 279 281  
 Hydroa aestivale 46 65  
   vacciniforme 46 65 278  
 Hyperkeratosis 500  
 Hyperlipemia 31 257 261  
   idiopathic 257 261  
   secondary 257 261  
 Hyperpigmentation 282  
 Ichthyosiform erythroderma congeni-  
   tal 41 65  
 Ichthyosis 40 41 42 286 287  
   hystrix 41 321 *See also* Nevus ver-  
   rucosus  
 Impetigo 65 166  
   Bockhart 168  
   perpetiformis 103 104 172  
 Inclusion bodies asteroid 190  
   Schaumann 142 189  
   viral 238 241 242 243 244 246  
   249 252 253  
 Incontinence of pigment 500  
 Incontinentia pigmenti 59 65 284  
   symptomatic 109 284 295 306 500  
 Infectious eczematoid dermatitis 67  
   72  
 Insect bites 137  
 Inter cellular edema in epidermis 68  
   500  
 Intracellular edema in epidermis 68  
   500  
 Intradermal nevus pigmentosus 450  
   452 456 459 461  
 Iododerma 152  
 Iron fixation and staining for 29  
 Isotropy of fat 31 *See also* Polariscopic  
   examination  
 Jadassohn Lewandowsky law 173 195  
 Junction nevus 342 450 452 453 458  
   459 460 464 465  
 Juxta articular nodes in syphilis 208  
   215  
 Kala azar 235  
 Kaposi's sarcoma 431 435 437  
   varicelliform eruption 244  
 Karyorrhexis 175 486 500  
 Keloid 408  
 Keratinization of epidermis 11  
   of hair cells 19  
 Keratohyalin 11  
 Keratosis arsenical 155 328 338  
   blennorrhagica 100 103 104 172  
   follicularis 48 *See also* Darier's dis-  
   ease  
   gonorrheal 100 103 104 172  
   palmaris et plantaris 42  
   punctata palmaris et plantaris 42  
   53  
   seborrheica 382. *See also* Basal-cell  
   papilloma  
   senilis 44 65 77 155 317 322 323  
   326 328 334 338 385  
 Keratotic nevus 321  
 Kerion Celsi 217 219  
 Krause's vulvae 158 161, 162  
 Krause bodies 22, 23 24  
 Lacunae 49 50 51, 52  
 Lames foliacées 451 457  
 Langerhans cells 10 13



- Langhans giant cells 36 175 189 213 222
- Lattice fibers 20 *See also* Reticulum fibers
- L.E. cell in lupus erythematosus 300
- Leiomyoma 437
- Leishmaniasis 36 202 204 232 235 237
- Lentigo 150 461
- Lepra bacilli 35 195
- cells 191
- Leprosy 36 175 190 193
- neural lesions in 194 196
- Letterer-Siwe disease 257 262 263 265 474
- Leukemia 221 231 170
- eosinophilic 495
- lymphatic 477 478 490 491
- monocytic Niegeli type 474 495
- Schilling type 474 475 490 495
- myeloid 32 474 493 495
- Leukemia 471 491
- Leukocytes basophilic 37
- eosinophilic 33
- neutrophilic 32
- Leukoplakia 110 161 162 317 328
- Lewandowsky rosacea like tuberculid of 124 176 182
- Libman Sacks syndrome 300
- Lichen amyloidosis 165 272 274
- corneus hypertrophicus 67 72
- nitidus 110
- planopilaris 106 109
- planus 34 38 65 71 106 111 150 151 284 328
- et acuminatus atrophicus (Feldman) 107
- bullosus 65 106 109
- follicularis 106 109
- hypertrophicus 106 109
- sclerosus et atrophicus 65 158 161 162
- scrofulosorum 176 185
- simplex chronicus 67 71 72 95 101 111
- spinulosus et folliculitis decalvans (Little) 107
- spinulosus of Crocker 286
- striatus 111
- urticatus 138
- Lichenificatio gigantea 67 72
- Lichenoid amyloidosis 165 272 274
- tuberculid 183
- Lipid granuloma 140
- histiocytosis 262
- reticulo endotheliosis 257 262
- Lipids 28 31 32 35
- fixation and staining of 28 29 30
- refractility 17 31 133 195 258 268 271 406
- Lipoblastosis systemic multicentric 413
- Lipogranuloma sclerosing 140
- Lipoid proteinosis 31 257 266
- Lipoidoses classification 257
- Lipoma 113 414
- Lipomelanotic reticulosis 75
- Lipomyoma 412
- Lipoplagic granuloma 133
- Liposarcoma 411 415
- Lipschutz bodies in herpes zoster 238 243
- Liquefaction degeneration of basal cells 500
- Lupus erythematosus 31 38 65 71 84 123 126 163 293 305 306 479
- acute systemic 293 297 303 305 310
- relationship to dermatomyositis and scleroderma 302
- visceral lesions in 298
- chronic discoid 126 293 294 303
- hematoxylin staining bodies 300 301
- le cell 300
- profundus 303
- subacute disseminate 293 296 303 307
- miliaris disseminatus faciei 176 183
- pernio (Desmier) 188
- vulgaris 142 176 177 180 183 190 215 334 380
- differentiation from sarcoidosis 179
- Lymphadenitis dermatopathic 75 480 490
- Lymphadenosis benigna cutis 137 491
- Lymphangioma 427
- Lymphangio sarcoma post mastectomy 437
- Lymphatic leukemia 477 478 490 491

- Lymphatic vessels of the skin 24  
 Lymphedema congenital (Milroy's disease) 135  
 Lymphoblast 34 470 472 477  
 Lymphoblastic lymphoma 471 473 477 480  
 Lymphocyte 21 34 37 470 472 479  
 Lymphocytic lymphoma 296 435 471 473 474 478 480 490 491  
 Lymphocytoma cutis 137 188 480 491  
 Lymphogranuloma venereum 199 246 252  
 Lymphoid cells 35 410  
 Lymphoma 34 68 74 75 223 231 303 435 470 491 492 493  
     classification 471  
     dermatitic 67 471 483 484  
     follicular 75 471 480 493  
     generalized exfoliative dermatitis in 68 74 75 471 474 478 479 481 483 484 488 490  
     Hodgkin's type 33 75 137 191 221 231 435 471 473 474 481 484 488 490 491  
     lymphoblastic 471 473 477 480  
     lymphocytic 296 435 471 473 474 478 480 490 491  
     mycosis fungoides 434 *See also* Mycosis fungoides  
     reticulum cell 55 410 471 473 474 484 488 490 492  
     Sprengel-Fendt type 137 188 480 491  
     stem cell 403 410 471 473 474 482  
 Lymphopathia venerea 199 246 252  
 Lymphosarcoma 471 484  
  
 Macrophage 39 34 35 36  
 Macular atrophy 163  
 Madura foot 228  
 Majocchi's purpura annularis telangiectodes 129  
 Malherbe's calcifying epithelioma 36 139 319 376 366 442  
 Malignant melanoma 330 333 342 401 410 450 456 457 458 459  
     criteria for malignancy 464  
     dopa reaction in 462  
 Mammary glands 16 353  
 Mast cells 28 37 38 57  
 Mast cells—(Continued)  
     differentiation from basophilic leucocytes 37  
     fixation and staining 28 29 30  
 Meibomian glands 17  
     carcinoma of 385  
 Meissner tactile corpuscles 13 21 23 451 455  
 Melanin 5 7 10 12 13 18 32 35 285  
     differentiation from hemosiderin 285  
 Melanocytes 5 7 9 12 13 18  
     in basal cell epithelioma 374  
     in basal cell papilloma 383  
     in blue nevus 466  
     in hair 18  
     in Mongolian spot 466  
     in nevi 450  
     origin of 5  
 Melanodermatitis toxica 283  
 Melanoma juvenile 457 458 463  
     malignant 478  
 Melanophores 12 35  
 Melanosis of Riehl 283  
 Meningococcemia purpura in 127 171  
 Merkel Ranvier tactile cells, 13 21  
 Merkel's disks 13  
 Metachromasia 37 38 57 272 279 281 297 500  
 Metaplasia 501  
 Metastasis in basal cell epithelioma 310  
     Bowen's disease 333  
     fibrosarcoma 409 410  
     lymphoma 470  
     malignant melanoma 458  
     squamous cell carcinoma 333  
 Metastatic carcinoma of the skin 389 473  
 Micro-abscesses of Munro 73 98 100 487 501  
     Fautrier 72 487 487 501  
 Micropapular tubercle 176 182  
 Microphage 32  
 Mikulicz cell of rhinoscleroma 202  
 Malaria 65 76  
 Miliary tuberculosis 176 177  
 Milium 47 278 279 324 326  
 Milium nodules 253  
 Milroy's disease 155

- Mixed carcinoma 379  
 tumor of the skin 360  
 Moll's glands 16  
 ..  
 Monocyte 34 35  
 Monocytic leukemia Naegeli type 474  
 495  
 Schilling type 474 475 490 495  
 Morphea 307  
 Mucin 28  
 fixation and staining for 28 29 30  
 in granuloma annulare 116  
 metastatic carcinoma 393  
 myxedema 280  
 myxoma 412  
 scleredema adultorum 281  
 Mucinosi papular 281  
 Mucoid degeneration of fatty tissue  
 299 303  
 Mucous membrane 11  
 Multiple benign cystic epithelioma  
 363 *See also* Trichoepthe-  
 loma  
 benign tumorlike growths of Schen-  
 inger and Buzzi 163  
 idiopathic hemorrhagic sarcoma of  
 Kaposi 431 435 437  
 myeloma 272  
 Munro micro abscesses 73 98 100 487  
 501  
 Muscles of the skin 25 438 439  
 Mycosis fungoides 33 71 75 105 157  
 153 204 305 306 330 435  
 471 473 474 484 490  
 d'emblee 484  
 involvement of internal organs 489  
 mycosis cell 485 486  
 poikiloderma atrophicum vasculare  
 in 484 489  
 Myelin sheaths of nerves 21  
 Myeloblast 32 474 491  
 Myelocyte 30 494  
 Myeloid leukemia 32 474 493  
 Myeloma multiple 272  
 Myelosis 32 470 493  
 Myoblastoma granular cell 334 440  
 Myoepithelial cells 14 17 350 353  
 355 358 359 361 363  
 of apocrine glands 17  
 Myoepithelial cells—(Continued)  
 eccrine glands 14  
 Myoepithelial sweat gland tumor 319  
 354 359 361 363  
 Myoepithelioma 319 354 359 361  
 363  
 clear cell 361  
 Myome dartoique 437  
 Myxedema 279  
 Myxoma 411  
 Myxomatous degeneration cyst of the  
 skin 413  
 Myxosarcoma 412  
 Nails 19  
 Necrobiosis lipoidica 31 116 257 269  
 Necrosis cavitating 175 212  
 Nerve end organs of dermis 21  
 Nerves fixation and staining 21 28  
 29 30  
 of dermis 21  
 epidermis 13  
 Neurinoma 418  
 Neurodermatitis circumscripta 72 *See*  
*also* Lichen simplex chronicus  
 disseminata 72 *See also* Atopic der-  
 matitis  
 Neurofibromatosis 38 411 413 416  
 417 457 458  
 Neurofibrosarcoma 420  
 Neuroid tubes 451 457  
 Neurolemoma 416 421  
 Neuroma 416  
 Neutrophils 32  
 Nevroid tumors 317 318 320 371  
 Nevroxantho-endothelioma 406 407  
 Nevus 320  
 araneus 427  
 blue 450 466  
 cell 450 452  
 histogenesis 450  
 definition of 320  
 epidermal 321  
 flammeus 422 423  
 hair 319  
 keratotic 321  
 linear 391  
 neural 458  
 organic 318 320  
 pigmentosus 320 392 450 458  
 compound 450 452 456

- Nevus—(Continued)  
   pigmentosus—(Continued)  
     intra-dermal type 450 452 456  
       459 461  
     junction type 342 450 452 453  
       458 459 460 461 465  
     lamellae foliaceous 451 457  
     neural type 458  
     neuroid tubes 451 457  
     et pilosus 458  
     verrucosus 458  
   sebaceous (Jadassohn) 319 343, 346  
     347 351  
   senile sebaceous 319 346  
   syngangliadenomatous papillif-  
     erous 319 349  
   unius lateralis 321 *See also* Nevus  
     verrucosus  
   vasculosus 422 423 425  
   verrucosus 41 282 283 317 321  
     346 385  
 Niemann Pick disease 257 265  
 Nodular subepidermal fibrosis 403  
   404  
   vasculitis 97  
 Nodules of Bizzozero 10  
 Nodulus cutaneus 403 *See also* Der-  
   matofibroma  
 Nonlipid histiocytosis of Foot and Ol-  
   cott, 262  
 North American blastomycosis 36 152  
   221 222 225 226 334  
 Nummular eczema 57 72  
  
 Ochronosis 286  
 Onychomycosis 218  
 Oriental sore (leishmaniasis) 235  
 Osler's disease 427  
 Ossification 364 412  
 Osteoma cutis 412  
 Ostium cystica Jungling (*sarcoidosis*)  
   191  
 Oxidase reaction 32 494  
  
 Pacini corpuscles 21 22  
 Pachyonychia congenita 42  
 Paget's disease 155 317 336 339  
   extramammary 339 342 389  
 Panniculitis relapsing febrile nodular  
   non-suppurative 33 131 135  
 Papilloma 219 317 321 322, 501  
 Papilloma—(Continued)  
   basal cell 322 323 328 382 385  
 Papillomatosis 501  
 Papular urticaria 138  
 Papulonecrotic tuberculid 376 383  
 Paraffinoma 36 140  
 Parakeratosis 11 501  
   variegata 104 105  
 Parapsoriasis 71 104 484  
   en plaques 104 105  
   guttata 100 104 105  
   lichenoides 101 105  
   varioliformis 101 105  
 Paravaccinia 253  
 Paschen bodies in variola 238 246  
 Pautrier micro-abscesses 72 485 487  
 Pellagra 71 283 287  
 Pemphigoid benign mucous mem-  
   brane 63 77 87  
   bullous 47 65 77 85 88 89  
 Pemphigus conjunctivae 63 77 87  
   erythematosus 65 77 84  
   familial benign chronic, 50 51, 65  
     77 80  
   foliaceus 65 77 82 84  
   Senear Usher type 65 77 84  
   vegetans 33 65 80 152  
   vulgaris 52 65 77 87 244  
 Perianteritis nodosa 96 150 310  
 Pericyte 24 431  
 Periderm 5  
 Perineurium 22  
 Periodic acid Schiff stain 8 30 73 99  
   218 224 225 230 231 272  
   279 297 299 501  
 Periductal cells 24 34 35 431 435  
 Peroxidase reaction 32 494  
 Phagocytosis 32 35  
 Phenolphthalein eruption 150  
 Phrynoderma 111 286  
 Pigment of the epidermis 12 *See also*  
   Melanin  
 Pigmentation due to gold 154  
   mercury 154  
   silver 153  
 Pigmented basal cell epithelioma 369  
   372 373  
   nevus 450 *See also* Nevus pigmen-  
     tosus  
   purpuric lichenoid dermatitis (Gou-  
     gerot and Blum) 129

- Mixed carcinoma 379  
tumor of the skin 360
- Moll's glands 16
- Molluscum contagiosum 238 244
- Mongolian spot 450 465
- Moniliasis 220 221
- Monocyte 31 35
- Monocytic leukemia Naegeli type 474 495  
Schulling type 474 475 490 495
- Morphea 307
- Mucin 28  
fixation and staining for 28 29 30  
in granuloma annulare 116  
metastatic carcinoma 393  
myxedema 280  
myxoma 412  
scleroderma adulatorum 281
- Mucinosis papular 281
- Mucoid degeneration of fatty tissue 299 303
- Mucous membrane 11
- Multiple benign cystic epithelioma 363 *See also* Trichoepithelioma  
benign tumorlike growths of Scheninger and Buzzi 163  
idiopathic hemorrhagic sarcoma of Kaposi 431 435 437  
myeloma 272
- Munro micro-abscesses 73 98 100 487 501
- Muscles of the skin 25 438 439
- Mycosis fungoides 33 71 75 105 137 153 204 305 306 350 435 471 473 474 484 490  
d'emblee 484  
involvement of internal organs 489  
mycosis cell 485 486  
poikiloderma atrophicans vasculare in 484 489
- Myelin sheaths of nerves 21
- Myeloblast 32 474 494
- Myelocyte 32 494
- Myeloid leukemia 32 474 493
- Myeloma multiple 272
- Myelosis 32 470 493
- Myoblastoma granular-cell 331 440
- Myo epithelial cells 14 17 350 353 355 358 359 361 363  
of apocrine glands 17
- Myo-epithelial cells—(Continued)  
eccrine glands 14
- Myo epithelial sweat gland tumor 319 354 359 361 363
- Myo epithelioma 319 351 359 361 363  
clear cell 361
- Myome dartoique 437
- Myxedema 279
- Myxoma 411
- Myxomatous degeneration cyst of the skin 413
- Myxosarcoma 412
- Nails 19
- Necrobiosis lipoidica 31 116 257 269
- Necrosis cicaeal 175 212
- Nerve end organs of dermis 21
- Nerve fixation and staining 21 28 29 30  
of dermis 21  
epidermis 13
- Neurinoma 418
- Neurodermatitis circumscripta 72 *See also* Lichen simplex chronicus  
disseminata 72 *See also* Atopic dermatitis
- Neurofibromatosis 38 411 413 416 417 457 458
- Neurofibrosarcoma 420
- Neuroid tubes 451 457
- Neurolemonia 416 421
- Neuroma 416
- Neutrophils 32
- Nevoid tumors 317 318 320 371
- Nevoxantho endothelioma 406 407
- Nevus 320  
araneus 427  
blue 450 466  
cell 450 452  
histogenesis 450  
definition of 320  
epidermal 321  
flammeus 420 423  
hair 319  
keratotic 321  
linear 321  
neural 458  
organic 318 320  
pigmentosus 320 322 450 458  
compound 450 452 456

- Reticulum—(Continued)  
   fibers 5 8 19 20 31 32 36  
   development into collagen 20 36 189  
   fixation and staining for 29  
   lymphoma 32 410 471 473 474 481 488 490 492  
   differentiation from reticuloendotheliosis 476  
   sarcoma 474 *See also* Reticulum-cell lymphoma  
 Rheumatic fever subcutaneous nodules of 117  
 Rheumatoid arthritis subcutaneous nodules of 116 117  
 Rhinoscleroma 37 202 232 236 237  
 Riehl's melanosis 283  
 Rosacea 124 183  
 Rosacea like tuberculid 124 176 182  
 Rothmund's syndrome 45  
 Round cell sarcoma 401 471  
 Ruffini bodies 22 23  
 Russell bodies 37 202  
  
 San Joaquin Valley fever (coccidiosis domyosia) 36 221 226  
 Sarcoid Spiegler Fendt type 137 188 480 491  
   subcutaneous of Darier Roussy 188 303  
 Sarcoidosis 36 141 142 179 183 188 196  
   asteroid inclusion bodies 190  
   differentiation from lupus vulgaris 179  
   erythoderma 189  
   osteitis cystica (Jungling) 191  
   Schumann's inclusion bodies 189  
   neoplasia (Heerfordt) 191  
 Sarcoma 41 123 333 403 471  
   multiple idiopathic hemorrhagic of Kaposi 431 432 437  
   reticulum-cell 403 471  
   round cell 403 471  
   spindle cell 403  
 Scabies 136  
 Schamberg's progressive pigmentary dermatosis 129  
 Schumann inclusion bodies 142 189  
 Schönlein-Henoch's purpura 32 33 89 127 130 311 313  
 Schüller-Christian disease 31 113 207 262 263 474  
 Schwannian cells 23 418 419 422 441 450 454  
   sheath 22 422  
 Schweninger-Buzzi type of macular atrophy 163  
 Scleredema adaliorum 281  
 Sclerema neonatorum generalized 133 nodular 133 134  
 Scleroderma 118 119 276 302 303 305 306 307 412  
   circumscribed 307  
   generalized 302 303 305 307 309  
   relationship to acute systemic lupus erythematosus 302  
   visceral lesions in 310  
 Sclerosing hemangioma 403 404 424  
   lipogranuloma 140  
 Scrofuloderma 176 177 180 188 215 227 334  
 Scurvy 127 288  
 Sebaceous adenoma 319 347 353 386  
   cyst 139 321 368 369  
   epithelioma 319 353 386  
   glands 4 17 21 318 315 341 346 347 353 377  
   carcinoma of 318 383  
   embryology of 4  
   generative cells 17 318  
   hyperplasia senile 316  
   nevus senile 313 346  
 Seborrheic dermatitis 67 68 73 100 101 263  
   keratosis 382 *See also* Basal-cell papilloma  
 Self-healing squamous cell carcinomata 336  
 Senear-Usher type of pemphigus 62 77 84  
 Senile degeneration of the skin 157 296  
   elastosis 56 157  
   hemangioma 420  
   keratosis 44 63 77 150 317 322 323 326 328 331 338 380  
   purpura 127 157  
   sebaceous hyperplasia 316  
   nevus 319 316  
   verruca 382 *See also* Basal cell papilloma

- Pityriasis lichenoides et varioliformis  
     acuta 101 105  
     rosea 71 105  
     rubra pilaris 111 287  
 Plasma cells 36  
 Poikiloderma atrophicum vasculare  
     (Jacobi) 15 305 481 489  
     in mycosis fungoides 484 489  
     reticulare (Civatte) 283  
 Poikilodermatomyositis 305  
 Polariscopes 31  
 Polariscopic examination 17 31 133  
     195 258 268 271 406  
 Polymorphonuclear leukocyte 32  
 Polysaccharides 30 31  
     staining for 30  
 Pompholyx 65 76  
 Porokeratosis Mibelli 13  
 Porphyrin 16 65 87 278  
 Port wine nevus 422  
 Post mastectomy lymphangiosarcoma  
     137  
 Precancerous dermatosis of Bowen 121  
     155 317 328 336 338 342  
 Primary epithelial germ 4 16 318  
     344 365 368 371 372 374  
     375  
     tumors of 318  
 Progeria of adults 45  
     children 45  
 Progressive pigmentary dermatosis of  
     Schamberg 129  
 Protozoa diseases caused by 235  
 Prurigo nodularis 95  
 Pseudoacanthosis nigricans 282  
 Pseudoepitheliomatous hyperplasia  
     50 95 157 151 170 179 180  
     200 202 222 232 334 336  
     380 441  
 Pseudopelade Brocq 126  
 Pseudoxanthoma elasticum 54 158  
 Psoriasis 34 71 73 98 208 211 484  
     487  
     exfoliative 98 101  
     pustular 101 102  
     of palms and soles 101 102  
 Purpura 126 150 313  
     angiothrombotic 32 33 89 127 130  
     311 313  
     annularis telangiectodes (Majocchi)  
         129  
     Purpura—(Continued)  
         due to drugs 127, 150  
         due to stasis 127  
         febrile with platelet thrombosis 127  
         inflammatory 127  
         in meningococcemia 127 171  
         periarteritis nodosa 127 311 313  
         scurvy 127 288  
         subacute bacterial endocarditis  
             127 170  
         noninflammatory 127, 288  
         pigmentosa progressiva (Majocchi  
             Schamberg) 129  
         Schoenlein Henoch 127  
         senile 127, 157  
         thrombocytopenic 127  
     Pustular bacterid 101 102  
     psoriasis 101, 102  
         of palms and soles 101 102  
     Pustule spongiform of Kogoj 102  
         103 104 172  
     Pustulosi palmaris et plantaris 101  
         102  
     Pyknosis of nucleus 175 186 501  
     Pyoderma gangrenosum 151 170 331  
     Pyodermite végétante 80 81  
  
     Quinacrine hydrochloride dermatitis  
         due to 150  
  
     Radiodermatitis 44 119 305 333 370  
         371 410  
     Reed Sternberg giant cells 75 481, 488  
         490  
     Refractility of fat 17 31 133 195 258  
         268 271 406  
     Reiter's disease 103 104 172 174  
     Relapsing febrile nodular nonsuppurative  
         pruriginous 33 131 135  
     Reticular cells 33 34 35 470 174  
         175 485  
         degeneration of epidermal cells 65  
             66 69 239 243 244 501  
     Reticuloendotheliosis lipid 257 262  
         nonlipid 262 263 476  
         differentiation from reticulum cell  
             lymphoma 476  
     Reticulo-histiocytoma 408  
     Reticulois lipomelanotic 75  
     Reticulum cell 21 31 35 37 470 472  
         471 485

## Reticulum—(Continued)

- fibers 5 8 19 20 34 35 36
- development into collagen 20 36 183
- fixation and staining for 29
- lymphoma 35 410 471 473 474 481 488 490 492
- differentiation from reticulo-endotheliosis 476
- sarcoma 474 See also Reticulum cell lymphoma
- Rheumatic fever subcutaneous nodules of 117
- Rheumatoid arthritis subcutaneous nodules of 116 117
- Rhinoscleroma 37 202 237 236 237
- Riehl's melanosis 283
- Rosacea 124 183
- Rosacea like tuberculid 124 176 182
- Rotl mund's syndrome 45
- Round cell sarcoma 401 471
- Ruffini bodies 22 23
- Russell bodies 37 202
- San Joaquin Valley fever (coccidiosis, stomycosis) 36 221 226
- Sarcoma Spiegler Fendt type 137 188 480 431
- subcutaneous of Darier Roussy 188 303
- Sarcoidosis 36 141 142 179 183 188 196
- asteroid inclusion bodies 190
- differentiation from lupus vulgaris 190
- erythroidemic 188
- ostitis cystica (Jungling) 191
- Schammann inclusion bodies 189
- uveoparitis (Heerfordt) 191
- Sarcoma 41 122 333 403 471
- multiple idiopathic hemorrhagic of Kaposi 431 435 437
- reticulum cell 403 471
- round cell 403 471
- spindle cell 403
- Scabies 136
- Schamberg's progressive pigmentary dermatosis 129
- Schammann inclusion bodies 142 189
- Schönlein-Henoch's purpura 32 33 81 127 130 311 313
- Schuller-Christian disease 31 113 257 262 265 474
- Schwannian cells 23 418 419 422 441 450 454
- sheath 22 422
- Schweninger-Buzzi type of macular atrophy 163
- Scleredema adultorum 281
- Sclerema neonatorum generalized 133 nodular 133 134
- Scleroderma 118 119 276 302 303 305 306 307 442
- circumscribed 307
- generalized 302 303 305 307 309
- relationship to acute systemic lupus erythematosus 302
- visceral lesions in 310
- Sclerosing hemangioma 403 404 424
- lipogranuloma 140
- Scrofuloderma 176 177 180 188 215 227 331
- Scurvy 127 288
- Sebaceous adenoma 319 347 353 386
- cyst 139 324 368 369
- epithelioma 319 333 386
- glands 4 17 21 318 343 344 346 347 353 377
- carcinoma of 318 385
- embryology of 4
- germinative cells 17 348
- hyperplasia senile 346
- nevus senile 319 346
- Seborrheic dermatitis 67 68 73 100 101 263
- keratosis 382 See also Basal cell papilloma
- Self healing squamous cell carcinomata 336
- Senear-Usher type of pemphigus 65 77 81
- Senile degeneration of the skin 157 296
- elastosis 56 157
- hemangioma 496
- keratosis 44 65 77 155 317 322 323 326 328 331 338 385
- purpura 127 157
- sebaceous hyperplasia 316
- nevus 319 346
- verruca 382 See also Basal-cell papilloma



- Silicon granuloma 140 142  
 Solar dermatitis 122  
 South American leishmaniasis 235  
 Spider nevus 427  
 Spiegler Fendt sarcoma 137 188 480 491  
 Spindle cell sarcoma 402  
 Spirochetes fixation and staining 28 29 30  
     in syphilis 208 209 211 215  
 Spongiform pustule of Kogoj 102 103 104 172 501  
 Spongiosis 500 501  
 Sporotrichosis 199 221 229  
 Squamous cell carcinoma 44 50 53 120 122 137 155 162 178 179 200 202 204 252 317 324 327 328 329 334 336 338 344 363 370 379 380 387 389 410 441 461 485  
     differentiation from basal cell epithelioma 380  
     pseudo epitheliomatous hyperplasia 334  
     due to arsenic 155  
     grading of 330  
     intra epidermal 336 *See also* Bowen's disease  
     multiple primary self healing 336  
     spindle celled 122 333 464  
 Squamous-cell layer 4 7 10  
 Staining methods 29 30  
     histochemical 30  
 Stasis dermatitis 67 68 73 130 334  
     purpura 127  
 Steatoma 325  
 Stellate abscesses of lymphopithia venerea 252  
 Stem cell lymphoid reticular 33 170 472 474  
     lymphoma 403 410 471 473 474 482  
 Sternberg Reed giant cell 75 181 188 490  
 Stratum basale 7  
     corneum 7 11  
     germinativum embryonal 3 4  
     granulosum 7 11  
     intermedium 3 41  
     lucidum 7 12  
     malpighii 4 7 10  
 Strawberry mark 422  
 Striae distensae 162  
 Subacute bacterial endocarditis 170  
 Subcutaneous fat necrosis of the newborn 133 134  
     nodules of rheumatic fever 117  
     rheumatoid arthritis 116 117  
     sarcoma of Darier Roussy 188  
 Succet Hoyer canal 24 427 429  
 Superficial basal cell epithelioma 1 369 372 374 405  
 Sweat glands adeno-acanthoma of apocrine 4 16 125 318 312 3 319 352 354 356 389  
     adenomas of 317 349 352  
     carcinoma of 318 312 389  
     embryology of 4  
     epitheliomas of 354  
     nevus of 319 346  
     eccrine 13 16 76 77 123  
     carcinoma of 318 386  
     embryology of 4  
 Swimming pool granuloma 142 180  
 Sycosis vulgaris 168 220  
 Synovial cyst of the skin 413  
 Syphilis 36 37 163 175 204 208 33  
     alopecia in 208 209  
     condyloma latum 208 209  
     congenital 209 215  
     differential diagnosis from tuberculosis 215  
     gummatous 181 188 208 213 216 331 380  
     internal lesions in 215  
     juxta articular nodes 208 215  
     primary 208 209  
         lymph nodes in 209  
     secondary 208 209 296  
     spirochetes in 208 209 211 215  
     tertiary 208 212  
 Syringocystadenoma papilliferum 319 349  
 Syringoma 319 351 367  
 Tactile cells of Merkel and Ranvier 13 21  
     corpuscles of Meissner 13 21 23 151 155  
 Telangiectasia familial hemorrhagic 427

- angiectatic carcinoma 390 392  
 anogenetic anhidrosis 76  
 aplastic purpura 127  
 ca, 217  
 urdae 217 219  
 pitus 217 219  
 aporis 217  
 uris 217  
 in color 218  
 ne, eos nophilia 33  
 onocyte 34  
 ishibilis 10  
 ulous 221 224 231  
 non giant cells 239 269 406 407  
 cho-epithelioma 319 356 357 358  
 363 379 382  
 relationship to basal-cell epithe-  
 lioma 364 379  
 berle 175  
 berulid 176 182  
 chen scrofulosorum 176 185  
 chenoid 185  
 apus miliaris disseminatus faciei  
 176 183  
 n cropapular 176 182  
 apulonecrotic 176 183  
 osacea like 124 176 182  
 berculosis 36 174 199 201 215  
 270 335  
 lassification 176  
 culis colligativa 177 180  
 culis officialis 176 177 181  
 d differentiation from syphilis 215  
 Chon complex 174 176  
 lupus vulgaris 142 176 177 180  
 185 190 215 334 380  
 miliary 176 177  
 primary 176  
 primary inoculation of the skin 174  
 176  
 reinfection 177  
 verrucosa cutis 142 176 177 180  
 224 226  
 tuberculous chancre 176  
 tuberos sclerotic 314  
 uliremia 199 231  
 umis epidermal 317  
 mesodermal 403  
 mixed of the skin 360  
 nevroid 317 318 320
- Tunica dartos 25  
 Turban tumors 357 See also Cylindroma  
 Tyrosinase 12  
 Tranch test 79
- Ulcer chronic undermining burrow-  
 ing 190  
 Urticaria 94 150  
 popular 138  
 pigmentosa 38 57  
 Uveoparotitis of Heerfordt (sarcoido-  
 sis) 191
- Vaccinia 241 243 244 251  
 virus 241 244  
 Varicella 65 238 239 243  
 Varicelliform eruption of Kaposi 244  
 Variola 65 238 239 242 246 254  
 Vasculitis allergic 32 89 127 171  
 311  
 nodular 97  
 Vegetative nerves of the skin 21  
 Verocay bodies (neurolemmas) 422  
 Verruca necrogenerica 180  
 plana 53 54 246 249  
 plantaris 246 249 251  
 senilis, 382. See also Basal cell papu-  
 loma  
 vulgaris 54 246 247 250 251 320  
 324
- Vesicle definition 501  
 Vesicles classification 65  
 mode of formation 64  
 Vili 49 50 51 52 80 350 502  
 Vichow cells of leprosy 194  
 Viruses cultivation of 238 246 252  
 diseases caused by 238  
 elementary bodies of 238  
 inclusion bodies 238 241 242 243  
 244 246 249 252 253
- Vitamins  
 A deficiency (phrynodermia) 111  
 286 288  
 B deficiency (pellagra) 71 283 287  
 C deficiency (scurvy) 127 288  
 Vitiligo 233  
 von Gierke's disease 261  
 von Recklinghausen's disease 38 411  
 413 416 417

Wandering connective tissue cell, 34, 35

Waterhouse Friderichsen syndrome, 171

Weber-Christian disease 33 131, 135

Wen, 324

Werner's syndrome, 45

Wire loop glomerulitis, 300, 310

Xanthelasma palpebrarum 31, 257, 260, 261, 269

Xanthoma, 257 *See also* Xanthomatosis

cells 35, 258, 260 406, 407

diabeticorum 261

disseminatum 263

eruptive, 261, 262

Xanthoma—(Continued)

planum et tuberosum, 257, 260 261 406 408

Xanthomatosis 31, 257, 274

biliary, 257, 260

Burger Grutz type, 261

normocholesteremic, 262

primary hypercholesteremic, 237, 260 261, 269 407, 408

secondary 257, 261, 262

Xanthomatous biliary cirrhosis 237 260

Xeroderma pigmentosum 44

X-ray dermatitis 44 119 303 333 370 371, 410

Zenker's solution 28



- Wandering connective tissue cell 34  
35
- Waterhouse Friderichsen syndrome  
171
- Weber-Christian disease 33 131 135
- Wen 324
- Werner's syndrome 45
- Wire loop glomerulitis 300 310
- Xanthelasma palpebrarum 31 257  
260 261 269
- Xanthoma 257 *See also* Xanthomato-  
sis  
cells 35 258 260 406 407  
diabeticorum 261  
disseminatum 263  
eruptive 261 262
- Xanthoma—(Continued)  
planum et tuberosum 257, 260 261  
406 408
- Xanthomatosis 31 257 274  
biliary 257 260  
Burger Gruz type 261  
normocholesteremic 262  
primary hypercholesteremic 257  
260 261 269 407 408  
secondary 257 261 262
- Xanthomatous biliary cirrhosis 257  
260
- Xeroderma pigmentosum 44
- Xyry dermatitis 44 119 305 333 370  
371 410
- Zenker's solution 28

